

P-ISSN2349-8528  
E-ISSN 2321-4902  
IJCS 2016; 4(1): 112-116  
© 2016 JEZS  
Received: 14-11-2015  
Accepted: 15-12-2015

**Abdu Z**  
Department of Chemical Sciences, Federal University Kashere P.M.B 0182 Gombe, Gombe State – Nigeria.

**Dimas K**  
Department of Chemistry, Modibbo Adama University of Technology Yola P.M.B 2076, Yola – Nigeria.

**Sunday AO**  
Department of Chemistry, Modibbo Adama University of Technology Yola P.M.B 2076, Yola – Nigeria.

**Isyaka MS**  
Department of Chemical Sciences, Federal University Kashere P.M.B 0182 Gombe, Gombe State – Nigeria.

**Sa'id J**  
Department of Chemical Sciences, Federal University Kashere P.M.B 0182 Gombe, Gombe State – Nigeria.

**Correspondence**  
**Abdu Z**  
Department of Chemical Sciences, Federal University Kashere P.M.B 0182 Gombe, Gombe State – Nigeria.

## Qualitative investigation of phytochemicals and brine shrimp lethality test of the root, stem bark and leaves extract of *Isoberlinia doka* (Fabaceae)

**Abdu Z, Dimas K, Sunday AO, Isyaka MS, Sa'id J**

### Abstract

*Isoberlinia doka* was selected in this study on the bases of its use in folk medicine by Lala tribe of Adamawa State - Nigeria to cure Jaundice and Headache diseases. Extracts prepared by soxhlet extraction with hot methanol from the plant were macerated and screened for the presence of secondary metabolites and activity in the brine shrimp (*Artemia salina* Leach) lethality test (BST). In the phytochemical screening, n-hexane extracts of the leaves, stem bark and root showed the presence of few secondary metabolites with the methanol extracts indicating the presence of most of the secondary metabolites (Tables 2-4). Extract of the root and stem bark exhibited high lethality on brine shrimp larvae with  $LC_{50} = 63.1\mu\text{g}/\text{ml}$  and  $70.8\mu\text{g}/\text{ml}$  respectively. The leaves extracts exhibited a moderate potent activity at  $LC_{50} = 436.5\mu\text{g}/\text{ml}$ . Maximum mortalities occurred at  $1000\mu\text{g}/\text{ml}$  concentration while the least mortalities happened to be at  $65.5\mu\text{g}/\text{ml}$  concentration. The mean results of mortality percentage of the brine shrimp versus the log of concentrations were plotted using the Microsoft Excel spreadsheet application, which also formulated the regression equations. These equations were later used to calculate  $LC_{50}$  values for the samples tested with consideration of value greater than  $1000\mu\text{g}/\text{ml}$ , suggesting that the extract is nontoxic.

**Keywords:** Phytochemicals, *Artemia salina*, Brine shrimp test, Folk medicine, *Isoberlinia doka*.

### 1. Introduction

The importance of plants is well known to us. Plant kingdom is a treasure house of potential drugs and in the recent years there has been an increasing awareness about the importance of medicinal plants. Drugs from the plants are easily available, less expensive, safe, efficient and rarely have side effects. According to WHO medicinal plants would be the best source to obtain variety of drugs. Large percentage of peoples from developed countries uses traditional medicines, which has compounds derived from medicinal plants. However, such plants should be investigated to better understand their properties, safety and efficiency (Athima *et al.*, 2005) [4]. Medicinal plants contain some natural products which perform definite physiological action on the human body and these bioactive substances include tannins, alkaloids, carbohydrates, terpenoids, steroids and flavonoids (Adoum, 2009) [2].

### Plant Collection and Identification

The fresh leaves, stem bark and root of *Isoberlinia doka* was collected in Kashere, Akko Local Government Area of Gombe State, Nigeria in May, 2015. The plant was identified by Alhaji Adamu of Kashere town and authenticated by Botanists in the Department of Biological sciences Federal University Kashere.

### Preparation of Plant Parts for Extraction

The plant samples were air dried and ground into powder with pestle and mortar and weighed. Around 60g of the powder was packed in a thimble of filter paper prepared manually. The thimble was then inserted into the Soxhlet apparatus, 500ml methanol was transferred down the thimble into the pot. A temperature of  $75^{\circ}\text{C}$  was maintained and extraction continued for 6 hours. Then the methanol extract was collected and the cake recovered from the thimble was kept. The methanolic extract was evaporated on a rotary evaporator (R110) at  $40^{\circ}\text{C}$ . Altogether, 200g of each sample were extracted and labeled F01. The crude extract F01 of the leaves, stem bark and root of *Isoberlinia doka* was macerated using solvents in ascending order of polarity (n-hexane, chloroform, ethylacetate, and methanol). For each solvent 20ml was

used to macerate the crude extract 5 times. The n-hexane soluble was collected, labeled as F02 and allowed to evaporate to dryness at 38 °C. The chloroform, ethylacetate, and methanol soluble were in the same way collected and labeled as F03, F04, F05 and allowed to evaporate to dryness at 38 °C (Omar, 2011).

#### **Phytochemical Screening of Plant Material**

All the plant extracts of the leaves, stem bark and root of *Isoberlinia doka* were screened for the presence of alkaloids, saponins, flavonoids, sterols, anthraquinones, phlobanins tannins, etc. using modified procedure described by Aduragbenro *et al.* 2009 [3].

#### **Test for tannins.**

A small quantity of the extract was mixed with distilled water and heated on a water-bath. The mixture was filtered and a solution of H<sub>2</sub>SO<sub>4</sub> followed by few drops of 5% FeCl<sub>3</sub> solution are added to the filtrate. A dark green or dark blue precipitate indicates presence of tannins.

#### **Test for glycosides (general test)**

Solution of H<sub>2</sub>SO<sub>4</sub> (10ml 50%) was added to 1mL of extract in a test tube, this mixture heated in boiling water for 5 minutes. 10mL Fehling's solution A, and B (5 mL each) was added and boiled. Brick red precipitate indicates positive test.

#### **Bornträger's Test for Anthraquinone Glycoside**

Small portion of the extract was mixed with dilute hydrochloric acid (HCl), extracted with carbon tetrachloride (CCl<sub>4</sub>) and then shaken with dilute ammonia (NH<sub>3</sub>). A rose pink to cherry red color indicates the presence of anthraquinone glycoside.

#### **Keller-Killiani Test for Cardiac Glycoside**

Small portion of the extract was mixed with 3ml of 3.5% solution of ferric chloride (FeCl<sub>3</sub>) in glacial acetic acid and transferred to the surface of 2ml conc. H<sub>2</sub>SO<sub>4</sub>. A reddish-brown color at the inter-phase with the upper layer pale-green in color which gradually turn blue was an indication of a deoxy sugar.

#### **Test for Cyanophoric Glycoside**

A portion (1ml) of the extract was placed in a test tube, a piece of sodium picrate paper was then suspended above the level of the extract by trapping the edge of the paper between the cork fitting and the wall of the test tube. The set-up was allowed to stand for 30 minutes in a warm water (45 °C). A change in color of the picrate paper from yellow to brick-red indicates the presence of cyanophoric glycoside.

#### **Test for resins**

Copper (II) Sulphate solution (2.5ml) was added to 2.5 mL of the extract. The resulting solution was shaken vigorously and allowed to separate. A green colour indicates positive test.

#### **Test for saponins (Frothing test)**

The extract solution (2ml) was vigorously shaken in test tube for two minutes followed by heating in a water-bath to boiling point. Frothing indicates positive test.

#### **Test for phlobatannins**

Distilled water (5cm<sup>3</sup>) was added to 5cm<sup>3</sup> of extract solution and boiled with 1% HCl for two minutes. A deep green colour indicates positive test.

#### **Test for flavonoids**

Extract solution (2ml) was heated with 10 ml of ethyl acetate on a water bath and allowed to cool for the layers to separate. A layer of red coloration (ammonia) indicates positive test.

#### **Salkowski Test for Steroidal Nucleus**

A small portion of the extract (0.5ml) was dissolved in 2ml CHCl<sub>3</sub>, followed by 2ml dilute solution of H<sub>2</sub>SO<sub>4</sub>. A reddish-brown color at the interphase indicates presence of steroid ring or nucleus.

#### **Liebermann-Burchard Test for Steroidal Nucleus**

A small portion of the extract was dissolved in CHCl<sub>3</sub> and filtered. The filtrate was mixed with 2ml acetic anhydride and 2 drops of conc. H<sub>2</sub>SO<sub>4</sub>. A color change from violet to blue or bluish-green indicates the presence of steroid nucleus or ring.

#### **Test for Phenols.**

Equal volumes of extract solution and FeCl<sub>3</sub> were mixed and allowed to stand for while. A deep bluish green solution confirms the presence of phenols.

#### **Test for volatile oils.**

The extract (0.2g) was mixed with 90% ethanol and 3 drops of ferric chloride was added. The appearance of green coloration confirms the presence of volatile oils.

#### **Test for carbohydrate. (Reducing sugars) [Fehling test]**

Mixtures of equal volumes (5ml) Fehling solution A and B was added to 2 ml of the extract in a test tube. The resultant mixture was boiled for two minutes. A brick red precipitate of copper oxide indicates a positive test.

#### **Test for alkaloids**

Concentrated solution of H<sub>2</sub>SO<sub>4</sub> (1ml) was added to 3ml of the extract and then treated with few drops of Wagner reagent. Reddish brown precipitate indicates positive test.

#### **Radulescu Test for Morphine Alkaloid**

1ml of the extract was evaporated to dryness and the residue was dissolved in a 0.6ml of 1% H<sub>2</sub>SO<sub>4</sub>, 2ml of distilled water followed by 2 drops of 10% NaNO<sub>3</sub> were added. The solution was made alkaline with dilute ammonia solution. A reddish-brown solution indicates the presence of morphine alkaloid.

#### **Thallequine Test for Quinoline Alkaloid**

To a portion of the extract was added 0.5ml of conc. HCl and a few crystals of KClO<sub>3</sub>, it was carefully evaporated and a drop of strong ammonia was added to it. A green coloration indicates a positive test.

#### **Test for Indole Alkaloid**

A few drops of conc. H<sub>2</sub>SO<sub>4</sub> and few crystals of K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> were added to a small portion of the extract. A coloration is an indication of presence of an Indole alkaloid.

#### **Vitalin-Morin Test for Tropane Alkaloid**

A small portion of the dry sample of the extract was dissolved in few drops of fuming HNO<sub>3</sub>, evaporated and allowed to cool. 2ml of acetone and a few drops of (3%) fresh alcoholic potash solution were added. A violet color indicates positive test.

#### **Test for Terpenoid (Salkowski test)**

The extract sample 0.2g was mixed with 2ml chloroform (CHCl<sub>3</sub>) and carefully followed by conc. H<sub>2</sub>SO<sub>4</sub> (3ml) to form a layer. A reddish brown coloration of the interface indicates positive test for the presence of terpenoids.

Preparation of test samples:

Samples were prepared by dissolving 2.0mg of the methanolic plant extract in 5 ml of a suitable solvent (Stock solution).

Dilution of this stock solution gives the series of concentrations required for testing. Three concentrations (3 replicates) were obtained for each series of tests. The negative and positive control solutions were simply distilled water and methanol respectively without test sample (plant extracts).

### Screening of Plants Extract in the Brine Shrimp Lethality Test

Brine shrimp eggs (*Artemia salina* premium grade) was generously provided by Professor O. A. Adoum of the Department of Pure and Industrial Chemistry Bayero University Kano. All the plants fractions were phytochemically screened for carbohydrates, tannins, saponin, anthraquinones, cardiac and cyanogenic glycosides, flavonoids, alkaloids etc using modified procedures outlined by Adoum *et al.* 2010.

*Artemia salina* eggs was added into a hatching chamber  $\frac{3}{4}$  filled with ocean sea water. The chamber was kept in an open space for 24hours, after which the eggs hatched into shrimp larvae. 4ml ocean water was then added and ten (10) larvae of *Artemia salina* were introduced into each vial. After 24 hours of introducing larvae, the number of survivals were counted in triplicate and recorded. To each sample vial, a drop of DMSO solvent was added, ten shrimps were transferred using a Pasteur pipette, and artificial seawater was added to make a total volume of 5 ml. The nauplii were counted against a lighted background. Counting for the chronic LC<sub>50</sub> began 24 hour after initiation of tests. Nauplii were considered dead if they were lying immobile at the bottom of the vials, and the percentage of deaths at each dose and at the control were determined (Omar 2011).

Microsoft Excel spreadsheet application was used to formulate the regression equations from the data of mean results of

percentage mortality of the brine shrimp versus the log of concentrations. These equations were later used to calculate LC<sub>50</sub> values for the samples tested with consideration of value greater than 1000  $\mu\text{g}/\text{ml}$ , suggesting that the extract is nontoxic (Abdul Rani *et al.*, 2010) [1].

## 2. Results and Discussion

**Table 1:** BST Assay Results of MeOH extract of *Isoberlinia doka* Leaves

Conc. ( $\mu\text{g}/\text{ml}$ )	Survivals			Deaths			% mortality	Log <sub>10</sub> Conc
	V	V	V	V	V	V		
1000	3	2	2	7	8	8	76.67	3
500	5	1	8	5	9	2	53.33	2.7
250	8	7	8	2	3	2	26.67	2.4
125	9	10	6	1	0	4	16.67	2.1
62.5	10	10	10	0	0	0	0.00	1.8
Ctrl(+)	0	0	0	10	10	10	100	
Ctrl(-)	10	10	10	0	0	0	0.00	

LC<sub>50</sub>( $\mu\text{g}/\text{ml}$ )  
436.5


**Table 2:** BST Assay Results of MeOH extract of *Isoberlinia doka* Stem Bark

Conc. ( $\mu\text{g}/\text{ml}$ )	Survivals			Deaths			% mortality	Log <sub>10</sub> Conc
	V	V	V	V	V	V		
1000	3	0	1	7	10	9	86.67	3
500	4	6	0	6	4	10	66.67	2.7
250	4	3	4	6	7	6	63.33	2.4
125	4	5	4	6	5	6	56.67	2.1
62.5	6	4	5	4	6	5	50.00	1.8
Ctrl(+)	0	0	0	10	10	10	100	
Ctrl(-)	10	10	10	0	0	0	0.00	



LC<sub>50</sub>( $\mu\text{g}/\text{ml}$ )  
70.8

**Table 3:** BST Assay Results of MeOH extract of *Isoberlinia doka* Roots

Conc. ( $\mu\text{g}/\text{ml}$ )	Survivals			Deaths			% mortality	Log <sub>10</sub> Conc
	V	V	V	V	V	V		
1000	0	3	2	10	7	8	83.33	3
500	2	4	2	8	6	8	73.33	2.7
250	3	3	3	7	7	7	70.00	2.4
125	2	4	5	8	6	5	63.33	2.1
62.5	5	8	4	5	2	6	43.33	1.8
Ctrl(+)	0	0	0	10	10	10	100	
Ctrl(-)	10	10	10	0	0	0	0.00	



**Fig 1:** graph of % mortality versus log<sub>10</sub> Conc of Stem bark

**Fig 2:** graph of % mortality versus log<sub>10</sub> Conc. for Root extract**Fig 3:** graph of % mortality versus log<sub>10</sub> Conc. for Leaves extracts

In the phytochemical screening, n-hexane extracts of the leaves, stem bark and root showed the presence of few secondary metabolites with the methanol extracts indicating the presence of most of the secondary metabolites.

Brine shrimp LC<sub>50</sub> values for the plant extracts evaluated in this study are reported in tables 1-3. All the extracts were found cytotoxic (LC<sub>50</sub> < 1000 µg/ml) in the brine shrimp bioassay. Based on the percentage of the mortality, the concentration that led 50% lethality (LC<sub>50</sub>) to the nauplii was determined by using the graph of mean percentage mortality versus the log of concentration.

### 3. Statistical Analysis

The mean results of mortality percentage of the brine shrimp versus the log of concentrations were plotted using the Microsoft Excel spreadsheet application, which also formulated the regression equations. These equations were later used to calculate LC<sub>50</sub> values for the samples tested with consideration of value greater than 1000 µg/ml, suggesting that the extract is nontoxic.

### 4. Conclusions

Our results indicated that methanol extracts of the root, stem bark, and leaves of *Isoberlinia doka* are active in the Brine Shrimp Assay. The results of the current study showed that activity of the extracts increases downwards from the leaves, stem bark and to the roots. The results does not concur with the use of this plant by traditional healers especially at higher doses to cure Jaundice and headache. A World Health Organization survey indicated that about 70-80% of the world's population rely on non-conventional medicine, mainly from herbal sources, in their primary healthcare, hence *Isoberlinia doka* may be used as a medicinal agent only in low dosages, especially in rural communities where conventional drugs are unaffordable because of the high cost. The reported active (cytotoxic) extracts in this study are worth of further pharmacological and phytochemical studies in order to define what kind of antitumor and antimalarial activity they have (if any) and to isolate the natural active constituents, which are responsible for the activity. Studies of this type are needed before a phytotherapeutic agent can be generally recommended for pharmaceutical use.

## 5. References

1. Abdul Rani, Muhamad Syahmi, Soundararajan Vijayarathna, Sreenivasan Sasidharan, Lachimanan Yoga Latha, Yuet Ping Kwan, Yee Ling Lau, Lai Ngit Shin, Yeng Chen. Acute Oral Toxicity and Brine Shrimp Lethality of *Elaeis guineensis* Jacq, (Oil Palm Leaf) Methanol Extract *Molecules*, 2010; 15:8111-8121.
2. Adoum OA. Determination of Toxicity Levels of Some Savannah Plants Using Brine Shrimp Test (BST) Bayero Journal of Pure and Applied Sciences. 2009; 2(1):135-138.
3. Aduragbenro DA, Yeside OO, Adeolu AA, Olanirewaju MJ, Ayotunde SA, Olumayokun AO *et al.* Blood Pressure Lowering Effect of *Adenanthera pavonina* seed extract on normotensive rats, Rec Nat Prod. 2009; 1(3):282-289.
4. Athima Saetung, Arunporn Itharat, Chawaboon Dechsukum, Chatchai Wattanapiromsakul, Niwat Keawpradub, Pranee Ratanasawan. Cytotoxic activity of Thai medicinal plants for cancer treatment J Sci Technol. 2005; 27(2):469-478.
5. Khaled A. Tawaha: Cytotoxicity Evaluation of Jordanian Wild Plants using Brine Shrimp Lethality Test Jordan J. J. Appl. Sci., 2006; 8(1):12-17
6. Oumar Al-Moubarak Adoum. Screening of *Cassia Singuaena*, *Commiphora Kerstingii*, *Khaya Senegalensis* for Brine Shrimp Lethality and Antiplasmodium Activity Chemsearch Journal. 2011; 2(2):50-52.
7. Syed M. Salman, Shaukat Ali, Ashfaq Ahmed, M Siddique Afridi, Amin Ur Rehman. Preliminary Phytochemical, Essential Element Analysis and Antimicrobial Activities of Ethanolic Extract of *Lotus Corniculatus* International Journal of Biosciences. 2015; 7(2):106-115.