



P-ISSN: 2349-8528

E-ISSN: 2321-4902

IJCS 2019; SP6: 272-274

Jai Prakash Gupta

Department of Agronomy,
Narendra Dev University of
Agriculture and Technology,
Kumarganj, Ayodhya, Uttar
Pradesh, India

Rajesh Kumar

Department of Horticulture,
COA, Azamgarh, Narendra Dev
University of Agriculture and
Technology, Kumarganj
Ayodhya Uttar Pradesh, India

Vimlesh Kumar

Department of Horticulture,
COA, Azamgarh, Narendra Dev
University of Agriculture and
Technology, Kumarganj
Ayodhya, Uttar Pradesh, India

(Special Issue -6)
3rd National Conference
On

**PROMOTING & REINVIGORATING AGRI-HORTI,
TECHNOLOGICAL INNOVATIONS
[PRAGATI-2019]
(14-15 December, 2019)**

Effect of nitrogen management and plant growth regulators on yield and yield attributes of wheat (*Triticum aestivum* L.)

Jai Prakash Gupta, Rajesh Kumar and Vimlesh Kumar

Abstract

The present investigation was conducted with the aim to know the effect of nitrogen management through organic and in-organic sources and plant growth regulators on the yield and yield attributes of the wheat cultivar HD-2968. The present study was conducted in the split plot design the treatments consist of four nutrient management, absolute control, RDF (Recommended dose of fertilizer) 150:60:40, 150% RDF.(225:90:60), 150% RDF+15 t FYM ha⁻¹ in main plots and four plant growth regulators, control- two sprays of water (400 lit water ha⁻¹) at first node and flag leaf stages. two spray of Chloromequat chloride (CCC) (Lihocin @ 0.2% (800 ml ha⁻¹) of commercial product dose at first node (Around 45 DAS) and flag leaf (Around 80 DAS) using 400 lit water ha⁻¹. two sprays of Tebuconazole (Folicur 430 SC) @ 0.1% at (400 ml ha⁻¹) First Node and Flag leaf with 400 lit water ha⁻¹. two spray combined application of Lihocin + Folicur in sub-plots.

Keywords: Wheat, nitrogen, yield, yield attributes and plant growth regulators

Introduction

Wheat (*Triticum aestivum* L.) belongs to family Poaceae, is the first important and strategic cereal crop for the majority of world population. It is the most important staple food of about two billion people (36.00% of the world population) worldwide. It provides nearly 55% of the carbohydrates and 20.00% of the food calories consumed globally and straw is a good source of feed for a large population of cattle in our country. It exceeds in acreage and production every other grain crop (including rice, maize, etc) and therefore, the most important cereal grain crop of the world, which is cultivated over a wide range of climatic conditions.

Among the different management practices, role of macro nutrients is crucial in crop nutrition for achieving higher yields (Raun & Johnson, 1999) [10]. The soils of India are deficient in nitrogen and are supplemented with chemical fertilizer for enhancing the crop productivity. Nitrogenous fertilizers play a vital role in modern farm technology, however only 20-50% of the soil applied nitrogen is recovered by the annual crops (Bajwa, 1992) [2]. The leftover nitrogen is lost from the soil system through denitrification, volatilization and leaching.

Efficient nitrogen (N) fertilization is crucial for economic wheat production and the protection of ground and surface waters (Alley *et al.* 1999) [1]. Nitrogen fertilizer rate and timing are the major tools available after planting for manipulating wheat growth and development to produce a greater grain yield per unit area (Simians 1982[11]. Alley *et al.* 1999) [1]. and such as intensive management systems is to increase N fertilizer rates and control lodging with PGRs ultimately to increase grain yields (Knapp and Harms, 1988 [9]. Van Sanford *et al.* 1989 [14]. Tripathi *et al.* 2003) [13].

Corresponding Author:**Jai Prakash Gupta**

Department of Agronomy
Narendra Dev University of
Agriculture and Technology,
Kumarganj Ayodhya, Uttar
Pradesh, India

The qualities of grain yield are influenced by fertilization treatment, environmental factors and by the genetic predispositions of a particular variety. It is commonly known that fertilization, especially nitrogen fertilization, has a fundamental position in the attainment of high yields and high quality grain. Nitrogen positively influences the leaf area and its chlorophyll concentration, thereby inducing crude protein content and the rheological properties of dough (Blandino and Reznari, 2009) [15]. and several field experiments showed that in winter wheat the number of spikes per unit area generally increases as the N rate increases while mean kernel weight usually declines (Batey and Reynish, 1976; Knapp and Harms, 1988 [9]. Alley *et al.* 1999) [1].

Materials and methods

The present study was conducted in the split plot design the treatments consist of four nutrient management, absolute control, RDF (Recommended dose of fertilizer) 150:60:40, 150% RDF.(225:90:60), 150% RDF+15 t FYM ha⁻¹ in main plots and four plant growth regulators, control- two sprays of water (400 lit water ha⁻¹) at first node and flag leaf stages. two spray of Chloromequat chloride (CCC) (Lihocin @ 0.2% (800 ml ha⁻¹) of commercial product dose at First node (Around 45 DAS) and flag leaf (Around 80 DAS) using 400 lit water ha⁻¹.two sprays of Tebuconazole (Folicur 430 SC) @ 0.1% at (400 ml ha⁻¹) First Node and Flag leaf with 400 lit water ha⁻¹. two spray combined application of Lihocin + Folicur in sub-plots. The treatments were replicated thrice.

Results and Discussion

Yield and yield attributes are the function of vegetative growth and development. Different nutrient levels influenced significantly yield and yield component during both the years of investigation. The results clearly indicated that ear head m², length of ear, number of grains/ear head, number of spikelet's per spike, test weight increased with increase in nutrient levels up to 150% RDF + 15 t FYM ha⁻¹. This may be attributed to favourable effects of nitrogen, throughout crop growth period on plant growth and development. As under adequate supply of nitrogen, plant height, tillers m⁻², leaf area index and dry matter accumulation were highest which results in improved yield attributes due to improved photosynthetic activity of leaves.

The increase in yield attributes was due to better nutrient translocation during reproductive and grain filling stages and increasing rate of photosynthesis. The ear formation is

directly related to number of tillers. Hence conditions that favour the formation of more no. of tillers were also responsible for the production of higher no. of ears. The number of grains determined primarily by the amount of nutrients absorbed and secondary by the amount of carbohydrates produced at the time of spikelet's differentiation. Hence under present study increase in number of grains might be due to better assimilation of carbohydrates in ear. Similar findings also were reported by Yadav *et al.* (2005) [15]. Chaturvedi *et al.* (2006) [6]. and Khan *et al.* (2008) [8].

Grain, straw and biomass yield increased significantly with increasing nitrogen levels. The maximum grain, straw and biomass yield was recorded with the application of 150% RDF + 15 t FYM ha⁻¹ during both the years. The increased grain, straw and biomass yield might be due to the improvement in growth and yield attributing characters and higher photosynthetic activity. The observations was in conformity with the findings of Sobh *et al.* (2000), Mosalem *et al.* (2002), Barthwale *et al.*(2013) and Singh *et al.* (2017).

It is clear from the result presented in the previous chapter that application of growth regulator with nutrients significantly influenced yield attributes *viz.*, ear-head m², length of ear head, number of grains/ ear head, number of spikelet's spike⁻¹, test weight recorded maximum under the two spray of Lihocin + Folicur during both the years. The minimum in values of yield attributes was recorded in the absolute control during both the years. These results can be discussed in light of fact that when plant growth especially height was restricted by the growth regulators. Similarly, crop yields were higher in treatments having better development of yield attributes than those with poor yield attributes. Zhang *et al.* (2017) [16]. reported that application of PGRs significantly reduced grain yield, and the combination of the two PGRs (Manipulator and Palisade) had a synergistic effect and lowered the yield.

The maximum grain, straw and biomass yield was recorded with the application of two spray of Lihocin + Folicur and minimum in control. This might be due to the growth of plants and the plant height was reduced due to the application of Lihocin + Folicur which leads to the conversion of photosynthets towards the reproductive parts of the plants. The earlier researcher Gurmani *et al.* (2006) [7]. reported that higher grain yield and 1000- grain weight was recorded by IR-6.

Table 1: Effect of nutrient management and plant growth regulators on yield attributes

Treatments	Earhead/m ²		Length of ear head		No. of grains/ ear head		No. of spikelet's/spike		Test weight	
	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18
(A) Nitrogen levels (kg ha⁻¹)										
1. Absolute control.	180.83	169.17	9.70	9.43	38.50	37.14	17.32	16.83	33.74	35.92
2. RDF (150:60:40)	293.75	312.58	11.10	9.96	45.50	39.80	19.82	17.79	35.53	38.16
3. 150% RDF.(225:90:60)	315.00	335.50	11.70	10.21	48.50	41.07	20.89	18.24	36.82	38.50
4. 150% RDF+15 t FYM ha ⁻¹ .	360.00	394.58	12.05	12.28	50.25	51.42	21.52	21.93	38.71	40.73
S.Em±	5.563	4.374	0.195	0.237	0.973	1.183	0.347	0.422	0.618	0.570
CD (P=0.05)	19.250	15.137	0.673	0.818	3.366	4.092	1.202	1.462	2.139	1.973
(B) Plant growth regulatores										
1. Control- two spray of water	247.50	280.58	10.60	9.75	43.00	38.75	18.93	17.41	35.25	37.09
2. Two spray of Lihocin @ .2%	299.58	309.50	11.35	10.91	46.75	44.57	20.27	19.49	36.78	38.92
3. Two sprays of Folicur @ .1%	285.00	299.00	11.00	10.15	45.00	40.77	19.64	18.13	35.70	36.82
4. Two spray of Lihocin + licur.	317.50	322.75	11.60	11.07	48.00	45.33	20.71	19.76	37.08	40.48
S.Em±	4.787	3.389	0.173	0.218	0.866	1.052	0.309	0.390	0.553	0.518
CD (P=0.05)	13.973	9.890	0.506	0.637	2.528	3.186	0.903	1.138	1.616	1.511

Table 2: Effect of nutrient management and plant growth regulators on Grain Yield, Straw Yield, (qha⁻¹), Harvest index (%) and Biomass

Treatments	Grain Yield (qha ⁻¹)		Straw Yield (qha ⁻¹)		Harvest index (%)		Biomass (q ha ⁻¹)	
	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18
(A) Nitrogen levels (kg ha⁻¹)								
1. Absolute control.	17.02	11.45	23.86	14.91	41.79	43.45	40.87	26.36
2. RDF (150:60:40)	47.66	47.64	69.13	65.70	40.89	42.13	116.79	113.34
3. 150% RDF.(225:90:60)	51.85	51.36	77.79	77.48	40.15	39.85	129.64	128.84
4. 150% RDF+15 t FYM ha ⁻¹ .	53.96	62.43	83.63	87.10	39.20	41.74	137.58	149.53
S.EM±	1.070	1.023	2.747	1.488	1.903	0.549	2.089	2.153
CD (P=0.05)	3.702	3.540	9.506	5.150	6.584	1.899	7.230	7.451
(B) Plant growth regulatores								
1. Control- two spray of water	38.32	41.00	57.20	58.73	40.55	41.54	95.51	99.74
2. Two spray of Lihocin @ 0.2%	44.84	43.98	66.91	62.53	40.49	41.71	111.75	106.51
3. Two sprays of Folicur+ @ 0.1%	40.11	42.46	59.86	59.33	40.48	42.14	99.97	101.80
4. Two spray of Lihocin + Folicur.	47.22	45.45	70.43	64.59	40.52	41.78	117.65	110.04
S.EM±	0.800	0.861	1.699	1.399	0.800	0.561	1.704	1.879
CD (P=0.05)	2.334	2.512	4.960	4.084	2.334	1.639	4.973	5.484

References

- Alley MM, Brann DE, Hammons JL, Brethren WE. Nitrogen Management for Winter Wheat: Principles and Recommendations. Crop and Soil Environmental Sci. 1999; (1):424-429.
- Bajwa MI. Soil fertility management for sustainable agriculture. In: Proceeding 3rd National Congress of Soil Science, Lahore, Pakistan. March. 1992; 20-22. 1990, 7-25.
- Barthwal A, Bhardwaj AK, Chaturvedi S, Pandiaraj T. Site specific NPK recommendation in wheat (*Triticum aestivum* L.) for sustained crop and soil productivity in *moll sols* of Tarai region. Indian Journal of Agronomy. 2011; 58(2):208-214.
- Batey T, Reynish DJ. The influence of Nitrogen Fertilizer on Grain Quality in Winter Wheat. J Sci. Food Agric. 1976; (27):983-990.
- Blandino M, Reznerim A. Effect of fungicide and foliar fertilizer application to winter wheat at a thesis on flag leaf senescence, grain yield, flour bread-making quality and DON contamination. Europe. J Agro. 2009; (30):275-282.
- Chaturvedi I. Effects of different nitrogen levels on growth, yield and nutrient uptake of wheat (*Triticum aestivum* L.) Internat. J. agric. Sci. 2006; 2(2):372-374.
- Gurmani AR, Bano A, Muhammad S. Effect of growth regulators on growth, yield and ions accumulation of rice (*Oryza sativa* L.) under salt stress. Pak. J Bot. 2006; 38(5):1415-1424.
- Khan P, Imtiaz MAslam M, Shah HK, Syed NY M, Siddiqui Saleem-ul-Haq. Effect of different nitrogen and phosphorus ratios on the performance of wheat cultivar chairman. Sarhad J Agric. 2008; 24(2):12-14.
- Knapp JS, Harms CL. Nitrogen fertilization and plant growth regulator effects on yield and quality of four wheat cultivars. J Prod. Agric. 1988; (1):94-98.
- Raun WR, Johnson GV. Improving nitrogen use efficiency for cereal production, 1999.
- Simons RG. Tiller and Ear Production of Winter Wheat. Field Crop Abst. 1982; (35):857-870.
- Singh VK, Brahma SD, Mishra RP, Shukla AK, Timsina J, Upadhyay PK *et al.* Yields, Soil Health and Farm Profits under a Rice-Wheat System: Long-Term Effect of Fertilizers and Organic Manures Applied Alone and in Combination. Agronomy. 2018; 9(1):1-22.
- Tripathi SB, Hazra CR. Sulphur requirement of groundnut (*Arachis hypogea*) under wheat (*Triticum aestivum* L.) based cropping system. Indian J Agri. Sci. 2003; 73(7):368-372.
- Van Sanford DA, Grove JH, Grabau LJ, MacKown CT. Ethephon and Nitrogen Use in Winter Wheat. Agron. J. 1989; (81):951-954.
- Yadav MP, Mohd Adlam, Kushwaha SP. Effect of integrated nutrient management in rice (*Oryza sativa*) wheat (*Triticum aestivum* L.) cropping system in Central Plain Zone of Uttar Pradesh. Indian Journal of Agronomy. 2005; 50(2):89-93.
- Zhang Y, Su S, Tabori M, Yu J, Chabot D, Baninasab B. *et al.* Effect of Selected Plant Growth Regulators on Yield and Stem Height of Spring Wheat in Ontario. Journal of Agricultural Science. 2017; 9(12):30-42.