



P-ISSN: 2349-8528

E-ISSN: 2321-4902

[www.chemijournal.com](http://www.chemijournal.com)

IJCS 2020; SP-8(4): 95-99

© 2020 IJCS

Received: 09-06-2020

Accepted: 11-05-2020

**MP Yeshika**

Department of Sericulture, UAS,  
GKVK, Bangalore, Karnataka,  
India

**KG Banuprakash**

Department of Sericulture, UAS,  
GKVK, Bangalore, Karnataka,  
India

**K Murali Mohan**

Department of Agricultural  
Entomology, UAS, GKVK,  
Bangalore, Karnataka, India

**KS Vinoda**

Department of Silkworm  
Genetics & Breeding, College of  
Sericulture, Chintamani,  
Karnataka, India

**Corresponding Author:**

**MP Yeshika**

Department of Sericulture, UAS,  
GKVK, Bangalore, Karnataka,  
India

## Effect of novel insecticide molecules in mulberry on reeling parameters of silkworm *Bombyx mori* L. cocoons

**MP Yeshika, KG Banuprakash, K Murali Mohan and KS Vinoda**

**DOI:** <https://doi.org/10.22271/chemi.2020.v8.i4b.9872>

### Abstract

Effect of novel insecticides treated mulberry leaves on the reeling performance of silkworm *Bombyx mori* L. were studied. The cocoon filament length was longest in untreated control and in the insecticide treatments viz., dinotefuron 20 SG @ 0.25 g/l, dichlorvos 76 EC @ 2.63 ml/l at different time intervals of 10, 20, 30 Days After Spraying. Lowest filament length was recorded when flonicamid 50WG sprayed @ 0.3 g/l and 20, 30 and 40 DAS. This was followed by azadirachtin 1 % @ 1 ml/l recording shorter filament length at 40 DAS. However, at 10 and 20 DAS azadirachtin 1 % @ 1 ml/l recorded longest cocoon filament which were on par with untreated control. At 10 DAS, filament denier across all the treatments was on par with each other except in the treatments dinotefuron 20 SG @ 0.12 g/l and dichlorvos 76 EC @ 2.63 ml/l. At 40 DAS, finer denier was observed in the treatments pymetrozine 50 WG @ 0.6 g/l & 0.3 g/l, flonicamid 50 WG @ 0.3 g/l & 0.15 g/l, dinotefuron 20 SG @ 0.12 g/l. In the remaining treatments, the filament denier was on par with untreated control. Azadirachtin 1 % @ 2 ml/l recorded finer denier at 30 and 40 DAS. Buprofezin 25 SC @ 1ml/l recorded coarse denier on par with control at 10, 20, 30 DAS, whereas denier was finer with the same treatment at 40 DAS. Among the various treatments, dinotefuron 20 SG @ 0.25 g/l has performed well with reference to the reeling parameters.

**Keywords:** *B. mori*, mulberry, dinotefuron, flonicamid, DDVP

### Introduction

The superiority of silk as a textile fibre has been recognised from the time immemorial and even today, no other fabric can match its lustre and elegance. Mulberry being the sole food for silkworm *Bombyx mori* L. is infested by several pests. The insecticides applied for the management of mulberry pests have greater impact on silkworm rearing and subsequently on cocoon reeling parameters. Pesticide residues on mulberry leaves pose a severe threat to the silkworms, which in turn effect the silk quality (Yokoyama, 1962) [8]. Field observations in India indicated loss of cocoon yield and quality of silk from silkworms fed on mulberry leaves sprayed with insecticides (Narasimhanna, 1988) [7]. Silkworm being an insect is known to be very sensitive to insecticides, chemicals meant for insect pest control also affect the silkworms in terms of silk quality when they come in contact with the insecticides either directly or due to consumption of mulberry leaves treated with insecticides. Therefore, any application of insecticides on mulberry should be taken up very carefully. Sufficient waiting period has to be followed before harvest, so as to make leaves safe for consumption by silkworms. Recommended and widely used insecticide, dichlorvos being a contact poison, has proven lethal against the silkworms if fed immediately after spraying. Therefore, leaves will be harvested only after 15 days of waiting or safety period. Pest management in mulberry sericulture is absolutely selective because silkworm cannot tolerate even sub lethal doses of toxic compounds. In the light of above context, a study was initiated using different novel insecticides along with dichlorvos, has been notified for ban from import and manufacture with effective from January 2019 and complete ban from usage on all crops with effective from December 2020 (Anonymous, 2016) [1]. Corresponding to the reeling parameters to silkworm like cocoon filament length and filament denier.

## Materials and Methods

The commercial cross breed PM X CSR<sub>2</sub> was used for the study which were fed with V<sub>1</sub> leaves. The larvae were provided with chopped mulberry leaves of required quantity and quality. At 3<sup>rd</sup> instar, after 30 minutes of initial feeding, 90 larvae were transferred to each experimental tray in three replications along with the mulberry leaves. The design used was Completely Randomized Design. The mulberry leaves sprayed with insecticides *viz.*, buprofezin 25 SC (Applaud), pymetrozine 50 WG (Chess), flonicamid 50 WG (Ulala), dinotefuran 20 SG (Osheen), azadirachtin 1% (Neemazol) and dichlorvos (Nuvan) were fed to silkworms separately as per schedule. The silkworms were fed with 10, 20, 30 and 40 DAS insecticides. The cocoons of different treatments were harvested and were subjected for the reeling parameters *viz.*, single cocoon filament length and filament denier.

Five cocoons per replication were selected and each cocoon was reeled using euppovette and silk filament length was recorded. Filament length was calculated by the formula,  $L = R \times 1.125 \text{ m}$ . Where, in L was length of the silk filament (m), R was number of revolutions and 1.125 m represented circumference of the euppovette.

Silkworm filament denier was calculated by using the formula,

$$\text{Denier} = \frac{\text{Weight of silk filament (g)}}{\text{Length of silk filament (m)}} \times 9000$$

## Results and discussion

### Single cocoon filament length (m)

Single cocoon filament length recorded significant different among the four set of treatment where the silkworm batches were reared on the mulberry leaf treated with different insecticides on 10, 20, 30 and 40 DAS. The treatment untreated control recorded longest filament length (734.67 m) in the batches reared with leaves fed 10 DAS which was statistically on par with the filament length of dichlorvos 76 EC @ 2.63 ml/l (725.67 m), dinotefuron 20SG @ 0.25 g/l (722.67 m) and azadirachtin 1 % @ 1 ml/l (717 m). The shortest filament length was recorded in the treatment buprofezin 25 SC @ 2 ml/l (614.67 m). The other treatment recorded single cocoon filament length ranging from 643.33 to 697.67 m (Table 1). In the batches of silkworm fed with mulberry 20 DAS the range of filament length varied 716.33 to 410.83 m. The longest filament length in the treatment dinotefuron 20SG @ 0.25 g/l (716.33 m) recorded was statistically on par with dichlorvos 76 EC @ 2.63 ml/l (standard Check) (711 m), azadirachtin 1 % @ 2 ml/l (701.67 m) and azadirachtin 1 % @ 1 ml/l (697.67 m). These were followed by Dinotefuron 20 SG @ 0.12 g/l (669 m) which was on par with buprofezin 25 SC @ 1 ml/l (666.33 m), untreated control (662.33 m) and dichlorvos 76 EC @ 1.32 ml/l (647.67 m). Filament length of 634.67 m was recorded by pymetrozine 50 WG @ 0.3 g/l which was on par with buprofezin 25 SC @ 2 ml/l (619.67 m). The shortest filament length was recorded in the treatment flonicamid 50WG @ 0.3 g/l (410.83 m) (Table 1; Fig. 1).

When the silkworms were fed with mulberry leaf 30 DAS of insecticide, the longest single cocoon filament length was recorded in dinotefuron 20SG @ 0.25 g/l (728.00 m) which was on par with the highest filament length of untreated control (718.67 m), azadirachtin 1 % @ 2 ml/l (706.00 m), dichlorvos 76 EC @ 2.63 ml/l (704.00 m), pymetrozine 50 WG @ 0.6 g/l (693.00 m), flonicamid 50WG @ 0.15 g/l (682.00 m) and dinotefuron 20 SG @ 0.12 g/l (672.33 m). Buprofezin 25 SC @ 2 ml/l exhibited short filament length of 615 m which was statistically on par with Buprofezin 25 SC @ 1ml/l (654.00 m), Pymetrozine 50 WG @ 0.3 g/l (663.33 m), Azadirachtin 1 % @ 1 ml/l (637.00 m) and Dichlorvos 76 EC @ 1.32 ml/l (643.33 m) (Table 1; Fig. 1).

The single cocoon filament length varied from 374 to 716 m in the batches treated with mulberry leaves 40 DAS of insecticides. The longest filament length was exhibited by untreated control (716 m) which was statistically on par with dinotefuron 20SG @ 0.25 g/l (705.33 m). Azadirachtin 1 % @ 2 ml/l and dichlorvos 76 EC @ 2.63 ml/l recorded the next best higher and statistically on par filament length of 697.33 m. Dinotefuron 20 SG @ 0.12 g/l recorded filament length of 668 m, which was statistically on par with pymetrozine 50 WG @ 0.6 g/l (679.33 m) and buprofezin 25 SC @ 1 ml/l (664 m). Flonicamid 50 WG @ 0.15 g/l recorded filament length of 657.33 m which was on par with pymetrozine 50 WG @ 0.3 g/l 651.33 m. The shortest single cocoon filament length was recorded in flonicamid 50WG @ 0.3 g/l (374.00 m) (Table 1; Fig. 1).

Silkworm cocoon filament length was longest in untreated control and in the insecticide treatments *viz.*, dinotefuron 20 SG @ 0.25 g/l, dichlorvos 76 EC @ 2.63 ml/l at different time intervals of 10, 20, 30 DAS. Lowest filament length was recorded in flonicamid 50WG @ 0.3 g/l at 20, 30 and 40 DAS. This was followed by azadirachtin 1 % @ 1 ml/l recording shorter filament length at 40 DAS. However, at 10 and 20 DAS azadirachtin 1 % @ 1 ml/l recorded highest cocoon filament length on par with untreated control. Dinotefuron 20 SG is a third generation neonicotinoid with strong systemic and translaminar action. Upon ingestion, it acts on the nervous system of the insect and disrupts the normal functioning of nervous system by mimicking the functions of acetylcholine. This molecule is known to be effective on multiple sucking insects on crops and also known to be safer for non sucking insects like silkworms (Ghosh *et al.*, 2014) [4].

Bandyopadhyay *et al.* (2005) [2] noticed higher filament length with dichlorvos 0.1% and azadirachtin at 1 % compared to other insecticides assessed. In another study where azadirachtin 0.15% EC showed higher silkworm cocoon filament length in comparison with other insecticides *viz.*, emamectin benzoate, flubendiamide, abamectin and *Bacillus thuringiensis* (Kordy. 2014) [5]. The similar trend was observed in the present investigation also.


Gayathri (2007) [3] reported the adverse effects insecticides on cocoon filament length. Here, when fourth instar silkworms were fed with organophosphate insecticide, methyl demeton (0.05 %), cocoon filament length was significantly lower than that of untreated control.

**Table 1:** Effect of insecticides on single cocoon filament length (m)

Treatments	Filament length (m)			
	10 DAS	20 DAS	30 DAS	40 DAS
T <sub>1</sub> Buprofezin 25 SC @ 1ml/l	652.67 <sup>cd</sup>	666.33 <sup>cd</sup>	654.00 <sup>cdef</sup>	664.00 <sup>cd</sup>
T <sub>2</sub> Buprofezin 25 SC @ 2 ml/l	614.67 <sup>f</sup>	619.67 <sup>f</sup>	615.00 <sup>f</sup>	620.00 <sup>f</sup>
T <sub>3</sub> Pymetrozine 50 WG @ 0.3 g/l	631.67 <sup>ef</sup>	634.67 <sup>ef</sup>	663.33 <sup>cdef</sup>	651.33 <sup>de</sup>
T <sub>4</sub> Pymetrozine 50 WG @ 0.6 g/l	697.67 <sup>b</sup>	686.33 <sup>bc</sup>	693.00 <sup>abcd</sup>	679.33 <sup>c</sup>

T <sub>5</sub>	Flonicamid 50 WG @ 0.15 g/l	647.00 <sup>cde</sup>	687.00 <sup>bc</sup>	682.00 <sup>abcde</sup>	657.33 <sup>d</sup>
T <sub>6</sub>	Flonicamid 50 WG @ 0.3 g/l	0.00 <sup>g</sup>	410.83 <sup>g</sup>	409.33 <sup>gbcde</sup>	374.00 <sup>h</sup>
T <sub>7</sub>	Dinotefuron 20 SG @ 0.12 g/l	665.00 <sup>e</sup>	669.00 <sup>cd</sup>	672.33 <sup>a</sup>	668.00 <sup>cd</sup>
T <sub>8</sub>	Dinotefuron 20 SG @ 0.25 g/l	722.67 <sup>a</sup>	716.33 <sup>a</sup>	728.00 <sup>a</sup>	705.33 <sup>ab</sup>
T <sub>9</sub>	Azadirachtin 1 % @ 1 ml/l	717.00 <sup>a</sup>	697.67 <sup>ab</sup>	637.00 <sup>ef</sup>	570.00 <sup>g</sup>
T <sub>10</sub>	Azadirachtin 1 % @ 2 ml/l	694.67 <sup>b</sup>	701.67 <sup>ab</sup>	706.00 <sup>abc</sup>	697.33 <sup>b</sup>
T <sub>11</sub>	Dichlorvos 76 EC @ 1.32 ml/l	643.33 <sup>dc</sup>	647.67 <sup>de</sup>	643.33 <sup>def</sup>	636.33 <sup>ef</sup>
T <sub>12</sub>	Dichlorvos 76 EC @ 2.63 ml/l (std. check)	725.67 <sup>a</sup>	711.00 <sup>ab</sup>	704.00 <sup>abc</sup>	697.33 <sup>b</sup>
T <sub>13</sub>	Untreated control	734.67 <sup>a</sup>	662.3 <sup>ed</sup>	718.67 <sup>ab</sup>	716.00 <sup>a</sup>
F-test		*	*	*	*
SE.m ±		6.451	10.436	18.982	6.115
CD at 5 %		18.753	30.337	55.179	17.776

\*Significant at 5 %, DAS - Days after spraying.



T<sub>1</sub>- Buprofezin 25 SC @ 1ml/l  
 T<sub>2</sub>- Buprofezin 25 SC @ 2 ml/l  
 T<sub>3</sub>- Pymetrozine 50 WG @ 0.3 g/l  
 T<sub>4</sub>- Pymetrozine 50 WG @ 0.6 g/l  
 T<sub>5</sub>- Flonicamid 50 WG @ 0.15 g/l  
 T<sub>6</sub>- Flonicamid 50 WG @ 0.3 g/l  
 T<sub>7</sub>- Dinotefuron 20 SG @ 0.12 g/l  
 T<sub>8</sub>- Dinotefuron 20 SG @ 0.25 g/l  
 T<sub>9</sub>- Azadirachtin 1 % @ 1 ml/l  
 T<sub>10</sub>- Azadirachtin 1 % @ 2 ml/l  
 T<sub>11</sub>- Dichlorvos 76 EC @ 1.32 ml/l  
 T<sub>12</sub>- Dichlorvos 76 EC @ 2.63 ml/l  
 T<sub>13</sub>- Untreated control  
 DAS- Days after spraying

**Fig 1:** Effect of insecticides on Single cocoon filament length (m)

### Filament denier (D)

The values pertaining to the denier which was recorded by the effect of different insecticides when they were fed at different intervals of DAS were found significant in all the four batches of rearing. The coarse denier at 10DAS was found in dinotefuron 20 SG @ 0.25 g/l (2.61) which was found statistically on par with the treatments like untreated control (2.57), pymetrozine 50 WG @ 0.6 g/l (2.56), azadirachtin 1% @ 2 ml/l (2.53), buprofezin 25 SC @ 2 ml/l (2.51), buprofezin 25 SC @ 1 ml/l (2.50) and pymetrozine 50 WG @ 0.3 g/l (2.45). However, dinotefuron 20 SG @ 0.12 g/l (2.23) has exhibited the finest denier. Dichlorvos 76 EC @ 2.63 ml/l (standard Check) had yielded coarse filament (2.61) which was found to be statistically on par with all the other treatments, except flonicamid 50 WG @ 0.15 g/l (2.13) and flonicamid 50 WG @ 0.3 g/l (2.07) when the silkworms were reared in the batches treated with 20 DAS of insecticides (Table 2 ; Fig. 2).

Azadirachtin 1 % @ 1 ml/l (2.59) was the most coarse filament given among all the treatments, when the silkworms were reared with mulberry leaves treated with 30 DAS of insecticides. The fine denier was found in flonicamid 50 WG @ 0.3 g/l (1.94) and azadirachtin 1 % @ 2 ml/l (2.18). Similarly azadirachtin 1 % @ 1 ml/l (2.60) has given coarse

denier among the 40 DAS of insecticide batches. This was found on par with buprofezin 25 SC @ 1 ml/l (2.55), dinotefuron 20 SG @ 0.25 g/l (2.51), pymetrozine 50 WG @ 0.3 g/l (2.48), untreated control (2.47) and buprofezin 25 SC @ 2 ml/l (2.38). The finest denier among the batch was found in buprofezin 25 SC @ 1 ml/l (2.09) which was on par with pymetrozine 50 WG @ 0.6 g/l (2.19), azadirachtin 1% @ 2 ml/l (2.23), flonicamid 50 WG @ 0.3 g/l (2.24), flonicamid 50 WG @ 0.15 g/l (2.25) and dinotefuron 20 SG @ 0.12 g/l (2.32) which possessed similar finer denier (Table 2 ; Fig. 2).

At 10 DAS, filament denier across all the treatments was on par with each other except in the treatments dinotefuron 20 SG @ 0.12 g/l and dichlorvos 76 EC @ 2.63 ml/l. Similar trend was noticed at 20 DAS and 30 DAS. At 40 DAS, finer denier was observed in the treatments pymetrozine 50 WG @ 0.6 g/l & 0.3 g/l, flonicamid 50 WG @ 0.3 g/l & 0.15 g/l, dinotefuron 20 SG @ 0.12 g/l. In the remaining treatments, the filament denier was on par with untreated control. Azadirachtin 1 % @ 2 ml/l recorded finer denier at 30 and 40 DAS. In contrast these observation, Bandyopadhyay *et al.* (2005)<sup>[2]</sup> reported higher denier value of 2.5 when silkworms were fed with azadirachtin 1500 ppm & 300 ppm treated leaves. However, this study suggested decrease in denier

among other insecticides *viz.*, monocrotophos, acephate and dichlorvos.


Buprofezin 25 SC @ 1ml/l recorded coarse denier on par with control at 10, 20, 30 DAS. However denier was finer with the

same treatment at 40 DAS. Similar to results witnessed at 40 DAS during the present investigation, feeding of buprofezin 25% WP treated leaves resulted in decline in denier as compared to control as reported by Maria *et al.* (2000) [6].

**Table 2:** Effect of insecticides on denier (D)

Treatments	Denier			
	10 DAS	20 DAS	30 DAS	40 DAS
T <sub>1</sub> Buprofezin 25 SC @ 1ml/l	2.50 <sup>abc</sup>	2.28 <sup>cdef</sup>	2.51 <sup>ab</sup>	2.09 <sup>e</sup>
T <sub>2</sub> Buprofezin 25 SC @ 2 ml/l	2.51 <sup>ab</sup>	2.52 <sup>abc</sup>	2.41 <sup>abc</sup>	2.38 <sup>abcd</sup>
T <sub>3</sub> Pymetrozine 50 WG @ 0.3 g/l	2.45 <sup>abc</sup>	2.50 <sup>abc</sup>	2.27 <sup>bc</sup>	2.48 <sup>abc</sup>
T <sub>4</sub> Pymetrozine 50 WG @ 0.6 g/l	2.56 <sup>ab</sup>	2.51 <sup>abc</sup>	2.41 <sup>abc</sup>	2.19 <sup>de</sup>
T <sub>5</sub> Flonicamid 50 WG @ 0.15 g/l	2.44 <sup>bed</sup>	2.13 <sup>ef</sup>	2.45 <sup>abc</sup>	2.25 <sup>cde</sup>
T <sub>6</sub> Flonicamid 50 WG @ 0.3 g/l	0.00 <sup>e</sup>	2.07 <sup>f</sup>	1.94 <sup>d</sup>	2.24 <sup>cde</sup>
T <sub>7</sub> Dinotefuron 20 SG @ 0.12 g/l	2.23 <sup>d</sup>	2.22 <sup>def</sup>	2.37 <sup>abc</sup>	2.32 <sup>bcde</sup>
T <sub>8</sub> Dinotefuron 20SG @ 0.25 g/l	2.61 <sup>a</sup>	2.58 <sup>ab</sup>	2.43 <sup>abc</sup>	2.51 <sup>ab</sup>
T <sub>9</sub> Azadirachtin 1 % @ 1 ml/l	2.33 <sup>bcd</sup>	2.46 <sup>abcd</sup>	2.59 <sup>a</sup>	2.60 <sup>a</sup>
T <sub>10</sub> Azadirachtin 1 % @ 2 ml/l	2.53 <sup>ab</sup>	2.51 <sup>abc</sup>	2.18 <sup>cd</sup>	2.23 <sup>cde</sup>
T <sub>11</sub> Dichlorvos 76 EC @ 1.32 ml/l	2.25 <sup>cd</sup>	2.33 <sup>bde</sup>	2.34 <sup>abc</sup>	2.55 <sup>ab</sup>
T <sub>12</sub> Dichlorvos 76 EC @ 2.63 ml/l (std. check)	2.36 <sup>bed</sup>	2.61 <sup>a</sup>	2.40 <sup>abc</sup>	2.41 <sup>abcd</sup>
T <sub>13</sub> Untreated control	2.57 <sup>ab</sup>	2.43 <sup>abcd</sup>	2.55 <sup>ab</sup>	2.47 <sup>abc</sup>
F-test	*	*	*	*
SE.m ±	0.099	0.133	0.103	0.088
CD at 5 %	0.287	0.387	0.299	0.256

\*Significant at 5 %, DAS - Days after spraying.



T<sub>1</sub>- Buprofezin 25 SC @ 1ml/l  
 T<sub>2</sub>- Buprofezin 25 SC @ 2 ml/l  
 T<sub>3</sub>- Pymetrozine 50 WG @ 0.3 g/l  
 T<sub>4</sub>- Pymetrozine 50 WG @ 0.6 g/l  
 T<sub>5</sub>- Flonicamid 50 WG @ 0.15 g/l  
 T<sub>6</sub>- Flonicamid 50 WG @ 0.3 g/l  
 T<sub>7</sub>- Dinotefuron 20 SG @ 0.12 g/l  
 T<sub>8</sub>- Dinotefuron 20 SG @ 0.25 g/l  
 T<sub>9</sub>- Azadirachtin 1 % @ 1 ml/l  
 T<sub>10</sub>- Azadirachtin 1 % @ 2 ml/l  
 T<sub>11</sub>- Dichlorvos 76 EC @ 1.32 ml/l  
 T<sub>12</sub>- Dichlorvos 76 EC @ 2.63 ml/l  
 T<sub>13</sub>- Untreated control  
 DAS- Days after spraying

**Fig 2:** Effect of insecticides on denier (D)

## Conclusion

The filament length was longest in untreated control and in the insecticide treatments *viz.*, dinotefuron 20 SG @ 0.25 g/l and dichlorvos 76 EC @ 2.63 ml/l at different time intervals of 10, 20, 30 DAS. Lowest filament length was recorded in flonicamid 50WG @ 0.3 g/l at 20, 30 and 40 DAS. Azadirachtin 1 % @ 1 ml/l recording shorter filament length at 40 DAS. Dinotefuron molecule is known to be effective on multiple sucking insects on crops and also known to be safer for non sucking insects like silkworms and azadirachtin is an effective eco friendly insecticide. Silk filament denier in the silkworm batch treated with the mulberry leaves 10 DAS has

recorded coarse denier in dinotefuron 20SG @ 0.25 g/l (2.61) and the finest denier was recorded in the batches treated with flonicamid 50WG @ 0.3 g/l. On 20 DAS the silkworm batches reared recorded coarse denier in the treatment dichlorvos 76 EC @ 2.63 ml/l (2.33) and flonicamid 50WG @ 0.3 g/l recorded the finer denier of 2.07. At 30DAS, the batches fed with mulberry leaves exhibited coarse denier of 2.18 in the azadirachtin 1% @ 1 ml/l and the finer denier was recorded in the treatment flonicamid 50 WG @ 0.3 g / l (1.94). In the silkworm batch treated with the mulberry leaves 40 DAS, the coarse denier was recorded in azadirachtin 1% @ 1 ml/l (2.60) and the finer denier was recorded in the batches

treated with pymetrozine 50WG @ 0.6 g/l (2.19). Flonicamid is known to act on insects also as a repellent along with its action on stylet penetrations. Due to its repellent action the silkworms exhibited lesser reeling parameters when compared to other treatments. Since dinotefuron 20SG @ 0.25 g/l and azadirachtin 1% @ 1 ml/l are performing well with respect to reeling parameters they can be used as an effective novel molecules to replace DDVP in sericulture.

### References

1. Anonymous. The gazette of India. 2016; 3(11):8.
2. Bandyopadhyay UK, Santhakumar MV, Saratchandra B. Role of Insecticides and Botanicals in Regulating Whitefly (*Dialeuropora decempuncta*) Incidence and Their Influence on some Economic Traits of Silkworm (*Bombyx mori* L.). Ann. Pl. Protec. Sci. 2005; 13(1):48-53.
3. Gayathri M. Studies on tukra and powdery mildew diseases in mulberry and their impact on the growth and development of silkworm, B. Mori. *M.Sc. (Agri.) Thesis*, ANGRAU Bapatla, 2007, 42-112.
4. Ghosh A, Samantha A, Chatterjee ML. Dinotefuran, a third generation neonicotinoid insecticide for management of rice brown planthopper. African. J. Agric. Res. 2014; 9(8):750-754.
5. Kordy AM. Residual effect of certain pesticides on the mulberry silkworm (*Bombyx mori* L.). Middle East J App. Sci. 2014; 4(3):711-717.
6. Maria EV, Paschalis CH, Sergios K. Effects of Applaud on the growth of silkworm (Lepidoptera: Bombycidae). J Eco. Ent. 2000; 93(2):290-292.
7. Narasimhanna MN. Manual on silkworm egg production. CSB, Bangalore, 1988, 146.
8. Yokoyama T. Synthesized Science of Sericulture. Japan, 1962, 39-46.