



P-ISSN2349-8528

E-ISSN 2321-4902

IJCS 2016; 4(4): 218-221

© 2016 JEZS

Received: 16-05-2016

Accepted: 18-06-2016

Prince Kumar

Extension Lecturer, M.Sc -
Chemistry (Physical Chemistry),
F.G.M Govt College, Adampur
(Hisar), Haryana, India

A study on nitrogen Sulphur donor ligands and their characterization

Prince Kumar

Abstract

A series of Bi(III) and As(III) complexes with two N∩S donor ligands, 1-(4-chloro-2-oxo-2H-chromen-3-yl)-methylene)- thiosemicarbazide (L1H) and N -[1-(2-oxo-2H-chrome-3yl-ethylidene)-hydrazinecarbodithionic acid benzyl ester (L2H) have been synthesized by the reaction of BiCl₃ and Ph₃As with ligands in 1 : 1 and 1 : 2 molar ratios. All the synthesized compounds were characterized by elemental analyses, melting point determinations, and a combination of electronic, IR, ¹H NMR, ¹³C NMR spectroscopic techniques, and X-ray diffraction for structure elucidation.

In order to evaluate the effect of metal ions upon chelation, both the ligands and their complexes have been screened for their antimicrobial activity against the various pathogenic bacterial and fungal strains. The metal complexes have shown to be more antimicrobial against the microbial species as compared to free ligands. Both the ligands and their corresponding metal complexes have been tested for their antifertility activity in male albino rats. The marked reduction in sperm motility and density resulted in infertility. Significant alterations were found in biochemical parameters of reproductive organs in treated animals as compared to control group.

Keywords: Sulphur, Nitrogen, Ligand, Donor

Introduction

Interest in coordination chemistry is increasing continuously with the preparation of organic ligands containing a variety of donor groups, and it is multiplied manifold when the ligands have biological importance ^[1]. The number and diversity of nitrogen and sulfur chelating agents used to prepare new coordination and organometallic compounds have increased rapidly during the past few years ^[2-4]. Sulfur compounds and their metal complexes have antimicrobial activity and showed a high dependence on their substituents ^[5, 6].

Organic compounds containing -C₆H₄S moiety are well known for their significant biological activities. The activity may be due to the presence of multi-coordination centers having the ability to form stable chelates with the essential metal ions which the organisms need in their metabolism. Interest in metal complexes of sulfur-nitrogen chelating agents, especially those formed from thiosemicarbazide ^[7] and S-alkyl/benzyl esters of dithiocarbamic acid has been stimulated by their interesting physicochemical properties and potentially useful pharmacological properties ^[8]. In the present paper, we describe the synthesis of metal complexes with Schiff bases using microwave-assisted technique.

Bismuth compounds have been used in medicine for more than two centuries ^[9]. Applications have been widespread, due to bismuth's antiseptic, astringent, protective, antacid, antisecretory, and local gastrointestinal properties. Organoarsenicals were the first compounds applied for the successful treatment of syphilis (they have since been replaced by penicillin) and have been used as feed additives in livestock to prevent bacterial and parasitic infections.

Organoarsenic(III) compounds are known to be biologically active and find applications in various fields. Antimicrobial activities of some arsenic(III) complexes with Schiff bases have been evaluated ^[10]. The focus of our present communication is on the exploration of the studies on the synthetic, structural, and biological aspects of Bi(III) and As(III) complexes of some stereochemical as well as biological interest with monobasic bidentate thiosemicarbazone and hydrazinecarbodithionic acid ligands.

Correspondence**Prince Kumar**

Extension Lecturer, M.Sc -
Chemistry (Physical Chemistry),
F.G.M Govt College, Adampur
(Hisar), Haryana, India

2. Experimental

2.1. Materials and Methods

BiCl₃ and Ph₃As as well as 3-acetylcoumarin and 4-hydroxycoumarin were purchased from Alfa Aesar and used as such. All the chemicals and solvents were dried and purified by standard methods. The reactions were carried out under strictly anhydrous conditions. Molecular weights were determined by the Rast camphor method. Chlorine was estimated by Volhard's method. Bismuth was estimated complexometrically. Arsenic was estimated iodimetrically. Nitrogen was estimated by the Kjeldahl method, and sulfur was estimated by the Messenger method ^[11].

Carbon and hydrogen analyses were performed at the CDRI, Lucknow, India. The electronic spectra were recorded on a Varian-Cary/2390 spectrophotometer at RSIC, IIT, Chennai, India. Infrared spectra of the ligands and their complexes were recorded on a Nicolet Magna FTIR- 550 spectrophotometer using KBr pellets. ¹H and ¹³C NMR spectra were recorded on a JEOL-AL-300 FT NMR spectrometer in DMSO-d₆. X-ray powder diffractogram of a representative compound was obtained on a Philip Model PW1840 automatic diffractogram using Cu(K α) target with Mg filter. The wavelength used was 1.540598 Å.

2.2. Preparation of the Ligands

Two different routes, microwave-assisted synthesis and conventional method, were employed for the synthesis of the ligands (see Scheme 1). 3-Formyl-4-chlorocoumarin. Phosphorus oxychloride (10 mL) was added dropwise to a solution of dimethylformamide (DMF) (20 mL) keeping the temperature below 5°C.

Solution of 4-hydroxycoumarin (4.0 g) in DMF (10 mL) was then gradually added to the mixture with constant stirring and maintaining the temperature of the reaction mixture below 5°C. The reaction mixture was then allowed to stand at room temperature for 2 h and then heated on a steam bath for 1 h. After cooling, the reaction mixture was poured onto crushed ice and neutralized with sodium carbonate. A solid product was immediately formed which was crystallized from ethanol to give a yellow solid (80%), MP 115°C ^[12].

2.2.1. Microwave-Assisted Synthesis

The ligands were prepared by the condensation of 3-formyl-4-chlorocoumarin with thiosemicarbazide and 3-acetyl coumarin with Sbenzylthiocarbazate in the presence of sodium acetate. The reaction mixture was irradiated in the microwave oven by taking 2-3 mL solvent. The reactions were completed in a short period (5–7 min). The resulting precipitate was then recrystallized with alcohol and dried under vacuum. These were characterized and analysed before use.

Elemental analyses (N and S) were conducted using the methods mentioned above, and their results were found to be in good agreement with the calculated values. The structures of the ligands have been shown in Figure 1.


The above ligands were also synthesized by a thermal method, where instead of few drops of alcohol the starting materials of the ligands were dissolved in ~100 mL of alcohol and the contents were refluxed for nearly 4 to 5 h. The solution was then concentrated under reduced pressure, which on cooling gave crystalline precipitates. These were recrystallized twice in alcohol.

2.3. Preparation of the Metal Complexes

2.3.1. Synthesis of the Bi(III) Complexes

For the synthesis of the complexes, BiCl₃ and sodium salt of the ligands (prepared by adding the corresponding weight of sodium metal to the ligand) in 5 mL of dry methanol in 1 : 1 and 1 : 2 molar ratios were irradiated inside a microwave oven for about 5–8 min. The products were recovered from the microwave oven and dissolved in few mL of dry methanol. The white precipitate of sodium chloride formed during the course of the reaction was removed by filtration, and the filtrate was dried under reduced pressure.

The resulting product was repeatedly washed with petroleum ether and then finally dried at 40–60°C/0.5 mmHg for 3-4 h. The purity was further checked by thin layer chromatography using silica gel-G. Chlorobismuth(III) complexes were also synthesized by the thermal method. The reaction mixtures were heated under reflux for 13–17 h and filtered to remove NaCl, and the solvent was removed by the same procedure mentioned above, which was adopted to get the complexes.






Fig 1: 3D molecular structures of the ligands

3. Biological Assay

All the compounds were evaluated for their antimicrobial properties. MIC was recorded as minimum concentration which inhibits the growth of microorganism. The results obtained were compared with those of the standard drug streptomycin for bacteria and bavistin for fungi. The microorganisms used were *Escherichia coli* (ATCC25922), *Bacillus subtilis* (ATCC6633), *Fusarium oxysporum* (ATCC7808) and *Rhizopus nigricans* (ATCC6227b). Both ligands and their corresponding metal complexes have been tested *in vivo* for their antifertility activity in male albino rats.

3.1. In Vitro Study

3.1.1. Antibacterial Screening

The newly prepared compounds were screened for their antibacterial activity against *Escherichia coli* (ATCC25922) and *Bacillus subtilis* (ATCC6633) by paper disc plate method^[13]. Each compound was dissolved in DMSO, and solutions of the concentrations (500 and 1000 ppm) were prepared separately. Paper discs of Whatman filter paper (no. 42) of uniform diameter (5 mm) were cut and sterilized in an autoclave. The paper discs soaked in the desired concentration of the complex solutions were placed aseptically in the petri dishes containing nutrient agar media (agar 20 g + beef extract 3 g + peptone 5 g) seeded with *Escherichia coli* (ATCC25922) and *Bacillus subtilis* (ATCC6633) separately.

The petri dishes were incubated at 37°C, and the inhibition zones were recorded after 24 h of incubation. The antibacterial activity of standard antibiotic streptomycin was also recorded using the same procedure. The medium with DMSO as a solvent was used as a negative control. The experiments were performed in triplicates.

3.1.2. Antifungal Screening

The antifungal activities of the standard fungicide (Bavistin), ligands, and complexes were tested for their effect on the growth of microbial cultures and studied for their interaction with *Fusarium oxysporum* (ATCC7808) and *Rhizopus nigricans* (ATCC6227b) using Czapek's agar medium^[14] having the composition, glucose 20 g, starch 20 g, agar-agar 20 g, and distilled water 1000 mL. To this medium was added requisite amount of the compounds after being dissolved in methanol so as to get the certain concentrations (100 and 200 ppm).

The medium then was poured into Petri Plates, and the spores of fungi were placed on the medium with the help of inoculum's needle. These Petri Plates were wrapped in polythene bags containing a few drops of alcohol and were placed in an incubator at $\pm 30^\circ\text{C}$. The controls were also run, and three replicates were used in each case.

3.1.3. Determination of Minimum Inhibitory Concentration (MIC)

The minimum inhibitory concentration (MIC) is the lowest concentration of the antimicrobial agent that prevents the development of viable growth after overnight incubation. The determination of the MIC involves a semiquantitative test procedure, which gives an approximation to the least concentration of an antimicrobial needed to prevent microbial growth.

The minimum inhibitory concentration was determined by microbroth dilution method. Inoculum of the overnight culture was prepared. In a series of tubes, 1 mL each of complex solution with different concentrations was taken and 0.4 mL of the inoculum was added to each tube.

Further 3.5 mL of the sterile water was added to each of the test tubes. These test tubes were incubated for 18 h and observed for the presence of turbidity. The absorbance of the suspension of the inoculum was observed with spectrophotometer at 555 nm. The end result of the test was the minimum concentration of the antimicrobial agent (test materials) which gave a clear solution, that is, no visual growth.

4. Results and Discussion

The elemental analysis and spectral data suggested the formation of the ligands (L1H and L2H) and their complexes of the type $[\text{Cl}_2\text{Bi}(\text{L}1)]$, $[\text{ClBi}(\text{L}1)_2]$, $[\text{Cl}_2\text{Bi}(\text{L}2)]$, $[\text{ClBi}(\text{L}2)_2]$, $[\text{Ph}_2\text{As}(\text{L}1)]$, $[\text{PhAs}(\text{L}1)_2]$, $[\text{Ph}_2\text{As}(\text{L}2)]$, and $[\text{PhAs}(\text{L}2)_2]$. The resulting complexes are colored solid, which were found to be soluble in DMF and DMSO. Molecular weight determinations showed that they are monomeric in nature. The molar conductances of 10⁻³ M solutions of the complexes in dry DMF lie in the 12–15 ohm⁻¹ cm² mol⁻¹ range, indicating that they are nonelectrolytes.

The electronic spectra of the ligands show a band due to the $>\text{C}=\text{N}$ chromophore observed at 365 nm that shifts to the lower wavelength region in the spectra of the complexes. Such a shift in the $n-\pi^*$ band is probably due to the donation of the lone pair of electrons by the nitrogen of the ligand to the central metal atom. Two bands at 319–320 and 298–300 nm assigned to $\pi-\pi^*$ transitions, within the benzene and coumarin rings and azomethine linkage also show some shifting in their position due to bond formation.

The broad band, due to ($-\text{NH}$) vibrations, disappears in the spectra of the complexes, indicating the deprotonation of this group on coordination with the metal atom. The marked shift (10–20 cm⁻¹) of ($>\text{C}=\text{N}$) band observed in all the complexes indicates the involvement of azomethine nitrogen upon complexation. The band due to ($>\text{C}=\text{S}$) is shifted towards lower frequencies in the complexes indicating coordination of sulfur to the central metal atom.

The spectra of the free ligands display two sharp bands at 3340–3500 and 3350–3490 cm^{-1} due to asym. and sym. vibrations of NH_2 group, respectively, which remain at almost the same positions in the spectra of the complexes, suggesting that the NH_2 group is not involved in chelation. Some new bands observed in the regions 439–442 and 422–425 cm^{-1} for ν ($\text{As} \leftarrow \text{N}$), ν ($\text{As}-\text{S}$), respectively. The band in 450–470 cm^{-1} range may be assigned to ν ($\text{As}-\text{Ph}$) vibrations in the respective complexes.

Further evidence for the coordinating mode of the ligands was obtained from ^1H NMR spectra. The ^1H NMR spectra of the ligands recorded in $\text{DMSO}-d_6$ exhibit a broad peak at δ 8.59–8.30 ppm due to $-\text{NH}$ proton. The $-\text{NH}$ proton signal of the ligands disappears in the complexes. The absence of this signal in these complexes suggests that this proton has been lost via thioenolization of $>\text{C}=\text{S}$ group and coordination of sulfur to the metal atoms, respectively, has taken place.

The $>\text{CH}=\text{N}$ proton moves downfield in the complexes in comparison with its original positions in the ligands due to coordination of $>\text{C}=\text{N}$ to the metal atom. The ^1H NMR spectrum of the ligand (L2H) exhibits $-\text{CH}_2$ proton signals at δ 4.15–4.16 ppm, and aromatic proton signals at δ 6.38–7.50 ppm and these remain at the same position in the spectra of the metal complexes. In the case of ^{13}C NMR spectra, the considerable shifts in the position of the carbon atoms attached to the different participating groups clearly indicate the bonding of azomethine nitrogen to the metal atom. The signal due to lactone $>\text{C}=\text{O}$ carbon atoms remains almost same position in the ^{13}C NMR spectra of the metal complexes while mark shifts in the position of the carbon atoms attached to the sulfur atom shows that proton is lost via thioenolization.

5. Conclusion

We describe the synthesis, characterization, and biological activity of $\text{Bi}(\text{III})$ and $\text{As}(\text{III})$ complexes. On the basis of analytical and spectral data, tetra- and pentacoordinated geometries have been proposed for the $\text{Bi}(\text{III})$ and $\text{As}(\text{III})$ complexes.

The antimicrobial activity results indicated that the complexes showed promising antibacterial and antifungal activities, but compounds $[\text{ClBi}(\text{L}1)_2]$ and $[\text{PhAs}(\text{L}2)_2]$ showed highest activity against bacterial strain *B. subtilis* ATCC6633 ($\text{MIC} = 10 \mu\text{g}/\text{mL}$) and fungal strain *F. oxysporum* ATCC7808 ($\text{MIC} = 16 \mu\text{g}/\text{mL}$). The results suggested that the ligand (L2H) is most effective in reducing fertility, and addition of bismuth and arsenic moiety to this ligand enhanced its activity.

References

- Kulkarni A, Avaji PG, Bagihalli GB, Patil SA, Badami PS. Synthesis, spectral, electrochemical and biological studies of $\text{Co}(\text{II})$, $\text{Ni}(\text{II})$ and $\text{Cu}(\text{II})$ complexes with Schiff bases of 8-formyl-7-hydroxy-4-methyl coumarin. *Journal of Coordination Chemistry*. 2009; 62(3):481-492.
- Sharma K, Singh RV, Fahmi N. Palladium (II) and platinum(II) derivatives of benzothiazoline ligands: synthesis, characterization, antimicrobial and antispermato-genic activity, *Spectrochimica Acta Part A*, 2011; 78(1):80-87.
- Basu Baul TS. Antimicrobial activity of organotin (IV) compounds, *Applied Organometallic Chemistry*, 2008; 22:195-204.
- Li M-X, Zhou J, Zhao H, Chen C-L, Wang J-P. Iron (III) complex of 2-acetylpyrazine thiosemicarbazone: synthesis, spectral characterization, structural studies and antitumoral activity. *Journal of Coordination Chemistry*. 2009; 62(9):1423-1429.
- Shanker D, Sharma RK, Sharma J, Rai AK, Singh YP, Metal-induced rearrangement of benzothiazoline ring: synthesis and characterization of some new organoantimony (V) derivatives of N, O, and S atom containing Schiff base ligands. *Heteroatom Chemistry*, 2007; 18(1):70-75.
- Singh K, Singh DP, Singh Barwa M, Tyagi P, Mirza Y. Some bivalent metal complexes of Schiff bases containing N and S donor atoms. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2006; 21(6):749-755.
- Yadav S, Singh RV. Ferrocenyl-substituted Schiff base complexes of boron: synthesis, structural, physico-chemical and biochemical aspects, *Spectrochimica Acta Part A*, 2011; 78(1):298-306.
- Akbar Ali M, Mirza AH, Voo CW, Tan AL, Bernhardt PV. The preparation of zinc(II) and cadmium(II) complexes of the pentadentate N_3S_2 ligand formed from 2,6-diacetylpyridine and S-benzylthiocarbamate (H_2SNNNS) and the X-ray crystal structure of the novel dimeric $[\text{Zn}_2(\text{SNNNS})_2]$ complex, *Polyhedron*, 2003; 22(27):3433-3438.
- Sun H, Li H, Harvey I, Sadler PJ. Interactions of bismuth complexes with metallothionein(II). *Journal of Biological Chemistry*. 1999; 274(41):29094-29101.
- Karra R, Singh YP, Rai AK. Synthesis and structural elucidation of some new phenylarsenic(III) derivatives of N(-substituted) S-benzyl dithiocarbazates. *Phosphorus, Sulfur and Silicon and Related Elements*, 2000; 166:125-135.
- Vogel AI. *A Textbook of Quantitative Chemical Analysis*, Pearson Education Ltd.: Thames Polytechnique, London, UK, 6th edition, 2006.
- Dawara L, Singh RV. Synthesis, spectroscopic characterization, antimicrobial, pesticidal and nematocidal activity of some nitrogen-oxygen and nitrogen-sulfur donor coumarins based ligands and their organotin(IV) complexes. *Applied Organometallic Chemistry*, 2011; 25(9):643-652.
- Gaur S, Fahmi N, Singh RV. Coordination behavior of unsymmetrical ligand complexes of diorganotin and diorganosilicon derived from Schiff bases, *Phosphorus, Sulfur and Silicon and the Related Elements*, 2007; 182(4):853-862.
- Shrivastava S, Fahmi N, Singh RV. Studies on chromium(III) complexes with active nitrogen, oxygen and sulfur donor ketimines synthesized under microwave conditions. *Journal of Sulfur Chemistry*, 2010; 31(6):515-524.