



P-ISSN 2349-8528
E-ISSN 2321-4902
IJCS 2017; 5(1): 01-04
© 2017 JEZS

Received: 01-11-2016
Accepted: 02-12-2016

Tekale AS
Research Laboratory,
Department of Chemistry,
Shivaji Mahavidyalaya, Udgir,
Dist. Latur (MS), India.

Shaikh SAL
Research Laboratory,
Department of Chemistry,
Shivaji Mahavidyalaya, Udgir,
Dist. Latur (MS), India.

Synthesis and characterisation of substituted quinoline by Vilsmeier-Haack reagent

Tekale AS and Shaikh SAL

Abstract

For the synthesis of substituted i.e. 2-chloro-3-formyl-8-methyl Quinoline an highly efficient, simple and convenient method through Vilsmeier-Haack reagent has been developed in high yields, less raw material, and less time requiring for completion of reaction. In the present research work the substituted quinoline and acetanilide are synthesized through Vilsmeier reaction has been reported to the substitution reaction by substituted acetanilide bearing electron donating group at ortho position and electron attracting group at ortho position are investigated and it is good in all case. Further, the nucleophilic substitution reaction of the substituted quinoline is also investigated. Similarly in quinoline the -CHO (formyl) group is subjected for further transformation of formyl group in to (4'-NO₂ nitro group, 3'-N-NH-C₆H₅) 3'-CH₃ group and 3'-N-benzylidene group.

Keywords: Synthesis and Characterisation of substituted Quinoline Vilsmeier reagent, 2-chloro-3-formyl-8-methyl Quinoline, TLC, Formylation.

1. Introduction

Substituted quinolines are very simply, efficiently and conveniently synthesized in less time and high yield with less amount of raw material under mild condition with the help of Vilsmeier-Haack reagent (DMF + POCl₃) and substituted acetanilide. Quinolines has been possesses a wide spectrum of biological activities. Now a days quinoline derivatives are used for a convenient starting material of various further substituted quinoline. Formylation of heterocyclic compounds [3] may be achieved by heating heterocyclic compound with Vilsmeier reagent. i.e (DMF + POCl₃) phosphorous oxy chloride and dimethyl formamide, the intermediate is hydrolysed in the presence of mildbase give 2-(ortho)-substituted heterocyclic compound [7]. This reaction is known as Vilsmeier-Haack reaction.

2. Methods and Materials

All chemicals were used of analytical grade from sigma Aldrich and S.d. Fine Chem. Limited, and the solvents used were purified by standard methods. All melting points were taken in open capillaries and are a correct. He progress of all reaction of the synthesized compound was monitored by TLC. i.e (thin layer chromatography). TLC was runned using TLC aluminium sheets adsorbent silica gel, chloroform solvent petether, ethylacetate (9:5, 0:5). Spot were visualized using solid iodine fumes, by the keeping of TLC plate in iodine chamber.

2.2 Experimental Procedure for

The completion of reaction and purity of synthesized products were monitored by silica gel TLC melting point determined by open capillary method. The substituted Quinoline compounds were synthesized by the method of Meth-Cohn et al [1-2].

D) Ortho- substituted acetanilide

Take 5 ml of ortho-substituted aniline, 5ml of acetic anhydride and 5 ml of glacial acetic acid in a 150 ml conical flask. Shake the reaction mixture thoroughly and then reflux, it for 60 minutes using an air condenser. After completion of requiring duration of reaction, pour the hot reaction mixture to 200 ml of ice cold water in a beaker with constant stirring. Filter the product and wash with water. Recrystallise the crude acetanilide either from boiling water or dilute acetic acid.

Correspondence


Tekale AS
Research Laboratory,
Department of Chemistry,
Shivaji Mahavidyalaya, Udgir,
Dist. Latur (MS), India.

II) Quinoline Derivative

(Synthesis of 2-Chloro-8-methyl-3-formyl Quinoline): Take Dimethyl formamide (DMF) 5 ml and is cooled to 0°C temperature in a flask equipped with a drying tube. Then 18 ml of POCl₃ (Phosphorous oxy chloride) is added drop wise with stirring to it. Then to this solution add 4 grams of ortho methyl acetanilide. After few minutes the solution of reaction mixture is refluxed for 6-8 hours using air condenser and the temperature of reaction mixture is maintained between 80-90 °C. After completion of requiring duration, the reaction mixture is cooled and poured in 100 ml beaker containing ice-cold water and stirred about half an hour then filter precipitated quinoline and wash with water and dried, recrystallized from ethyl acetate.

III) (Substituted benzylidene) Quinoline

In a clean and dry round bottom flask take recrystallized 2-chloro-3-formyl-8-methyl Quinoline about (2 mmole) and (2 mmole) of 4-Nitro aniline, add 8 ml of methanol and 4-5 drops of glycolic acetic acid. Shake the reaction mixture thoroughly and then reflux it for about 1-2 hours using air condenser. After completion of requiring duration of reaction cool and pour the reaction mixture in an ice-cold water, filter and dried, Recrystallized it. From ethyl acetate.

Reaction**2.3 Characterization data of the synthesized compound**

Acetanilide: Yield: 54.54%; m.p.:114⁰ C; Anal. Calcd for C₈H₉NO: C:71.11%; H:6.66%; N:10.37;O:11.85%; IR (KBr)cm-1: 3295 (N-H),1664 (CO), 1584 (C=C). 1H-NMR (300 MHz, DMSO-*d*₆); δ 8.72 (s, 1H, NH), 2.1 (s, 3H, CH₃), 7.2 (d, 1H, Ar-H), 7.1 (s, 1H, Ar-H), 7.0 (d, 1H, Ar-H).

13C-NMR (75 MHz, DMSO-*d*₆); δ 138.2 (ipso), 169.5 (CO), 24.1 (CH₃), 120.4, 128.7, 124.1.

2-Methyl Acetanilide: Yield: 52.55%; m.p.: 109-112⁰C; Anal. Calcd for C₉H₁₁NO: C: 72.11%; H: 7.38%; N:9.39%; IR (KBr) cm-1: 3279 (N-H), 1658 (CO), 1581 (C=C), 2851 (Ar-CH₃-C-H).

1H-NMR (300 MHz, DMSO-*d*₆); δ 8.69 (s, 1H, NH), 2.1 (s, 3H, CH₃), 4.25 (s, 3H, Ar-CH₃), 7.2 (d, 1H, Ar-H), 7.1 (s, 1H, Ar-H), 7.0 (d, 1H, Ar-H).

13C-NMR (75 MHz, DMSO-*d*₆); δ 137.1 (ipso), 170(CO), 21, (CH₃), 137.5(ArC-CH₃),119, 127, 122.

2-Nitro Acetanilide: Yield: 57.62%; m.p.:90-94⁰C; Anal. Calcd for C₈H₈N₂O₃:C:53.33%; H:4.44%; N:15.55%.O:26.66%; IR (KBr) cm-1: 3298 (N-H),1680 (CO), 1585 (C=C),1514 (Ar-NO₂).1H-NMR (300 MHz, DMSO-*d*₆); δ 8.73 (s, 1H, NH), 2.2 (s, 3H, CH₃), 7.2 (d, 1H, Ar-H), 7.0 (s, 1H, Ar-H), 6.9 (d, 1H, Ar-H).

13C-NMR (75 MHz, DMSO-*d*₆); δ 136.9 (ipso), 169 (CO), 24.2 (CH₃), 148.3(ArC-NO₂),121,127,123.

4-Chloro Acetanilide: Yield: 53.11%; m.p.: 176-178 °C; Anal. Calcd for C₈H₈NOCl: C:56.80% H:4.73%; N:8.28%.O:9.46%; Cl:20.71%;IR (KBr) cm-1: 3292 (N-H), 1665 (CO), 1585 (C=C), 759 (C-Cl).1H-NMR (300 MHz, DMSO-*d*₆); δ 8.70 (s, 1H, NH), 2.1 (s, 3H, CH₃), 3.6 (HC-Cl), 7.2 (d, 1H, Ar-H), 7.1 (s, 1H, Ar-H),7.0 (d, 1H, Ar-H).

13C-NMR (75 MHz, DMSO-*d*₆); δ 138 (ipso), 169 (CO), 23.9 (CH₃), 156 (Ar-Cl), 131,126,120.

2-Chloro-3-formyl Quinoline: Yield: 62.82%; m.p.:143 °C; Anal. Calcd for C₁₀H₆NOCl: C: 62.82% H: 3.14%; N:7.32%.O:8.37%; Cl:18.32%; (CDCl₃); δ 1H NMR : 10.5 (s,1H,CHO),8.8 (s,1H,H-4),8.1(m,1H,H-6),7.7(m,1H,H-7). 13C-NMR (CDCl₃); δ 189.12.

MS m/z: 193(M⁺⁺+2), 191(M⁺⁺), 190,162,155,127.

2-Chloro-3-formyl-8-Methyl Quinoline: Yield: 63%; m.p.:138 °C; Anal. Calcd for C₁₁H₈NOCl: C: 64.39% H: 3.90%; N: 6.82%.O:7.80%; Cl: 17.07%; (CDCl₃); δ 1H NMR : 10.4 (s, 1H,CHO),8.7 (s,1H,H-4), 8.1- 7.4 (m,3H,H-5,H-6 andH-7), 2.8 (s,3H, CH₃).13C-NMR (CDCl₃); δ 189.51.

2,6-Dichloro-3-formyl Quinoline :Yield:69%; m.p.:138 °C; Anal. Calcd for C₁₀H₅NOC1₂: C:53.33% H:2.22%; N:6.22%.O:7.11%; Cl:31.11%; (CDCl₃); δ 1H NMR : 10.8 (s,1H,CHO),8.6 (s,1H,H-4),8.1 (m,1H,H-8),7.7 (m,1H, H-5).13C-NMR (CDCl₃); δ 189.49.IR (KBr) cm-1: 3050, 1693, 1628, 1379, 1038.

2-Chloro-3-formyl-8-Nitro Quinoline: Yield: 65%; m.p.:124⁰C; Anal. Calcd for C₁₀H₅N₂O₃Cl: C: 50.84% H: 2.11%; N:11.86%.O:20.33%; Cl:14.83%; (CDCl₃); δ 1H NMR : 10.5 (s,1H,CHO),8.9 (s,1H,H-4), 7.7 (m,1H, H-5), 8.1 (m,1H,H-6,H-7). 13C-NMR (CDCl₃); δ 189.31.

2-Chloro-3-(3'-methyl benzylidene)-8-Methyl Quinoline: Yield: 57%; m.p.:197⁰ C; Anal. Calcd for C₁₈H₁₅N₂Cl: C:73.46% H:5.10%; N:9.52%; Cl:11.90%; IR (KBr) cm-1: 2925 (-CH₃), 1333 (C-N), 1622 (C=N) imine,776 (C-Cl),1588 (C=C) Aromatic. (CDCl₃); δ 1H NMR: 7.2-7.6 (8Hm Ar-H), 8.9 (1H,sCH=N),2.5 (6H,s,-CH₃).

2-Chloro-3-(4'-nitro benzylidene)-8-Methyl Quinoline: Yield: 57%; m.p.:140 °C; Anal. Calcd for C₁₇H₁₂N₃O₂Cl: C:62.76% H: 3.69%; N: 12.92%.O:9.84%; Cl: 10.76%; IR (KBr) cm-1: 2930 (-CH₃), 1342 (C-N), 1635 (C=N) imine,775 (C-Cl), 1580 (C=C) Aromatic 1525 (N=O). (CDCl₃); δ 1H NMR: 7.1-7.63 (8Hm Ar-H), 8.3 (1Hs, CH=N),2.48 (3H,s,-CH₃).

2-Chloro-3-(benzylidene)-8-Methyl Quinoline: Yield: 65%; m.p.:181 °C; Anal. Calcd for C₁₇H₁₃N₂Cl: C: 72.85%;H:4.64%; N:10%; Cl:12.5%; IR (KBr) cm⁻¹: 2922 (-CH₃), 1323 (C-N), 1628 (C=N) imine, 770 (C-Cl), 1585 (C=C) Aromatic.(CDCl₃); δ 1H NMR: 7.1-7.68 (9Hm Ar-H), 8.8 (1Hs, CH=N),2.43 (3Hs,- CH₃).

2-Chloro-3-(benzylidenehydrazido)-8-Methyl Quinoline: Yield: 62%; m.p.:207°C; Anal. Calcd for C₁₇H₁₄N₃Cl: C:69.17% H: 4.74%; N: 14.23%; Cl: 11.86%; IR (KBr) cm⁻¹: 2932 (-CH₃), 1340 (C-N), 1637 (C=N) imine, 777 (C-Cl), 1582 (C=C) Aromatic. (CDCl₃); δ 1H NMR: 7.1-7.8 (9Hm Ar-H), 8.73 (1Hs, CH=N), 2.44 (3H, s,- CH₃),3.7 (=N-NH-).

Table 1: Physical data of synthesized compounds

SR. No.	Molecular formula of Compound	M.P. in °C	Yield in GM	% Yield in GM	Molecular Weight.	Elemental analysis in%				
						C	H	N	O	Cl
1	C ₈ H ₉ NO	114 °C	3.90	54.54	135.16	71.11	6.66	10.37	11.85	
2	C ₉ H ₁₁ NO	109-112 °C	3.60	52.55	149	72.48	7.38	9.39	10.73	
3	C ₈ H ₈ N ₂ O ₂	90-94 °C	3.40	57.62	164	58.53	4.87	17.07	19.51	
4	C ₈ H ₈ NOCl	176-178 °C	3.50	53.11	169	56.80	4.73	8.28	9.46	20.71
5	C ₁₀ H ₆ NOCl	143 °C	6	62.82	191	62.82	3.41	7.32	8.37	18.32
6	C ₁₁ H ₈ NOCl	138 °C	3.46	63	205	64.39	3.90	6.82	7.80	17.07
7	C ₁₀ H ₅ N ₂ O ₃ Cl	124 °C	3.40	65	236	50.84	2.11	11.86	20.33	14.83
8	C ₁₀ H ₅ NOCl ₂	138 °C	3.93	69	225	53.33	2.22	6.22	7.11	31.11
9	C ₁₈ H ₁₅ N ₂ Cl	197 °C	1.67	57	294	73.46	5.10	9.52		11.90
10	C ₁₇ H ₁₂ N ₃ O ₂ Cl	140 °C	1.86	57	325	62.76	3.69	12.92	9.84	10.76
11	C ₁₇ H ₁₄ N ₃ Cl	207 °C	1.82	62	295	69.15	4.74	14.23		11.86
12	C ₁₇ H ₁₃ N ₂ Cl	181 °C	1.82	65	280	72.85	4.64	10		12.5

Table 2: Molecular formula, Retention factor and Time period at 80-90 °C of Synthesized compound

SR. No.		Molecular Formula	Rf Value	Time Period at 80-90 °C
1	1-a	C ₈ H ₉ NO	0.63	60 min
2	1-b	C ₉ H ₁₁ NO	0.67	60 min
3	1-c	C ₈ H ₈ N ₂ O ₃	0.68	10-15 min
4	1-d	C ₈ H ₈ NOCl	0.71	60 min
5	2-a	C ₁₀ H ₆ NOCl	0.63	4 hours
6	2-b	C ₁₁ H ₈ NOCl	0.65	6-8 hours
7	2-c	C ₁₀ H ₅ N ₂ O ₃ Cl	0.70	4 hours
8	2-d	C ₁₀ H ₅ NOCl ₂	0.74	4 hours
9	3-a	C ₁₈ H ₁₅ N ₂ Cl	0.72	1-2 hours
10	3-b	C ₁₇ H ₁₂ N ₃ O ₂ Cl	0.69	1-2 hours
11	3-c	C ₁₇ H ₁₄ N ₃ Cl	0.68	1-2 hours
12	3-d	C ₁₇ H ₁₃ N ₂ Cl	0.71	1-2 hours

3. Results and Discussion

Different biological active heterocycles (table-I) have been successfully synthesized in higher yield using Vilsmeier-Haack reagent. In the present study on formylation (Aromatic Electrophilic Substitution) was found to be better support on basis of the yield of the product. The substituted quinolines and its derivatives was synthesized by Vilsmeier-Haack reagent in good yield (60-80%). It was prepared by refluxing 2-chloro-3-formyl quinoline and substituted aromatic aniline for 2-3 hours using air condenser. The required substituted 2-chloro-3-formyl quinoline was prepared by two stage procedure. Thus, in that first the starting material which is substituted acetanilide is prepared by the reaction between (various) substituted aniline and acetic anhydride in presence of acetic acid, then this reaction mixture reflux for about 60 minutes at 80-90 °C on air condenser. Then next stage consisting the Vilsmeier cyclisation of acetanilide, in which POCl₃ (phosphoryl chloride) adding to the substrate (acetanilide) in DMF (N,N-dimethyl formamide) at 0-5 °C then reaction mixture refluxed for about 4 hours at 80- 90 °C to get 2-chloro-3-formyl quinoline. Only methyl group substituted quinolines requires 4-10 hours heating. The structure of all synthesized compounds were confirmed by IR,

¹H NMR spectra and melting point. And all the reactions are monitored by Thin Layer Chromatography. In which we checked the progress of reaction mixture and we found that Rf values in between 0-1. The Rf value is used to determine the value of the movement of material along the TLC plate. The Rf value is equal to the distance travelled by the substrate (material) spotted on TLC plate divided by the distance travelled by solvent, its value is always between zero and one. A TLC analysis might be summarized something like, Using silica gel plate and ethyl acetate as the development solvent and after visualization we observed that the less polar compound has travelled further and more polar compound has travelled less far. The following table showing Rf value of all synthesized products.

4. Conclusion

Different heterocyclic compounds have been synthesized by simple, rapid and environmentally benign method using Vilsmeier reagent. i.e (DMF + POCl₃) phosphorous oxy chloride and dimethyl formamide. Synthesized heterocycles possess good yield. All synthesized compounds were characterized by TLC, Melting Point. A efficient and convenient method for the preparation of substituted

quinoline and its derivatives with good yield has been developed. The reaction / method successfully expands the synthetic scope of the multi-component Vilsmeier-Haack reaction. Further, the present procedure may be readily used to large scale synthesis and generation of the substituted quinolines, The physical data, such as molecular formula, molecular weight, melting point, Yield, % Yield and elemental analysis were shown in table no-I. And Rf value and time period of reaction are shown in table no-II

5. Acknowledgement

The authors are thankful to UGC. This work was financially supported by UGC western regional office Pune. And the authors are also thankful to the principal of Shivaji College, Udgir and Head department of chemistry for providing facilities in the department.

6. References

1. Meth-cohn O, Narine B. Versatile new synthesis of quinolines, thienopyridines and related fused pyridines. (Tetrahedron lett.) 1978; 19:2045-2048
2. Meth-cohn O, Narine B, Tarnowski B. A Versatile new synthesis of quinolines and related fused pyridines. The synthesis of 2-chloroquinoline-3- carbaldehydes. J. Chem. soc; perkins Trans-1- 1981, 1520-1530.
3. Ambika Srivastava, Singh KM. Indian journal of chemistry 2005, 44B:1868-1875.
4. Bandgar BP, Makane SS. synlett, 2003, 262.
5. Venugopal M, Perumal PT, Rajaduraj S. Tetrahedron let.1974, 913.
6. Judit Toth, GaborBlasko, AndrasDancso. Synthetic communication, - (Taylor and francis). 2006, 3653581-3589.
7. Yogeshmurti, Sujeet K Gupta, Pathak. Derpharma chemical 2010, 2(4):271-277.
8. Sekar M, Prasad RKJ. J. Chem, Technol, Biotechnol, 1998, 32-52.
9. Sheeja Devi K *et al.* / Asian. J. of pharmaceutical and medicinal chemistry. 2013; 1(1):39-47.
10. Bakr F, Abdel-Wahab, Rizk E. khidre. ARKIVOC 2012; (1):211-276.
11. Bakr F, Abdel-Wahab, khidre. (Review article) (Hindwai) Journal of chemistry 2013; ID851297:13.
12. Hemant Badwik, Vanktraman S, Deepa Thakur. (A Jr c) Asian journal research chem 2011, 4(6).
13. Jaya kumar swamy BHM, Praveen Y. Pramod; Journal of Pharmacy Research 2012; 5(5):2735-2737.
14. Ameya A Chavan, Nandini R. Pai. Molecules 2007; 12:2467-2477.
15. Rajput AP, Girase P. (review article) International journal of Pharma-ceutical, chemical and biological sciences. (IJPCBS) 2012; 3(1):25-43.
16. Badgujar DM, Talawar MB, Asthana SN, Maulikar PP. J. of scientific and industrial research 2008, 67.