



P-ISSN: 2349-8528
E-ISSN: 2321-4902
IJCS 2017; 5(4): 1851-1853
© 2017 IJCS
Received: 28-05-2017
Accepted: 30-06-2017

Bollaveni Sathish Kumar
Department of agronomy,
collage of agriculture, Sardar
Vallabhbhai Patel University of
Agriculture & Technology,
Meerut, Uttar Pradesh, India

Raj Vir Singh
Proffeser, Department of
agronomy, collage of agriculture,
Sardar Vallabhbhai Patel
University of Agriculture &
Technology, Meerut Uttar
Pradesh, India

Adesh Singh
Assistant Proffeser, Department
of agronomy, collage of
agriculture, Sardar Vallabhbhai
Patel University of Agriculture
& Technology, Meerut Uttar
Pradesh, India

BP Dhyani
Proffeser, Department of Soil
Science, collage of agriculture,
Sardar Vallabhbhai Patel
University of Agriculture &
Technology, Meerut Uttar
Pradesh, India

Pooran Chand
Proffeser, Department of
Genetics and Plant Breeding,
collage of agriculture, Sardar
Vallabhbhai Patel University of
Agriculture & Technology,
Meerut Uttar Pradesh, India

Correspondence

Bollaveni Sathish Kumar
Department of agronomy,
collage of agriculture, Sardar
Vallabhbhai Patel University of
Agriculture & Technology,
Meerut, Uttar Pradesh, India

Effect of nitrogen levels and cutting management on growth parameters of fodder oat (*Avena sativa* L.)

Bollaveni Sathish Kumar, Raj Vir Singh, Adesh Singh, BP Dhyani and Pooran Chand

Abstract

A field experiment was conducted during *Rabi* season 2015-16 at Crop Research Centre (Chirauri) of Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.) to evaluate the effect of nitrogen levels and cutting management on performance of fodder oat and chemical properties of soil. The area lie at a latitude of 29° 40' North and longitude of 77° 42' East with an elevation of 237 meters above mean sea level. The soil of the experimental field was well drained, sandy loam in texture and slightly alkaline in reaction. It was medium in available nitrogen and phosphorus but high in available potassium with an electrical conductivity (1:2, soil: water suspension) of 1.6 dS/m. The treatments comprised 5 nitrogen levels (0, 40, 80, 120 and 160 kg ha⁻¹) and 2 cutting management (single at 50% flowering and double at 60 DAS and 50% flowering), replicated 4 times in a factorial randomized block design. The data on growth, physiology, green forage yield and quality and its contributing traits were calculated on net plot area basis (16 m²), whereas content and uptake in straw along with soil available nutrients and production economic, were recorded as per the standard procedure. The results indicated that growth parameters (plant height, number of tillers, leaf area index and dry matter accumulation) at all the stages were significantly superior at 160 kg nitrogen ha⁻¹. Likewise, single cut resulted into higher values of above mentioned parameters than their respective counterparts like double cutting (60DAS and 50 % flowering).

Keywords: growth parameters, *Avena sativa* L, Crop Research Centre

Introduction

Livestock population is the largest in India comprising 182.50 million cattle, among these, 61.30 million buffaloes, 76.65 million goats, 41.30 million sheep, 10.0 million pigs and 3.04 million other animals. (Jat *et al.* 2014) [4]. India is having the largest livestock population, 15% of the world's livestock population (Neelar, 2011) [7]. Livestock contributing 7% to national GDP and source of employment and ultimate livelihood for 70% population in rural areas. Deficiency in feed and fodder has been identified as one of the major components in achieving the desired level of livestock production. At present, the country faces a net deficit of 63% green fodder, 24% dry crop residues and 64% feeds (Kumar *et al.*, 2012) [6]. The productivity of our livestock often remains low due to inadequate and nutritionally unbalanced supply of feed and fodder. Half of the total losses in livestock productivity are attributed to the inadequacy in supply of feed and fodder.

Deficiency of green fodder will be about 64.9% and for dry fodders up to 24.9% in 2025 A.D. Under such circumstances the only way to bridge the gap between demand and supply of the fodder is to ameliorate the forage resources. Among forages, oat (*Avena sativa* L.) is an important winter feed and forage crop. Owing to its excellent growth habit, better regeneration capacity and good quality forage, it is a promising forage crop. Oat being rich in carbohydrates, vitamins, minerals and total digestible nutrients, is widely used as supplement, hay and silage.

Oat (*Avena sativa* L.) is the important cereal (*Gramineae*) and forage crop, grown during *rabi* season and is next to berseem in nutritive value. It is also rich in energy, protein, vitamin B, phosphorous and iron (Tiwana *et al.* 2008) [10]. It is cultivated in an area of 1021 million ha with an annual production of 233 million tons in the world (Anonymous, 2009) [1]. The total area covered under oat cultivation in India is about 1.0 million ha with 350-500 q ha⁻¹ green fodder productivity (IGFRI, 2011) [3]. In India, it is grown in Punjab, Haryana, Jammu &

Kashmir Himachal Pradesh, Uttar Pradesh, Madhya Pradesh, Rajasthan, Maharashtra and West Bengal. The crop occupies maximum area in Uttar Pradesh (34%) followed by Punjab (20%), Bihar (16%), Haryana (9%) and Madhya Pradesh (6%).

It is important winter crop in many parts of the world and is grown as multipurpose crop for grain, pasture, forage or as a rotation crop. Oat plant has excellent growth habit, quick recovery after cutting and good quality herbage. It is a palatable, succulent and nutritious crop. The protein quality of oat is excellent. Oat requires a long cool season for its growth; therefore, it is successfully grown in the northern plains and hilly areas of the country. Oat is mostly fed as green but surplus is converted into silage or hay to use during fodder deficit periods. Oat as a forage crop, has the advantage of being winter hardy and serves as catch crop.

There are several factors, which affect the productivity and quality of forage oat. Nitrogen is a one of the major component to influence the forage growth, yield and quality. Nitrogen play the vital role in the growth of fodder through the impact on cell elongation, cell division and inter-nodal expansion, it also play a major role in early establishment of the crop. Nitrogen is useful for the improvement of the leaf area by synthesis of enzymes and chlorophyll and also improves the leaf weight. Nitrogen improves the fodder yield through enhancement of growth parameters like plant height, number of tillers, leaf area index, number of leaves, leaf: stem ratio and dry matter accumulation. It is an essential part of protein and is a constituent of physiologically important compounds like nucleotides, vitamins, enzymes and hormones that promotes growth and development in crop plants (Kumar *et al.*, 2001)^[5] and also improve the meristematic activity, it is useful for absorbing of nutrients from the soil efficiently and enhance the protein content of the crop through improvement in synthesis of carbohydrates. Forage oat especially multi-cut oat cultivars are heavy feeder of nutrients and remove large amount of nutrients from the soil. Nitrogen availability to the plant directly influence the forage yield, it is the reason to provide the nitrogen with the split applications. Split application of nitrogen improves the availability of the nutrient to the crop and improves the nitrogen use efficiency. In general farmers use the nitrogen 120 kg ha⁻¹ with improper doses in oat crop but nitrogen @160 kg ha⁻¹ with split applications gave the higher yields than farmer practices (Chouhan *et al.*, 2014)^[2].

Cutting management is the one of the important factor to influence the fodder crop growth, yield and quality. In general cutting management may followed in fodder crops for higher yields. Cutting exhibits the effect on nutrient and natural resource utilization by the crop. As compared to single cut multi cut crops absorb more nutrients, which directly influence the nitrogen content, protein content and other quality parameters of the crop. Cutting is one of the main factor to influence the green and dry forage yield (Patel *et al.*,

2013)^[8], because it plays major role in biomass synthesis. In oat crop commonly gave two cuttings are at different stages, but 1st cut at 60 DAS and 2nd cut at 50% flowering gave the better growth and yield (Sharma *et al.* 2001)^[9].

Materials and Methods

The field experiment was conducted at Crop Research Centre of Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (U.P.) during *rabi* season 2015-16. The climate of this region is sub-tropical and semi-arid and climate characterized with summers and extremely cold winters. The mean maximum temperature of this region is about 43 °C to 45 °C is not uncommon during summer while very low temperature (1-2 °C) accompanied by frost may be experienced in December-January. The winters are cool; frost generally occurs towards the end of December and may continue till the end of January. The monsoon generally begins during the third week of June and desists by the end of September. The total precipitation and its distribution in this region various largely about 75 to 85% of it's received during July to September and few showers are also a common feature during the month of December to January and in late spring season. Mean weekly meteorological data for the season during experimentation period (2015-2016) based on observations collected at meteorological observatory of SVPUA&T, Modipuram, Meerut. Total precipitation of 211.30 mm was received during the experimentation period. The maximum temperature being 32.71°C was recorded in 13th standard week. While, the minimum temperature was 8.57°C in 51th standard week and the Relative Humidity ranged from 45.4 to 66.2 % recorded, respectively in the morning. Ten treatment combinations comprising of five levels of nitrogen viz., 0 kg ha⁻¹ N(N0), 40 kg ha⁻¹ N (N1), 80 kg ha⁻¹ N (N2), 120 kg ha⁻¹ N (N3) and 160 kg ha⁻¹ and two cuttings, single cut at 50% flowering stage and first cut at 60 DAS and second cut at 50% flowering. Oat seed were drilled by adopting a spacing of 25 cm. Kent cultivar was used in the study.

Five plants were tagged randomly from the sampling area for recording height (cm) at 25, 55, 85 DAS and at 115 DAS. The height was measure from ground surface to the base of fully opened leaf before the ear emergence and up to the base of ear heading. Total number of tillers 0.5 meter⁻¹ row length was recorded from five places (marked with sticks) in each plot at 25, 55, 85 DAS and at 115 DAS. The leaf area index was calculated at 25, 55, 85 DAS and at 115 DAS. Plants were harvested from gram per plant from the one row per plot at 25, 55, 85 DAS and at 115 DAS. The plants were sun dried separately and then oven dried at 72 ± 0.5°C till the constant weight is obtained. The dry matter accumulation was expressed in g plant⁻¹.

Results and Discussion

Table 1: Effect of nitrogen levels and cutting management on plant height (cm) of fodder oat at various growth stages

Treatment	Plant height (cm)			
	25 DAS	55 DAS	85 DAS	115 DAS
Nitrogen levels (kg ha ⁻¹)				
N - 0	25.6	61.5	71.1	112.3
N - 40	29.4	64.9	78.6	119.1
N - 80	30.4	71.2	82.5	130.5
N - 120	32.5	75.2	88.2	143.2
N - 160	34.9	81.4	94.0	152.5
S.Em (±)	0.5	0.76	0.79	0.84

CD at 5%	1.45	2.21	2.29	2.45
Cutting management				
50% flowering	30.6	70.9	111.0	147.2
At 60 DAS and 50% flowering	30.5	70.8	54.8	115.9
S.Em (\pm)	0.31	0.48	0.50	0.53
CD at 5%	NS	NS	1.45	1.55

Table 2: Effect of nitrogen levels and cutting management on number of tillers 0.5 per metre row length at various growth stages

No. of Tillers				
Treatment	25 das	55 das	85 das	115 das
Nitrogen levels (kg ha ⁻¹)				
N - 0	25.3	50.4	52.6	32.8
N - 40	27.3	53.1	59.3	36.4
N -80	29.4	60.4	64.2	39.7
N -120	33.5	66.2	68.8	42.3
N - 160	35.6	72.7	74.5	45.0
S.Em (\pm)	0.30	0.75	0.58	0.35
CD at 5%	0.89	2.18	1.69	1.03
Cutting management				
50% flowering	30.3	60.6	65.3	43.3
At 60 DAS and 50% flowering	30.1	60.5	62.5	34.8
S.Em (\pm)	0.19	0.47	0.36	0.22
CD at 5%	NS	NS	1.07	0.65

Table 3: Effect of nitrogen levels and cutting management on leaf area index at various growth stages

Leaf area index				
Treatment	25 das	55 das	85 das	115 das
Nitrogen levels (kg ha ⁻¹)				
N - 0	0.44	2.14	1.98	3.05
N - 40	0.53	2.67	2.31	3.35
N -80	0.67	3.43	2.93	4.18
N -120	0.91	4.10	3.62	6.56
N - 160	1.04	4.22	4.20	9.07
S.Em (\pm)	0.00	0.09	0.02	0.04
CD at 5%	0.02	0.19	0.06	0.12
Cutting management				
50% flowering	0.72	3.13	4.09	5.20
At 60 DAS and 50% flowering	0.72	3.31	1.93	5.19
S.Em (\pm)	0.00	0.02	0.01	0.02
CD at 5%	NS	NS	0.03	0.08

Table 4: Effect of nitrogen levels and cutting management on dry matter accumulation (g plant⁻¹) at various growth stages.

Dry matter accumulation (g plant ⁻¹)				
Treatment	25 das	55 das	85 das	115 das
Nitrogen levels (kg ha ⁻¹)				
N - 0	7.42	11.04	20.62	58.20
N - 40	7.64	11.38	26.83	62.50
N -80	7.93	11.87	35.15	70.50
N -120	8.27	12.42	44.48	75.87
N - 160	8.56	13.07	50.75	81.00
S.Em (\pm)	0.01	0.01	0.73	0.76
CD at 5%	0.03	0.04	2.12	2.23
Cutting management				
50% flowering	7.97	11.96	51.30	73.66
At 60 DAS and 50% flowering	7.96	11.95	20.62	65.60
S.Em (\pm)	0.002	0.006	0.46	0.48
CD at 5%	NS	NS	1.34	1.41

Nitrogen levels exhibited a significant effect on all growth Parameters (plant height, number of tillers, leaf area index and dry matter accumulation) at all the crop growth stages. While, in case of cutting management effect was noticed only

after 85 DAS. The related data are represented in above tables.

Among the all nitrogen levels, 160 kg N ha⁻¹ accumulated significantly more plant height, number of tillers, leaf area index and dry matter than lower doses. Growth parameters are at every growth stage increased significantly with increasing in N levels. Cutting management also exhibited significant effect on growth parameters after 85 DAS. Single cut recorded significantly higher results than double cut at 85 and 115 DAS.

Conclusion

Plant height, number of tillers, leaf area index and dry matter accumulation increased with crop age only upto 1st cut (60 DAS), and thereafter, decreased progressively under all levels of nitrogen. At all the stages, application of nitrogen 160 kg ha⁻¹ recorded significantly higher results over 0, 40, 80 and 120 kg ha⁻¹.

References

1. Anonymou Available in: <[http://www.fao.org/statistical database](http://www.fao.org/statistical_database)> [accessed 24 5 2016], 2009.
2. Chouhan M, Gudadhe NN, Dinesh Kumar, Kumawat AK, Kumar R. Transplanting dates and nitrogen levels influences on growth, yield attributes, and yield of summer pearl millet. The Biasean. 2014; 10(3):1295-1298.
3. IGFRI. Vision 2030: In: Pandey, KC, Roy, AK. (ed) Forage Crops Varieties. IGFRI, Jhanshi, India, 2011, 23-27.
4. Jat RK, Patel AG, Shviran A, Bijarnia AL. Bhunwal V. Response of oat (*Avena sativa* L.) on quality and economics to nitrogen and phosphorus levels under North Gujarat agro-climatic conditions. Journal of Crop and Weed. 2014; 10(2):492-494.
5. Kumar A, Jaiswal RS, Verma ML, Joshi YP. Effect of nitrogen level and cutting management on yield and quality of different varieties of oat fodder. Indian Journal of Animal Nutrition. 2001; 18(3):262-266.
6. Kumar S, Agrawal RK, Dixit AK, Rai AK, Singh JB, Rai SK. Forage Production Technology for Arable Lands. *Technology Bulletin*. 2012; 39(9):255-260.
7. Neelar A. Response of oat genotypes to seed rate and nitrogen levels on forage yield and quality under irrigation. M.Sc. (Ag) thesis submitted to Department of Agronomy, University of Agricultural Sciences, Dharwad, India, 2011.
8. Patel TU, Arvadia MK, Patel DD, Thanki JD, Patel HM. Response of oat (*Avena sativa* L.) to cutting management and times of nitrogen application. *Crop Reaserch and Research on Crops*. 2013; 14(3):902-906.
9. Sharma SK, Bhunia SR. Response of oat (*Avena sativa* L.) to cutting management, method of sowing and nitrogen. Indian Journal of Agronomy. 2001; 46(3):563-567.
10. Tiwana US, Puri KP, Chaudhary DP. Fodder productivity quality of multicut oat grown pure and in mixture with different seed rates of sarson. *Forage Research*. 2008; 33(4):224-226.