



P-ISSN: 2349-8528

E-ISSN: 2321-4902

IJCS 2017; 5(5): 306-308

© 2017 IJCS

Received: 15-05-2017

Accepted: 16-06-2017

Rajendra NasnwaDepartment of Plant Pathology,
COA Bikaner, Rajasthan, India**SL Godara**Professor and head, Department
of Plant Pathology, COA
Bikaner, Rajasthan, India**AK Meena**Assistant Professor, Department
of Plant Pathology, COA
Bikaner, Rajasthan, India**GL Kakraliya**Ph.D. Scholar, Department of
Plant Pathology, SKNCOA
Jobner, Rajasthan, India**Sajjan Choudhary**Ph.D. Scholar, Department of
Plant Pathology, SKNCOA
Jobner, Rajasthan, India**Correspondence****Rajendra Nasnwa**Department of Plant Pathology,
COA Bikaner, Rajasthan, India

International Journal of Chemical Studies

Efficacy of Fungicides against *Drechslera setariae* Causing Leaf Spot Disease of Pearl Millet *Pennisetum glaucum* (L.) *in vitro* condition

Rajendra Nasnwa, SL Godara, AK Meena, GL Kakraliya and Sajjan Choudhary

Abstract

The present investigations were carried out to study the effects of selected fungicides *viz.* Dithane M-45, Tilt, Score, Kavach, Contaf, Companion and Folicur on mycelial growth of *Drechslera setariae in vitro*. The result revealed that all the fungicides *viz.*, Dithane M-45, Tilt, Score, Kavach, Contaf, Companion and Folicur, significantly inhibited the mycelial growth of *D. setariae* at 1, 10, 50, 100 and 200 ppm. Tilt was most inhibitory proved followed Folicur and Tilt to mycelial growth of *D. setariae in vitro*. Out of seven fungicides tested under field conditions to manage leaf spot disease of pearl millet. Tilt (0.2%) and folicur (0.2%) provided effective against this disease. There was significant increase in the grain yield of these two treatments.

Keywords: leaf spot, *drechslera setariae*, severity, fungicides, management

Introduction

Pearl millet [*Pennisetum glaucum* (L.) R. Br.] 2n = 14 locally known as bajra, bari, sajja, combo, ganti or kambam, is an allogamous crop having protogynous nature. It belongs to the family *Poaceae*. It is an important food and forage crop in Africa and Asia and important forage in America. It has great potential because of its suitability to the extreme limits of agriculture. The pearl millet grains are very nutritious and form the staple diet of approximately 10 per cent of the population in India. It has high protein with slightly superior amino acid profile. It is a good source of protein (11.5%), fat (4.1-6.4%), carbohydrate (59.8-78.2%) and also rich good amount of minerals particularly phosphorus and iron (2.8%). India is the largest producer of pearl millet with an annual production of 9.25 million tonnes from an area of 7.0 million hectares with productivity being 1250 kg/ha (Anonymous, 2014-15a) [1].

Leaf spot of pearl millet caused by *Drechslera setariae* is a common foliar disease. *Drechslera setariae* was isolated from the, rotted seeds and infected parts of seedlings. The seed used was obtained from Mysore, India and was found to be infected with *D. setariae*, which has been reported from the United States to be seed-borne in pearl millet, causing seed rot, blight and leaf spotting (Wells & Winstead, 1965 and Wells & Burton, 1967) [16, 15]. The disease has also been noticed in India by Bhowmik (1972) [4] and Balasubramanian (1980) [3].

Materials and Methods

Evaluation of fungicides against of *Drechslera setariae in vitro*

Different fungicides were tested at five different concentrations *viz.*, 1, 10, 50, 100 and 200 ppm in Petri plates. Poisoned food technique (Grover and Moore, 1961) [6] was adopted for management of plant pathogen *in vitro*.

Technical formulation	Trade name	Dose (ppm)
Mancozeb 75 WP	Dithane M-45	1, 10, 50,100,200
Chlorothalonil 75WP	Kavach	1,10, 50,100,200
Hexaconazole 5 EC	Contaf	1,10, 50,100,200
Propiconazole 25 EC	Tilt	1,10, 50,100,200
Difenconazole 25EC	Score	1,10, 50,100,200
Carbendazim 12 WP + Mancozeb 63 WP	Companion	1,10, 50,100,200
Tebuconazole 25 EC	Folicur	1,10, 50,100,200
Control*	-	-

*Without fungicide

~ 306 ~

Poisoned food technique

In this experiment the antifungal activity of various fungicides were tested *in vitro*, against mycelial growth of the pathogen by poisoned food technique at five different concentrations *viz.*, 1, 10, 50, 100 and 200 ppm.

Desired quantities of fungicides were incorporated in potato dextrose agar medium (PDA) aseptically, thoroughly mixed and 20 ml of medium was poured in each Petri plate. These Petri plates were seeded with 5 mm disc taken from periphery of seven-days old culture of the pathogen and incubated at 28 ±1° C. Experiment was replicated thrice. A control was also maintained where no components was added in potato dextrose agar medium (PDA).

Observations were recorded on radial growth of mycelium with two diagonals through the center of colony after seven days. Growth inhibition percentage was calculated by the following formula (Bliss, 1934) [5].

$$\text{Per cent inhibition (I)} = \frac{C - T}{C} \times 100$$

Where,

I = Per cent inhibition of mycelial growth (%).

C= Mycelial growth observed in control (mm).

T= Mycelial growth observed in treatment (mm).

Results and Discussion**Evaluation of fungicides against *Drechslera setariae* in *in vitro***

The efficacy of fungicides *viz.*, Dithane-45, Kavach, Contaf, Tilt, Score, Companion and Folicur on mycelial growth of *Drechslera setariae* was studied *in vitro* using potato dextrose agar medium with five concentrations *viz.*, 1, 10, 50, 100 and 200 ppm. The results given in (Table- 1), revealed that the mycelial growth of *Drechslera setariae* was significantly less in potato dextrose agar medium amended with seven individual fungicides. The results also indicated that the inhibition of mycelial growth increased with the increase in concentration of fungicides (Table- 2), out of seven fungicides tested, Tilt was found most inhibitory to the mycelial growth of *Drechslera setariae* followed by Folicur. Companion also significantly checked the mycelial growth of the fungus, however, it was found less effective as compared to Tilt and Folicur. The inhibition of mycelial growth was relatively less in Companion, Score, Dithane M-45 and Contaf treatments. Kavach was least inhibitory to the mycelial growth of *D. setariae*.

Table 1: The efficacy of fungicides on mycelial growth of *Drechslera setariae*

Fungicides	mycelial growth (mm)					Mean
	1 ppm	10 ppm	50 ppm	100 ppm	200 ppm	
Dithane M-45	38.19	29.49	25.29	21.38	18.33	26.54
Kavach	42.37	33.33	27.29	24.19	20.51	29.54
Contaf	35.15	28.39	24.30	19.06	16.09	24.59
Tilt	10.07	0.00	0.00	0.00	0.00	2.01
Score	30.10	20.69	15.39	10.19	5.11	16.30
Companion	25.06	15.62	13.79	10.09	0.00	12.91
Folicur	15.13	5.07	0.00	0.00	0.00	4.04
Control	90.00	90.00	90.00	90.00	90.00	90.00
Mean	35.76	27.82	24.51	21.86	18.76	24.75
SEm±	0.21	0.19	0.18	0.17	0.15	0.16
CD at 5%	0.63	0.56	0.53	0.49	0.45	0.45

Table 2: The efficacy of fungicides on inhibition on mycelial growth of *Drechslera setariae*

Fungicides	Per cent inhibition of mycelial growth					Mean
	1 ppm	10 ppm	50 ppm	100 ppm	200 ppm	
Dithane M-45	57.57 (49.35)*	63.23 (52.67)	71.90 (57.99)	76.24 (60.82)	79.63 (63.17)	69.71 (56.60)
Kavach	52.93 (46.67)	62.97 (52.51)	69.68 (56.58)	73.12 (58.77)	77.21 (61.48)	67.18 (55.04)
Contaf	60.92 (51.30)	68.46 (55.83)	73.00 (58.69)	78.82 (62.59)	82.12 (64.98)	72.66 (58.47)
Tilt	88.81 (70.45)	100.00 (90.00)	100.00 (90.00)	100.00 (90.00)	100.00 (90.00)	97.76 (81.39)
Score	66.56 (54.67)	77.01 (61.34)	82.90 (65.57)	88.68 (70.33)	94.32 (76.21)	81.89 (64.81)
Companion	72.16 (58.15)	82.64 (65.37)	84.68 (66.95)	88.79 (70.43)	100.00 (90.00)	85.65 (67.73)
Folicur	83.19 (65.79)	94.37 (76.27)	100.00 (90.00)	100.00 (90.00)	100.00 (90.00)	95.51 (77.76)
Mean	68.88 (56.09)	78.38 (62.29)	83.17 (65.77)	86.52 (68.45)	90.47 (72.01)	81.48 (64.51)
	SEm±	CD at 5%				
Fungicide (F)	0.54	1.51				
Concentration (C)	0.40	1.12				
F x C	1.20	3.37				

* Figures given in parenthesis are angular transformed values diseases.

References

1. Anonymous. Ministry of Agriculture, Govt. of India, 2014-2015a.
2. Anonymous. Department of Agriculture, Govt. of Rajasthan, 2014-2015b.
3. Balasubramanian KA. Association of *Drechslera setariae* with downy mildew affected pearl millet. *Curr. Sci.* 1980; 49(6):233-234.
4. Bhowmik TP. *Bipolaris setariae* on two new host in india. *Indian Phytopathol.* 1972; 25:590-591.
5. Bliss CL. The methods of probits. *Science.* 1934; 79:38.
6. Grover RK, Moore JD. Adaptations of *Sclerotinia fructicola* and *Sclerotinia laxa* to grow in higher concentration of fungicides. *Phytopath.* 1961; 51:399-401.
7. Gupta V, Shamas N, Razdan VK, Sharma BC, Sharma R, Kaur K et al. Foliar application of fungicides for management of brown spot disease in rice (*Oryza sativa* L.) caused by *Bipolaris oryzae*. *African J Agril Res.* 2013; 8(25):3303-3309.
8. Hemayatullah, Mahapatra SS. Chemical management of brown spot disease of rice in Odisha. *J Pl. Protec. Environ.* 2014; 11(2):67-70.
9. Hossain I, Dey P, Hossain MZ. Efficacy of Bion, Amistar, Tilt in controlling brown spot and narrow brown spot of rice cv.BR11 (Mukta). *J Bangladesh. Agril. Univ.* 2011; 9(2):201-204.
10. Kiran Kumar AC, Nagaraja A, Raghavendra BT, Ravikumara BM. Evaluation of fungicides against *Drechslera setariae* causing brown leaf spot of foxtail millet [*Setaria italica* (L.)]. *Environ. Ecol.* 2013; 31(2A):801-803.
11. Lore JS, Thind TS, Hunjan MS, Goel RK. Performance of different fungicides against multiple diseases of rice. *Indian Phytopath.* 2007; 60(3):296-301.
12. Mac Neal FH, Konzak CF, Smith EP, Tats WS, Russell TS. A uniform system for recording and processing cereal research data USDA. D. C., ARS, Washington. *Agric. Res. Serv.* 1971; 42:34-121.
13. Sharma JR, Basuchoudhary KC. Assesment of seed mycoflora of pearl millet and their control. *Indian Phytopath.* 1975; 28:288-390.
14. Sunder S, Singh R, Dodan DS, Mehla DS. Effect of different nitrogen levels on brown spot (*Drechslera oryzea*) of rice and its management through host resistance and fungicides. *Pl. Dis. Res.* 2004; 20(2):111-114.
15. Wells HD, Burton GW. *Helminthosporium setariae* on pearl millet, *Pennisetum typhoides*, as affected by age of host differences. *Crop Sci.* 1967; 7:621-622.
16. Wells HD, Winstead EE. Seed borne fungi in Georgia grown and western grown pearl millet seed on safe in Georgia during 1960. *Pl. Dis. Rep.* 1965; 49:487-489.