



P-ISSN: 2349-8528
E-ISSN: 2321-4902
IJCS 2017; 5(5): 635-640
© 2017 IJCS
Received: 28-07-2017
Accepted: 30-08-2017

Abel M Maharramov
Department of Organic
Chemistry, Baku State
University, Z. Khalilov, 23.
Baku, Azerbaijan.

Afet T Huseynova
Department of Organic
Chemistry, Baku State
University, Z. Khalilov, 23.
Baku, Azerbaijan.

Kerim A. Kuliev
Department of Analytical and
Organic Chemistry, Azerbaijan
State Pedagogical University, U.
Gadjibekov, 68. Baku,
Azerbaijan

Kamala O Isgenderova
Department of Analytical and
Organic Chemistry, Azerbaijan
State Pedagogical University, U.
Gadjibekov, 68. Baku,
Azerbaijan

Yavar C Gasimova
Department of Analytical and
Organic Chemistry, Azerbaijan
State Pedagogical University, U.
Gadjibekov, 68. Baku,
Azerbaijan

Ali Z Zalov
Department of Analytical and
Organic Chemistry, Azerbaijan
State Pedagogical University, U.
Gadjibekov, 68. Baku,
Azerbaijan

Correspondence
Ali Z Zalov
Department of Analytical and
Organic Chemistry, Azerbaijan
State Pedagogical University, U.
Gadjibekov, 68. Baku,
Azerbaijan

Extraction and Spectrophotometric Determination of Copper (II) With 1-(2- Metoxiphenylamin)-3-Metoksiopropanthiol-2

Abel M Maharramov, Afet T Huseynova, Kerim A Kuliev, Kamala O Isgenderova, Yavar C Gasimova and Ali Z Zalov

Abstract

1-(2-Metoxiphenylamin)-3-metoksiopropanthiol-2 (MPAMPT) is proposed as an analytical reagent for the extractive spectrophotometric determination of copper (II). MPAMPT forms blue colored complex with copper (II) in the pH range 5.4–6.8. Beer's law is obeyed in the concentration range up to $16 \mu\text{g mL}^{-1}$. The yellowish Cu (II)– MPAMPT complex shows a maximum absorbance at 605 nm, with molar absorptivity of $4.32 \times 10^4 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$ and Sandell's sensitivity of the complex obtained from Beer's data is 1.48 ng cm^{-2} . The composition of the Cu(II)– MPAMPT complex is found to be 1:2 (M/L). The interference of various cations and anions in the method were studied. Thus the method can be employed for the determination of trace amount of copper (II) in pharmaceutical, food and plant sample.

Keywords: Copper, 1-(2-Metoxiphenylamin)-3-metoksiopropanthiol-2, determination, chromogenic reagent

Introduction

The alloys of copper find extensive use in automobile radiators, heat exchangers, home heating systems and panels for absorbing solar energy. Copper (II) is a biologically active metal, its compounds affect the vital activity of plant and animal organisms. Copper is one of essential elements required for normal human metabolism^[1]. Copper (II) is known to play a significant role in biological systems and also as a pharmaceutical agent^[2]. Its antibacterial properties have been known for thousands of years. Synthetic copper(II) complexes have been reported to act as a potential anticancer and cancer inhibiting agents and a number of copper complexes have been found to be active both *in vitro* and *in vivo*^[3,4], therefore it is necessary to control that the content of copper (II) ions in the environmental objects is within the permissible concentrations.

Hence, it is necessary to seek highly, accurate and selective analytical methods for quantitative determination of copper at trace levels.

Various spectrophotometric methods have been proposed for the determination of copper contents of the various samples including natural waters and pharmaceutical samples^[5]

For the spectrophotometric determination of copper in various sites (in water, alloys and pharmaceutical samples, synthetic mixtures) suggested naphthazarin (5,8-dihydroxy-1,4-naphthoquinone)^[6], 1-phenyl-1-hydrazonyl-2-oximino propane -1, 2- dion^[7], 4-[N, N-(dimethyl) amino] benzaldehyde thiosemicarbazone^[8], 2-(5-bromo-2-oxindolin-3-ylidene) hydrazine carbothioamide as an analytical reagent^[9], 4-(4'-nitrobenzylidene imino)-3-methyl-5-mercapto-1, 2, 4-triazole^[10], 2-hydroxy-3-methoxy benzaldehyde thiosemicarbazone^[11], 2-acetylthiophene thiosemicarbazone^[12], o-hydroxyacetophenone isonicotinoylhydrazone^[13].

Oxyphenolate and dithiophenolate complexes of copper are insoluble in chloroform, while mixed-ligand complexes with hydrophobic amines and aminophenols easily dissolve in various organic solvents^[14-16].

Copper (II) which forms an intense blue coloured complex with 1-(2-Metoxiphenylamin)-3-metoksiopropanthiol-2 (MPAMPT), which is completely extracted into chloroform. This forms the basis of the proposed extractive spectrophotometric method for determination of copper after its extraction in the form of Cu(II)- MPAMPT complex. The method has been successfully applied to the analysis of a large variety of samples with diverse matrices such as pharmaceutical, food and plant samples.

2. Materials and Methods

2.1. Reagents and Apparatus

The stock solution (1mg / ml) of Copper (II) was prepared by dissolving weighed amount of Copper Sulphate (CuSO₄) in doubly distilled deionized water [17]. More dilute standard solutions were prepared from this stock solution as and when required.

Solutions of MPAMPT in chloroform (0.01M) were used.

To create the optimal acidity, 0.1M solutions of KOH and HCl or ammonium acetate buffers were applied. Acetate buffer solution, prepared by mixing of 2 mol x L⁻¹ aqueous solutions of CH₃COOH and NH₄OH.

The stock solution of various metal ions and anions were prepared by dissolving the appropriate metal salts in distilled water or with suitable dilute acids and making up to a known volume. The extractant was purified chloroform.

The absorbance of the extracts was measured using a Shimadzu UV1240 spectrophotometer and KFK 2 photocolormeter (USSR). Glass cells with optical path of 5 or 10 mm were used. pH of aqueous phase was measured using an I-120.2 potentiometer with a glass electrode. Muffle furnace was used for dissolution of the samples. The process of thermolysis of the compounds was studied using derivatograph system «ShimadzuTGA-50H». IR spectra were recorded on a spectrophotometer "Specord M80" (Germany).

2.3. General procedure

2.3.1. General procedure for the determination of Copper

Portions of stock solutions of Copper (II) varying from 0.1 to 1.0 mL with a 0.1-mL step, a 2.0 mL portion of a 0.01 M solution of MPAMPT, and a 2.5 mL portion of a 0.01M solution of Am were placed in to calibrated test tubes with ground-glass stoppers (the volume of the organic phase was 5 mL). The required value of pH was adjusted by adding 0.1M NaOH.

The volume of the aqueous phase was increased to 20 mL using distilled water. In 10 min after the complete separation of the phases, the organic phase was separated from the aqueous phase and the absorbance of the extracts was measured on KFK-2 at room temperature and 590 nm (l=0.5cm).

2.3.2. Procedure for Determination of Cu (II) in Pharmaceuticals samples

Boiling with 10 ml of aqua regia dissolved 0.5gm of Pharmaceuticals samples. The solution was evaporated to dryness and the residue was dissolved in 10 ml of 1M HCl filter, if required and resulting solution was diluted to 100 ml with doubly distilled water. The working solution was prepared by appropriate dilution of stock solution. From an aliquot of this solution 1 ml was analysed for Cu(II) by the procedure as described earlier.

2.3.3. Determination of copper in plant

A portion of beans (10 g) was crushed and dried in a porcelain dish at 105 °C. The dry residue is heated in a muffle furnace at 500°C. The ash was dissolved in diluted (1: 1) HNO₃ and evaporate to moist salts, which are then dissolved in water, filtered into a volumetric flask of 100 ml. The copper content is determined with MPAMPT

2.3.4. Determination of copper in the gelatin

5 grams of gelatin in a porcelain dish soak 50 ml of distilled water for 2-3 hours. By the swollen gelatin was added 25 mL (1: 1) HNO₃ and heated in a boiling water bath for 2 hours. The solution was filtered and neutralized with NH₄OH (1: 1) was transferred into a volumetric flask of 50 ml. In solution, the copper content is determined with MPAMPT as well as dyetyldytyokarbamate.

3. Results and Discussion

The reagent solution in chloroform has a yellowish color. The maximum light absorption is observed at 370 nm. Structure of ligand was confirmed by using IR spectra [18, 19, 20].

IR (KBr): 3410 cm⁻¹ ν (NH), 3045 cm⁻¹ ν(CH), 2565cm⁻¹ ν(SH), 2968 и 2875 cm⁻¹ ν(-CH₃), 1570cm⁻¹ δ(C₆H₅), 1380 cm⁻¹ δ_{as} (-CH₃). The chemical structure of the reagent is shown in Fig.1.




Figure 1. The chemical structure of MPAMPT

Cu(II) reacts with MPAMPT and gives a blue colored complexes. These complexes are soluble in non-polar solvents.

3.1. The choice of the extractant

The extraction of the complex has been tried with several solvents: chloroform, 1, 2-dichloroethane, tetrachloromethane, dichloromethane, benzene, chlorobenzene, toluene, xylol, butanol, isoamyl alcohol, cyclohexane, ethyl acetate, isobutanol, isoamyl acetate and their mixes. Extractibility of complexes was estimated in coefficient of distribution and extent of extraction. Thus basicity of amines has no noticeable impact on conditions and extraction of complexes. Fast division of layers and the maximum value of molar coefficient of absorption were received at extraction of complexes by chloroform. After a single extraction with chloroform, 97,6% of copper was extracted as an colored complex (in a case the dichloroethane and carbontetrachloride was removed 95,7 % of Copper). Further researches were conducted with chloroform. The concentration of Copper in the organic phase was determined with rubenic acid [21, 22] by photometric measurements afterback extraction, while in the aqueous phase it was determined by the difference.

3.2. Extraction as a function of pH

Change in pH affected the complexation of Cu(II)–MPAMPT. Therefore, the absorbance of complex was studied between pH 1 to 10 by using dilute HCl and NaOH solutions. The absorbance values of extracted complex were measured. The maximum absorbance was obtained in the pH range 5.4 to 6.8(fig. 2). Beyond this pH range, the observed absorbance values were lower. Thus further extraction and determination carried out at 6.




Fig 1: Absorbance of mixed-ligand complexes as a function of the pH of the aqueous phase $C_{Cu} = 1.875 \times 10^{-5} M$; $C_{MPAMPT} = 5.0 \times 10^{-4} M$, KFK-2, $\lambda = 590 \text{ nm}$, $l = 0.5 \text{ cm}$

3.3. Absorption spectrum

1-(2-Metoxiphenylamin)-3-metoksipropanthiol-2 forms a sparingly soluble complex with copper (II). The complex can readily be extracted quantitatively into chloroform in the pH range 5.4–6.8 absorption maximum in the visible region at 605 nm (fig. 3). The reagent has a negligibly small It is evident from the spectrum that the blue solution of the complex in chloroform shows an absorbance at the λ_{max} of the complex and, hence, does not interfere with the determination of copper. Thus, further absorbance measurements of the complex were made at 590 nm.




Fig 2: Absorption of complex Cu – MPAMPT $C_{Cu} = 1.875 \times 10^{-5} M$; $C_{MPAMPT} = 5.0 \times 10^{-4} M$, Shimadzu UV1240, $l = 1 \text{ cm}$.

3.4. Effect of reagent concentration and of shaking time

For the formation and extraction of complex, a 10-15-fold excess of complexing reagent is required; for example, the optimal conditions for formation and extraction of these compounds are provided by $5.0 \times 10^{-4} M$ MPAMPT. However it was found that the presence of excess of the reagent solution does not alter the absorbance of the color reaction. Under optimum conditions the absorbance of the complex formed in the aqueous phase after equilibration with toluene

increased initially and then achieved a constant and maximum value for 10–180 s. Therefore, 30 s was selected as the optimum equilibration time for each extraction during further studies and has been used in the proposed procedure. Based upon the above study optimum conditions providing maximum, stable and reproducible absorbance values were selected and incorporated in the proposed procedure. The equilibration time of 3.0 minute is sufficient for the quantitative extraction of Copper. The stability of colour of the Cu(II)- MPAMPT complex with respect to time shows that the absorbance due to extracted species is stable up to 36 hours, after which slight decrease in absorbance is observed.

3.5. Stoichiometry of the complexes and the mechanism of complexation

The stoichiometry of the Cu(II): MPAMPT complex was determined by Starik-Barbanel relative yield method, equilibrium shift method, crossed lines method and Asmus' methods [23]. It shows that the composition of Cu(II): MPAMPT complex is 1:2 (Fig 4).




Fig 4: Determination of the ratio of components by equilibrium shift method for Cu - MPAMPT $C_{Cu} = 1.875 \times 10^{-5} M$; $C_{MPAMPT} = 5.0 \times 10^{-4} M$, pH=6, $\lambda = 590 \text{ nm}$, KFK-2. $l = 1 \text{ cm}$.

The probable structure of the complex was supported by the IR spectra, in which absorption bands in the $3250\text{-}3620 \text{ cm}^{-1}$ with a maximum at 3475 cm^{-1} observed in the spectrum of MPAMPT, says that the -NH group is involved in the formation of the complex (Fig.4). The observed decrease in the intensity, absorption bands in the area 2580 cm^{-1} shows that the -SH groups involved in the formation of coordination bond. The (C-N) bands occur as a sharp peak in the ranges $1421 - 1431 \text{ cm}^{-1}$. Detection of the absorption bands at 1410 cm^{-1} indicates the presence of a coordinated -NH group [18-20]. New bands were observed between $400 - 600 \text{ cm}^{-1}$ region in the complex, which were absent in the spectrum of ligand. The bands between 455 cm^{-1} were assigned to stretching frequencies of $\nu(\text{Cu-S})$ and the band between 575 cm^{-1} have been assigned to the stretching frequencies $\nu(\text{Cu-N})$ respectively.

The iron content in the complexes was determined after their decomposition aqua regia photometrically using phenantroline. The purity of the compound was checked by the elemental analysis. Elemental analysis individually complexes are given in Table 1.

Table 1: Elemental analysis of ligand L and complex Fe-L

Compound	%	C	S	N	Cu
L	Found	58.55	14.10	6.25	-
	Calculated	58.41	14.16	6.19	-
Cu- L	Found	51.21	12.48	5.37	12.48
	Calculated	51.16	12.40	5.42	12.40

Thermogravimetric study of the complex Cu- MPAMPT shown that thermal decomposition of the complex takes place in two stages: at 60-110°C water evaporates, at 430-500°C-decomposed MPAMPT. The final product of the termolysis of the complex is CuO.

The stability constant of complex Cu(II)- MPAMPT was calculated and found to be $\lg\beta = 12.84$ at room temperature. The sizes of equilibrium constant K ecalculated on a formula $\lg K_e = \lg D - \lg [Am]$ were presented in Table 4.

Additional experiments by the Akhmedly's method [24] showed that the complex exists in monomeric form in the organic phase (the obtained coefficient of polymerization γ was equal to 1.08).

In conclusion the analytical parameters pertaining to the proposed method are given in Table 4.

It was found using the Nazarenko method that Cu(II) in the complexes was present in the form of Cu^{2+} . The number of protons replaced by cobalt in one MPAMPT molecule appeared to be one [25, 26].

3.6. Influence of foreign ions

To evaluate the complex applicability for photometric determination of copper, we examined the influence of foreign ions and reagents. The tolerance limit of the ions shows minimum deviation ($\pm 2\%$) in absorbance. Influence of a number of cations and anions on the accuracy of determination of Cu (II) was studied. Experiments were performed according to the recipe, by which established Calibration curves, with the only difference that a solution other than Cu (II) injected a certain amount of the corresponding ions.

The interference of various cations was removed by using suitable masking agents. The ions which show interference in the spectrophotometric determination of Copper were overcome by using appropriate masking agents. The effect of various ions and reagents on the extraction-spectrophotometric determination of 30 mg copper (II) is summarised in Table 2. It can be assumed that large amounts of alkaline ions, alkaline-earth ions, Cl^- , Br^- , SO_3^{2-} , SO_4^{2-} , NO_2^- , NO_3^- and $C_2O_4^{2-}$. Tartrate, citrate, F^- , J^- , CN^- , thiourea interfere determination of Cu(II). Co(II), Ni(II), Fe(II, III), V(IV,V), W(VI), Mo(VI), Ti(IV) and Mn(II) interfere determination of Cu(II). However, the interfering effect of some of these ions can be reduced by masking with oxalate, citrate or EDTA.

Interference of Fe(III) eliminated oxalic acid; Ti(IV) - tiron or sodium fluoride; Hg (II) ion -sulfit; Nb (V) and Ta (V) - oxalic acid, and Mo(VI) and W(VI) - sodium fluoride and oxalic acid. When using a 1% solution of ascorbic acid does not interfere with determination Mn(VII), V(IV), Nb(V), Cr(VI), Mo(VI) и Fe(III). When using 0.01M oxalic acid definition not interfere V(IV), Nb(V), Ta(V), Cr(III), Mo(VI), W(VI) и Fe(III).

Table 2: Influence of interfering ions on the determination of copper (II) with MPAMPT (30.0 μg Cu added)

Ion	Molar excess of the ion	Masking agent	Found Cu, μg	S_r
Co(II)	60		30.1	0,05
Ni(II)	60	Sodium cyanide	29.8	0,03
Fe(II)	35		29.8	0,04
Cd(II)	250		29.6	0,05
Al(III)	200		29.7	0,02
Fe(III)	40	Sodium Fluoride	30.5	0,05
Zr(IV)	50		29.8	0,03
W(VI)	45		29.8	0,04
Hg(II)	38	$Na_2S_2O_3$	30.2	0,05
Ti(IV)	60	Sodium Fluoride	29.8	0,04
V(IV)	30		29.8	0,06
Mo(VI)	20	Citrate	30.5	0,04
Cr(III)	120		30.4	0,04
Nb(V)	50	Sodium Fluoride	30.3	0,03
Ta(V)	50	Sodium Fluoride	29.5	0,04
Pb (II)	25		30.3	0,05
Pd (II)	10		29.2	0,04
Pt(II)	10		30.4	0,05
Ag (I)	25	Potassium iodide	30.2	0,05
UO_2^{2+}	50		29.5	0,06
Bi(III)	50		30.5	0,05
Acetate	110		30.2	0,05
Tartarate	110		30.3	0,05
Sulphate	125		29.9	0,02
Thiourea	25		29.7	0,05
Fluoride	110		30.5	0,06
Thiosulphate	36		30.2	0,04

The proposed method compares favourably with the existing ones (Table 3) and offers the advantages of better simplicity, rapidity, sensitivity and selectivity.

Table 3: Comparative characteristics of the procedures for determining of copper

Reagent	pH (solvent)	λ , nm	$\epsilon \times 10^{-4}$
Dyétyldydyokarbamat [21, 22]	4-11(carbon tetrachloride)	436	1.40
Cuproin [21, 22]	4-7(isoamyl alcohol)	546	0.64
Neokuproin [21, 22]	3-10(isoamyl alcohol)	454	0.79
DTMP+Phen	6.7-7.9 (Chloroform)	630	3.45
DTMP +BPhen	6.6-8.1 (Chloroform)	635	4.37
DTMP +Dip	6.5-7.7 (Chloroform)	629	3.28
MPAMPT	5.4-6.8 Chloroform)	605	4.32

3.7. Beer's law and analytical characteristics

Table 4: Optical characteristics, precision and accuracy of the spectrophotometric determination of Cu (II) with MPAMPT

Parameter	Value
Color	blue
The pH range of education and extraction	3.0-9.0
The pH range of maximum extraction	5.4-6.8
λ_{\max} (nm)	605
Molar absorptivity ($L \cdot mol^{-1} \cdot cm^{-1}$)	$4.32 \cdot 10^4$
Sandell's sensitivity ($ng \cdot cm^{-2}$)	1.48
R, %	97.6
The equation of calibration curves	$0.025+0.041x$
Correlation coefficient	0.9964
$lg k_e$	5.76
Stability constant (β)	14.5
Beer's law range ($\mu g \cdot ml^{-1}$)	0.05-16
Limit of detection (LOD): $ng \cdot mL$	13
Limit of quantification (LOQ): $ng \cdot mL^{-1}$	42

The adherence to Beer's law was studied by measuring the absorbance value of the series of solutions containing

Table 5: Determination of Cu (II) in Pharmaceuticals samples (n=3, P=0.95)

Sample	Method	\bar{X} (mg/tablet)	RSD(%)	$\bar{X} \pm \frac{t_p \cdot S}{\sqrt{n}}$
Zincovit (Apex)	Diethyldithio carbamate	0.50	1.21	0.50 ± 0.007
	MPAMPT	0.51	1.21	0.51 ± 0.007
Multi vitamin capsule (A-Z)	Diethyldithio carbamate	0.88	0.95	0.89 ± 0.009
	MPAMPT	0.93	0.79	0.93 ± 0.008
Revital (Ranbaxy)	Diethyldithio carbamate	0.49	1.27	0.49 ± 0.006
	MPAMPT	0.52	1.15	0.51 ± 0.006
Supradyn	Diethyldithio carbamate	0.98	0.92	0.98 ± 0.009
	MPAMPT	1.03	0.85	1.03 ± 0.009

Supradyn: calcium (phosphate Pantothenate) 51.3 mg, magnesium (phosphate, stearate, oxide) 21.2 mg, iron (carbonate, sulfate) 10 mg, manganese (sulphate) 0.5 mg, phosphorus (phosphate) 23.8 mg, Copper (Sulfate) 1 mg, Zinc (Sulfate) 0.5 mg, molybdenum 0.1 mg.

Multi vitamin capsule A to Z ns (alkem): elemental copper 0.9 mg, elemental manganese 2 mg, elemental selenium 55 mcg, elemental zinc 10 mg.

different concentrations of the metal ion. A linear calibration graph drawn between absorbance and the metal ion concentration indicates that Cu(II) may be determined in the range 0.5-16 $\mu g/ml$ [27]. The equations of the obtained straight lines and some important characteristics concerning the application of the ternary complexes for extractive-spectrophotometric determination of Cu(II) are listed in Table 4.

The proposed method compares favourably with the existing ones (Table 3) and offers the advantages of better simplicity, rapidity, sensitivity and selectivity [27].

In conclusion the analytical parameters pertaining to the proposed method are given in Table 4.

4. Application

The proposed method was successfully applied for the determination of copper from various pharmaceutical, food and in plant sample. The results found to be in good agreement with those obtained by the standard known method (Table 5 and Table 6).

Revital (Ranbaxy): zinc - 10 mg, ferrous fumarate - 17 mg, iodine - 0.1 mg, magnesium - 3 mg, manganese - 0.5 mg, calcium - 75 mg, copper - 0.5 mg, potassium - 2 mg.

Zincovit (Apex): selenium (50 mcg), manganese (0.9 mg), zinc (22 mg), copper (0.5 mg), chromium (chromic) (25 mcg), molybdenum (25 mcg).

Table 6: Determination of copper in food (mg / kg). n = 6, P = 0.95

Sample	Method	\bar{X} $\mu g/kg$	S	S _r	$\bar{X} \pm \frac{t_p \cdot S}{\sqrt{n}}$
Beans	rubeanic acid	6.05	0.304	0.048	6.05 ± 0.30
	MPAMPT	5.93	0.207	0.035	5.93 ± 0.21
Gelatin	rubeanic acid	12.10	0.508	0.042	12.10 ± 0.53
	MPAMPT	11.74	0.305	0.026	11.74 ± 0.32
Wheat bran	rubeanic acid	5.65	0.198	0.035	5.65 ± 0.21
	MPAMPT	5.33	0.150	0.028	5.33 ± 0.15
Rye	rubeanic acid	4.30	0.1892	0.044	4.30 ± 0.198
	MPAMPT	4.15	0.1618	0.039	4.15 ± 0.170

5. Conclusions

The results obtained show that the newly developed method in which the reagent MPAMPT was used, can be effectively used for quantitative extraction and estimation of Cu(II) from aqueous media. Complex of copper(II) with MPAMPT have been investigated by spectrophotometric method. Extraction of complex is maximal at pH 5.4-6.8. The proposed method is quick and requires less volume of organic solvent. The optimal conditions for the formation and extraction of

compound have been found and the ratios of components I the complexes have been determined.

The Beer's law was applicable in the range of 0.5-16 $\mu g/ml$. A simple, rapid and sensitive methods proposed for the determination of trace amounts of copper. The method is very precise, faster and simpler than other methods. The proposed method was successfully applied for the determination of copper from various pharmaceutical, food and in plant sample. The obtained by independent methods similar values

of some of the mentioned above characteristics are the evidence for the correctness of the performed experiments.

6. References

1. Veitía MSI, Dumas F, Morgant G, Sorenson JRJ, Frapart Y, Tomas A. Synthesis, structural analysis and anticonvulsant activity of a ternary Cu(II) mononuclear complex containing 1,10-phenanthroline and the leading antiepileptic drug valproic acid. *Biochimie*. 2009; 9:1286-1293
2. Kannan D, Arumugham MN. Synthesis, characterisation, DNA-binding studies and antimicrobial activity of copper(II) complex with 1,10-phenanthroline, L-tyrosine and urea as ligands. *Int. J. Inorg. Bioinorg. Chem.* 2013; 3:8-15
3. Bales BC, Kodama T, Weledji YN, Pitié M, Meunier B, Greenberg MM. Mechanistic studies on DNA damage by minor groove binding copper-phenanthroline conjugates. *Nucleic Acids Res.* 2005; 33:5371-5379.
4. Fernandes C, Parrilha GL, Lessa JA, Santiago LJM, Kanashiro MM, Boniolo FS *et al.* Synthesis, crystal structure, nuclease and *in vitro* antitumor activities of a new mononuclear copper(II) complex containing a tripodal N3O ligand. *Inorg. Chim. Acta* 2006; 359:3167-3176.
5. Turkoglu O, Soylak M. Spectrophotometric Determination of Copper in Natural Waters and Pharmaceutical Samples with Chloro (phenyl) glyoxime. *Journal of the Chinese Chemical Society*. 2005; 52:575-579
6. Chaisuksant R, Palkawong-na-ayuthaya W, Grudpan K. Spectrophotometric determination of copper in alloys using naphthazarin. *Talanta*, 2000; 53(3, 4):579-585.
7. Tekale P, Tekale S, Lingayat S, Pabrekar PN. Extractive Spectrophotometric Determination of Copper (II) using 1-phenyl-1- hydrazonyl-2-oximino propane –1, 2 – dione. *Science Research Reporter*. 2011; 1(2):83-87.
8. Karthikeyan J, Naik PP, Shetty AN. A rapid extractive spectrophotometric determination of copper(II) in environmental samples, alloys, complexes and pharmaceutical samples using 4-*N,N* (dimethyl)amino] benzaldehydethiosemicarbazone. *Environmental Monitoring and Assessment*. 2011; 176(1):419-426.
9. Madan PU, Barhate VD. Extractive spectrophotometric determination of copper (II) using 2-(5- bromo-2-oxindolin-3-ylidene) hydrazine carbothioamide as an analytical reagent *European Journal of Biomedical and Pharmaceutical sciences*. 2016; 3(4):392-396
10. Shaikh B, Barache UB, Anuse MA, Gaikwad SH. 4-(4'-Nitrobenzylideneimino)-3-methyl-5-mercapto-1, 2, 4-triazole, A New Chromogenic Reagent for Extractive Spectrophotometric Determination of Copper (II) in Pharmaceutical and Alloy Samples *S. Afr. J. Chem.*, 2016; 69:157-165,
11. Patil YK, Lokhande RS, Rana PK. Extractive spectrophotometric determination of copper (II) by using 2-hydroxy-3-methoxy benzaldehyde thiosemicarbazone as an analytical reagent. *World Journal of Pharmacy and Pharmaceutical Sciences* 2016; 5(2):965-970
12. Rao, M Sayaji Prasad, Reddy NBL, Hussain K. Spectrophotometric determination of copper (II) in alloys and edible oils using 2-acetylthiophene thiosemicarbazone. *Indian Journal of Chemistry -Section A (IJC-A)* IJC-A 2006; 45A(07):1659-1662
13. Reddy GT, Kumar Reddy PN, Gangi Reddy NC, A rapid extractive spectrophotometric determination of Cu(II) in biological, geological and pharmaceutical samples using o-hydroxyacetophenone isonicotinoylhydrazone. *Scholars Research Library Der Pharmacia Lettre*, 2015; 7(3):274-286
14. Kuliev KA, Sh. A Mamedova. Extractive spectrophotometric determination of copper (II) by using 2, 6-dithiol-4-ethylphenol as an analytical reagent. *International Journal of Current Research* 2017; 9(02):46576-46582
15. Kuliev KA. Extractive spectrophotometric determination of copper(II) using 2, 6-dithiol-4-methylphenol and heterocyclic diamines. *World Journal of Pharmacy and Pharmaceutical Sciences*. Bulgariya, 2016; 5(11):11-16
16. Magerramov AM, Verdizade NA, Zalov AZ. Study of the reaction of the complexation of copper(II) with 2-hydroxythiolphenol and its derivatives in the presence of hydrophobic amines // *Bulletin of the St. Petersburg University. Series 4: Physics. Chemistry*. 2015; 2(60, 4):374-385
17. Korostelev PP. Preparation of solutions for chemical analysis works. M. Publishing house of Academy of Sciences of the USSR. 1964, 401.
18. Shrayner R, F'yuzon R, Kortin D, Morrill T. *Identifikatsiya organicheskikh soedineniy: Per. s angl. - M.: Mir, 1983. (In Russian)*
19. Nakamoto K, *IK-spectr KR. neorganicheskikh I koordinatsionnikh soedineniy (Infrared and Raman Spectra of Inorganic and Coordination Compounds)*. Moscow: Mir. 1991, 536.
20. Bellami L. *Infrakrasnie spectra slojnikh molecule (The infrared spectra of complex molecules)*. Moscow: Mir. 1991, 592.
21. Umland F. *Kompleksnye soedineniya v analiticheskoi khimii: teoriya i praktika primeneniya [Complex compounds in analytical chemistry: theory and practice of application]*. M., Mir, 1975, 531.
22. Marczenko Z, Baltsejak MK. *Metodi Spectrophotometrii v UF I vidimoy oblasti (Spectrophotometrically in the UV and visible regions in inorganic analysis)*. M. Binom. *Laboratoriya znaniy*, 2007, 711.
23. Bulatov MI, Kalinkin IP. *Prakticheskoe rukovodstvo po fotokolorimetriceskim I spektrofotometriceskim metodam analiza (Practical Guide on Photocolorimetric and Spectrophotometric Methods of Analysis)*, Moscow: Khimiya, 1972.
24. Akhmedly MK, Kly'gin AE, Ivanova LI, Bashirov EA. On the chemistry of interaction of gallium ions with a number of sulphophtaleins. *Zhurnal Neorganicheskoi Khimii*. 1974; 19(8):2007-2012.
25. Nazarenko VA, Biriuk EA. *Issledovanie khimizma reaktsii ionov mnogovalentnykh elementov sorganicheskimi reagentami [Research of chemism of reactions of ions of multivalent elements with organic reagents]*. *Zhurn. analit. khimii*. 1967; 22(1):57. (In Russian)
26. Nazarenko VA. *Vzaimodeistvie ionov mnogovalentnykh elementov s organicheskimi reagentami [Interaction of ions of multivalent elements with organic reagents]*. *Trudy komis. Po analit. khimii AN SSSR [Proc. of the Commission on analytical chemistry of Academy of Sciences of the USSR]*. Moscow, Nauka, 1969; 17:22. (In Russian)].
27. Dorokhova EN, Prokhorova GV. *Analytical chemistry (physical-chemical methods of analysis)*. M: Higher School, 1991, 250.