



P-ISSN: 2349-8528
 E-ISSN: 2321-4902
 IJCS 2017; 5(6): 2150-2152
 © 2017 IJCS
 Received: 23-09-2017
 Accepted: 24-10-2017

MP Ahir
 Department of Floriculture and
 Landscape Architecture, Aspee
 College of Horticulture and
 Forestry, Navsari Agricultural
 University, Navsari, Gujarat,
 India

Alka Singh
 Department of Floriculture and
 Landscape Architecture, Aspee
 College of Horticulture and
 Forestry, Navsari Agricultural
 University, Navsari, Gujarat,
 India

SJ Patil
 Department of Floriculture and
 Landscape Architecture, Aspee
 College of Horticulture and
 Forestry, Navsari Agricultural
 University, Navsari, Gujarat,
 India

Correspondence
MP Ahir
 Department of Floriculture and
 Landscape Architecture, Aspee
 College of Horticulture and
 Forestry, Navsari Agricultural
 University, Navsari, Gujarat,
 India

Response of different salinity levels on growth and yield of tuberose cv Prajwal

MP Ahir, Alka Singh and SJ Patil

Abstract

During 2013-2014 and 2014-2015, tuberose cv Prajwal was irrigated with water of different salinity levels (2.0, 4.0, 6.0, 8.0, 10.0 and 12.0) along with control (best available water) to evaluated for growth and yield parameters at Department of floriculture, Aspee College of Horticulture and Forestry, Navsari Agricultural University, Navsari. Different levels of salinity were significantly reduced growth and yield parameters. The growth parameters exhibited significantly reduced tuberose plant growth above 4.0 dSm⁻¹ salinity level. Flower yield, bulbs per plant, bulblets per plant were prominently decreased above 4.0 dSm⁻¹ salinity level. In case of leaf chlorophyll content was also decreased with increase in salinity level. Based on results, tuberose cv Prajwal was found moderate salinity sensitive flower crop.

Keywords: Salinity levels, growth, yield, sensitive flower crop

Introduction

The commercial floricultural industry includes flower crops, garden plants, potted flowering and foliage plants. A dramatic aspect of floriculture is water consumption: it has been estimated that 100-350 kg of water are needed to produce 1 kg of plant dry matter, but it can vary with species and variety, cultivation system and plant growing season (Fornes *et al.*, 2007) [10]. About half of the fresh water available to support a growing world population is already used for human consumption (Rozema and Flowers, 2008) [14]. However, the majority of water on earth is seawater (98%), with only about 1 % being fresh-water. Ever increasing demands for good quality water for domestic and industrial uses in developing countries like India create scarcity of good quality water for agriculture use. Many developing countries are now facing this situation, especially in arid and semiarid regions where limited water availability is already a severe constraint to development.

India has coastal line of 7516 kms, Gujarat state having large coastal line in country covering about 1660 kms. The misty and enchanting atmosphere of coastal area attracts more and more people to build their hotels and homes at seashore. In saline soils, the presence of excessive salts in the root zone lead to various physiological changes in the plants which ultimately affect the growth and flowering of the plant. Plants that are able to survive in rugged coastal environment must withstand the prevailing winds, tolerate the salt spray and be capable to set their roots in saline conditions. Coastal landscape can be enhancing by using flowering plants, which can tolerate salinity up to some limit. The effort towards utilization of saline soil and water for growing flowering plants mainly aims to make the beauty of seashore landscape even more enchanting. Various irrigation strategies devised can be used for the purpose of controlling salinity with in the threshold limit of plants, through the conjunctive use of saline irrigation water by avoiding salt stress at critical period of their growth.

Material and Methods

The experiment was carried out at Department of floriculture and landscape architecture, Aspee College of Horticulture and Forestry, Navsari Agricultural University, Navsari (Navsari, India 20° 57' N latitude and 72° 54' E longitudes) during the *rabi* season of 2013-14 and 2014-15. The soil of the experimental pots was collected from Regional Horticultural Research Station, Navsari. Corms of tuberose (*Poliantha tuberosa*- L) cv Prajwal were planted in pot having capacity of 20 kg filled with mixture of soil and FYM along with NPK as per recommended dose. Average two-three corms per pot were placed by dibbling in the soil and irrigated immediately with normal water (best available water). Different levels of saline water (2.0, 4.0, 6.0, 8.0, 10.0 and 12.0 dSm⁻¹) were made up with sea water (approx. 55 dSm⁻¹)

mixed with fresh water (1.48 dSm^{-1}). The experiment was laid out in completely randomized design. Each treatment was applied in four separate repetitions. All the plants were initially grown with fresh water up to one month; saline water treatment was given in October, 2013 and October, 2014. In each treatment, irrigation water was checked every time for EC and pH. Appropriate drain holes should be kept for leaching.

The soil samples were drawn at time planting (before experiment) and finally at harvesting. Growth and flowering data were collected at regular interval during both the seasons, including plant height (cm), number of leaves per plant, necrotic area percentage (%), leaf area (cm^2), root length (cm), days taken to flower initiation, number of spike per plant, flower yield, corms per plant, cormels per plant and chlorophyll content. The influence of saline water irrigation on plant growth, flower production, bulb production, chlorophyll content and anthocyanin content were assessed through ANOVA.

Results and Discussion

Growth Parameters

The growth parameters *viz.*, plant height, number of leaves per plant, leaf area and root length of tuberose cv Prajwal were affected significantly due to different levels of salinity of irrigation water. No plant survival was observed in irrigation treatment 12.0 dS^{-1} water. Necrotic area percentage (NAP) was also increased with increase in salinity level of irrigation water and 100 % necrosis was recorded in treatment with 12.0 dS^{-1} saline water. Above 2.0 dS m^{-1} salinity level of irrigation water, growth in tuberose cv Prajwal was prominently decreased.

Reduced plant growth is a common phenomenon when plants are grown under increased salinity and usually expressed as stunted plant growth. The first responses of plants to salinity is a decreased rate of plant growth primarily due to osmotic effect of salt around the roots, which leads to reduction in water supply to plant cells as explained by Blum, (1986) [5]. Further, Wild (1988) [22] and Shannon & Grieve (1999) [15] stated inhibition of root growth and its function owing to high

external salt concentration. Explaining, the mechanism of salt tolerance in plants, due to increasing EC_{iw} probably resulting into limited cell expansion (Munns and Tester, 2008) [16]. Reduction in cell elongation and division in plant cell reduce their final size, resulting in decrease in plant height, number of leaves leaf area and root growth as elucidated by earlier workers (Cabrera, 2003; Cassaniti *et al.*, 2009) [6, 8]. Growth reduction in different ornamental plants due to salinity have been also reported in gladiolus (Ahir and Singh, 2017) [1], in *Nerium oleander* (Banon *et al.*, 2005), in marigold (Valdez-Aguilar *et al.*, 2009) [20], in gladiolus (Haouala and Sahli 2011) [13], in gladiolus and heliconia (Cerquera *et al.*, 2008) and in zinnia (Zivder *et al.*, 2011) [23].

Flowering Parameters

In flowering parameters *viz.* days taken to flower initiation, number of spikes per plant, flower yield, corms per plant and cormels per plant were significantly influenced by salinity levels of irrigation water. Growth and flowering parameters were reduced steeply after irrigation treatment 4.0 dSm^{-1} and above.

Delay in flowering due to the specific mechanism that alter the growth stage of flowering have been known to occur due to multiple stresses (osmotic imbalance, nutritional deficit and cellular toxicity) exerted by salinity (Risse and Shenk, 1990; Stanton *et al.*, 2000) [18, 17]. Besides, reduction in root biomass caused due to salinity has also been indicated as a factor impeding flowering by affecting energetic reserves (Van Zandt and Mopper, 2002) [11]. Saline water irrigation reduced crop growth and production in sensitive species (Volkmar *et al.* 1998) [21] due to negative effects on water and mineral relations, carbon assimilation and biomass partitioning. Crop response to salinity depends on cultivar and growing conditions (e.g. Bass *et al.*, 1995; De Kreij and Van Os, 1989; Sonneveld *et al.*, 1999) [4, 9, 19]. In our work, tuberose plants appeared much moderate sensitive to salinity. The use of saline water irrigation significantly reduced plant growth, flower production and bio-chemical parameters (table-1).

Table 1: Effect of different levels of salinity of irrigation water on plant growth and flowering of tuberose cv Prajwal

Treatments	Growth parameters				Flowering parameters					Chlorophyll content (mg/g FW)
	Plant height (cm)	No. of leaves/plant	Leaf area (cm^2)	Root length (cm)	Days taken to flower initiation	No. of spike/plant	Post harvest life of flowers	No. of bulbs/plant	No. of bulblets/plant	
Salinity 2.0 dSm^{-1}	59.27	30.88	82.97	11.55	86.65	2.78	6.39	5.24	11.39	11.32
Salinity 4.0 dSm^{-1}	55.56	28.29	76.07	11.10	88.63	2.65	5.94	4.99	10.39	10.59
Salinity 6.0 dSm^{-1}	47.03	24.06	64.31	7.50	92.19	2.28	5.31	4.45	8.91	9.19
Salinity 8.0 dSm^{-1}	45.75	23.68	64.05	5.98	93.95	2.25	5.06	4.18	7.32	8.55
Salinity 10.0 dSm^{-1}	41.13	22.41	61.45	4.10	95.30	2.11	4.60	3.98	6.91	7.61
Salinity 12.0 dSm^{-1}	39.02	21.20	60.10	3.98	96.05	2.06	4.49	3.62	6.27	6.88
Best Available Water	62.03	31.64	91.26	12.11	85.11	2.95	6.50	5.31	12.03	13.66
SEM \pm	2.31	1.11	2.97	0.37	2.30	0.11	0.17	0.24	0.19	0.28
CD at 5 %	6.78	3.25	8.72	1.08	6.77	0.33	0.49	0.71	0.55	0.83
CV	9.23	8.50	8.29	9.11	5.05	9.29	6.05	10.63	4.13	5.85

Conclusion

Based on the results of the above experiment, with the data on growth and flowering parameters indicates level of salinity tolerance in tuberose cv Prajwal has been found salt sensitive crop. Above 4.0 dSm^{-1} salinity level growth and flowering was prominently decreased.

References

1. Ahir MP, Alka Singh. Effect of different levels of irrigation water on growth and yield of gladiolus cv.

American Beauty. Trends in Bio. 2017; 10(43):9011-9013.

2. Arnold MA, Lesikar BJ, McDonald GV, Bryan DL, Gross A. Irrigating landscape bedding plants and cut flowers with recycled nursery runoff and constructed wetland treated water. J. Environ. Hort. 2003; 21(2):89-98.

3. Bañón S, Conesa E, Valdés R, Miralles J, Martínez JJ, Sánchez Blanco MJ. Effect of saline irrigation on phytohormone-treated chrysanthemum plants. Acta Hort. 2012; 937:307-312.

4. Bass R, Nijssen HMC, Van Den Berg TJM, Warmenhoven MG. Yield and quality of carnation (*Dianthus caryophyllus* L) and gerbera (*Gerbera jamesonii* L) in a closed nutrient system as affected by sodium chloride. *Sci. Horticulturae*. 1995; 61:273-284.
5. Blum A. Salinity resistance, In: *Plant Breeding for stress environments*, A Blum (Ed.), 1163-1169, CRC Press, Boca Raton. 1986.
6. Cabrera RI. Mineral nutrition, In; Roberts, A. V., Debener, T. and Gudin, S (eds.). *Encyclopedia of rose science*. Academic Press, Oxford, UK. 2003, 573-580.
7. Cerqueira L, Fadigas FDS, Pereira FA, Gloaguen TV, Costa JA. Growth of *Heliconia psittacorum* and *Gladiolus hortulanus* irrigated with treated domestic waste water. *Revista Brasileira de Engenharia*. 2008; 12(6):606-613.
8. Cassaniti C, Leonardi C, Flower TJ. The effect of sodium chloride on ornamental shrubs. *Sci. Hort*. 2009; 122:586-593.
9. De Kreijl C, Van Os PC. Production and quality of gerbera in Rockwool as affected by electrical conductivity of the nutrient solution. In: *Proc. 7th Int. Cong. Soilless Culture*. 1989, 225-264.
10. Fromes F, Belda RM, Carrion C, Noguera V, Garcia-Agustin, Abad M. Pre-conditioning ornamental plants to draught by means of saline water irrigation as related to salinity tolerance. *Scientia Horticulturae*. 2007; 113:52-59.
11. Van Zandt PA, Mopper S. Delayed and carryover effects of salinity on flowering in *Iris hexagona*. *American J. Botany*. 2002; 89:1847-1851.
12. Grime JP. *Plant Strategies and Vegetation Process*, Wiley, New York. 1979.
13. Haouala F, Sahli L. NaCl effects on growth, flowering and bulbing of gladiolus (*Gladiolus grandiflorus* hort.). *Revue Suisse de viticulture arboriculture horticulture*. 2011; 43(6):378-383.
14. Rozema J, Flowers TJ. Crops for Stalinized world. *Science*. 2008; 322:1478-1480.
15. Shannon MC, Grieve CM. Tolerance of vegetable crops to salinity, *Scientia Horticulturae*. 1999; 78:5-38.
16. Munns R, Tester M. Mechanism of salinity tolerance, *The Annual Review of Plant Biology*. 2008; 59:651-681.
17. Stanton ML, Roy BA, Thiede DA. Evolution in stressful environments. I. Phenotypic variability, phenotypic selection and response to selection in five distinct environmental stresses. *Evolution*. 2000; 54:93-111.
18. Rise I, Shank M. Influence of Cl^- , Na^+ and SO_4^{2-} in irrigation water on the growth of azaleas. *Gartenbauwissenschaft*. 1990; 55:252-258.
19. Sonneveld C, Baas R, Nijssen HMC, Hoog J. Salt tolerance of flower crops grown in soilless culture. *J. Plant Nutrition*. 1999; 22(6):1033-1048.
20. Valdez-Aguilar LA, Grieve CM, Poss JA. Salinity and alkaline pH of irrigation water affect marigold plants. I. Growth and shoot dry mass partitioning. *Hort. Sci*. 2009a; 44:1719-1725.
21. Volkmar KM, Hu Y, Steppihn H. Response physiologique des plantes a la salinite: Mise au point bibliographique. *Can. J. Plant Sci*. 1998; 78:19-27.
22. Wild A. *Russels's soil condition and plant growth*. 11th Edn. Harlow, Longman. 1988.
23. Zivder S, Khaleghi E, Dehkordi FS. Effect of salinity and temperature on seed germination indices of *Zinnia elegans* L. *J. Applied Horti*. 2011; 13(1):48-51.