



P-ISSN: 2349-8528
E-ISSN: 2321-4902
IJCS 2018; 6(2): 08-11
© 2018 IJCS
Received: 03-01-2018
Accepted: 04-02-2018

Vinod Kumar Prajapati

Department of Soil Science, Sam Higginbottom Institute of Agriculture, Technology & Sciences (Deemed-to-be-University), Allahabad, Uttar Pradesh, India

Narendra Swaroop

Department of Soil Science, Sam Higginbottom Institute of Agriculture, Technology & Sciences (Deemed-to-be-University), Allahabad, Uttar Pradesh, India

Ashish Masih

Department of Soil Science, Sam Higginbottom Institute of Agriculture, Technology & Sciences (Deemed-to-be-University), Allahabad, Uttar Pradesh, India

Reena Lakra

Department of Soil Science, Sam Higginbottom Institute of Agriculture, Technology & Sciences (Deemed-to-be-University), Allahabad, Uttar Pradesh, India

Correspondence**Vinod Kumar Prajapati**

Department of Soil Science, Sam Higginbottom Institute of Agriculture, Technology & Sciences (Deemed-to-be-University), Allahabad, Uttar Pradesh, India

Effect of different levels of NPK and vermicompost on chemical properties of maize [*Zea Mays* (L.)] Cv. MM2255

Vinod Kumar Prajapati, Narendra Swaroop, Ashish Masih and Reena Lakra

Abstract

Field experiments were conducted during 2015-2016 on clay loam, sandy loam soil to evaluate "to study the effect of different levels of NPK and Vermicompost on physico-chemical properties of soil" comparison to inorganic fertilizers nitrogen, phosphorous, potassium. Vermicompost had increased the pore space, reduced particle and bulk density, increased water holding capacity, cation exchange capacity, reduced pH and electrical conductivity, increased organic carbon content, available nitrogen, phosphorous, potassium and microbial population and activity in all the soil types, particularly clay loam soil. On the contrary, the application of inorganic fertilizers has resulted in reduced porosity, compaction of soil, reduced organic carbon and reduced microbial activity. For soil amendment and soil quality improvement. Observed that, the application of NPK with Vermicompost were excellent source to fertilization than fertilizers only in soil.

Keywords: soil properties, nutrients sources, maize, NPK and vermicompost content, etc.

Introduction

Maize (*Zea mays* L.) belongs to Gramineae family maize is considered as the native to the Central America & Mexico where many diverse types of maize are found Rai (2006) [17]. Maize is one of the most important cereal crops in the world agriculture economy both as food for man and feed for animal. It is a miracle crop. It has very high yield potential. There is no cereal on the earth which has so immense potentiality and that is why it is called "queen of cereals" maize is grown in almost all the states of India. Maize grain contains about 10% protein, 4% oil, 70% carbohydrate 2.3% crude fiber, 10.4% aluminizes, 1.4% ash. Maize protein 'Zein' is in tryptophan and lysine two essential amino acids Singh *et al.*, (2007) [20]. Excessive use of chemical fertilizers, decline in soil and food quality due to loss of soil organic matter is the main characteristic of the conventional farming systems which are more pronounced in arid and semi-arid areas (Singh *et al.* (2007) [20]. Melero *et al.* (2008) [14]. & Liu *et al.* (2009) [12]. Increasing public awareness of the negative environment impacts, growing consumer demand for healthier products and criticism of high input production systems. Alternate agricultural practices such as organic farming, eco-farming, biodynamic farming and traditional farming practices are considered important alternatives to increase soil fertility and soil health. Inorganic farming the application of organic manure especially vermicompost is recommended. It is ecofriendly, non-toxic, consumes low energy input for composting and is a recycled biological product (Lourduraj and Yadav (2005) [13]. Vermicomposts are organic materials broken down by interactions between microorganisms and earthworms in a mesophilic process (up to 25°C), to produce fully stabilized organic soil amendments with low C: N ratios. They have a high and diverse microbial and enzymatic activity, fine particulate structure, good moisture-holding capacity, and contain nutrients such as N, P, K, Ca and Mg which are readily taken up by plants (Lavelle and Martin. (1992) [11]. Prabha *et al.* (2005) [16], Arancon and Edwards (2009) [1]. The application of vermicompost helps to improve and conserve the fertility of soil. Vermicompost imparts a dark colour to the soil and thereby helps to maintain the temperature of soil. Vermicompost is one of the manures used by the farmer in growing crops because of early availability and presence of almost all the nutrients required by plants. The composition of vermicompost is 0.6-1.2% N, 0.13-0.22% P and 0.40-0.75% K Pawar (2007). Nitrogen is a most important element for the synthesis of protoplasm, which is responsible for rapid cell division (plant shape and size).

It increased the production of grain yield in maize as well as it is important for the quality of produce like increase proteins in grain. It increases utilization of P and K to an appreciable extent Singh *et al.* (2003) [19]. Phosphorus its plays a *vital* role in photosynthesis, respiration, energy storage transfer cell division, cell elongation and several other processes in living plants. Phosphorus is also a structural component of the cell constituents and metabolically active compound Ahmad *et al.* (2004). Potassium maintains the cellular organization by regularity the permeability of cellular membrane and keeping the protoplasm in a proper degree of hydration by stabilizing the emulsion of highly colloidal particles. Thus help in maintaining turgor pressure and eliminates water imbalance in plants Singh *et al.* (2003) [19].

Materials and Methods

A field Experiment was conducted on research farm of department of Soil Science, Allahabad School of Agriculture, Sam Higginbottom Institute of Agriculture, Technology & Sciences (Deemed-to-be-University) Allahabad, (U.P.) India. The soil of experimental area falls in order Inceptisol and the experimental field is alluvial in nature. The design applied for statistical analysis was carried out with 3² factorial randomized block design having three factors with three

levels of NPK @ 0, 50, and 100% ha⁻¹, three levels of Vermicompost @ 0, 50 and 100% ha⁻¹ respectively. Treatments were T₀ – (L₀ V₀) @ 0 % NPK ha⁻¹ + 0% Vermicompost ha⁻¹, T₁ – (L₀ V₁) @ 0% NPK ha⁻¹ + 50% Vermicompost ha⁻¹, T₂ – (L₀ V₂) @ 0% NPK ha⁻¹ + 100% Vermicompost ha⁻¹, T₃ – (L₁ V₀) @ 50% NPK ha⁻¹ + 0% Vermicompost ha⁻¹, T₄ – (L₁ V₁) @ 50% NPK ha⁻¹ + 50% Vermicompost ha⁻¹, T₅ – (L₁ V₂) @ 50% NPK ha⁻¹ + 100% Vermicompost ha⁻¹, T₆ – (L₂ V₀) @ 100% NPK ha⁻¹ + 0% Vermicompost ha⁻¹, T₇ – (L₂ V₁) @ 100% NPK ha⁻¹ + 50% Vermicompost ha⁻¹, T₈ – (L₂ V₂) @ 100% NPK ha⁻¹ + 100% Vermicompost ha⁻¹. having the treatments was replicated thrice. The source of inorganic nutrients sources as Urea, SSP, MOP, and organic nutrients sources as Vermicompost respectively. Basal dose of fertilizer was applied in respective plots according to treatment allocation unifurrows opened by about 5cm. depth before sowing seeds in soil at the same time sowing of seeds was sown on well prepared beds in shallow furrows, at the depth of 5cm,.

The soil analysis was done in the laboratory of Soil Science, SHIATS.-DU, and Allahabad with following standard methods,

Results and Discussions

Table 1 (a): Chemical analysis of pre-harvest soil.

Particulars	Rating	Method
1. Soil pH, Soil water Suspension	7.19	Digital pH meter (Jackson 1958)
2. EC (dS m ⁻¹)	0.23	Digital Conductivity meter (Wilcox 1950) [24]
3. Organic carbon (%)	0.70	Walkley and Black (1947) [23]
4. Available N (Kgha ⁻¹)	209.25	Alkaline Permanganate Method (Subbiah and Asija. 1956)
5. Available P (Kgha ⁻¹)	19.61	Calorimetric Method (Olsen <i>et al.</i> , 1954) [15]
6. Available K (Kgha ⁻¹)	132.45	Flame photometric Method (Toth and Price. 1949) [22]

Table 2 (c): Soil Properties post Analysis

Treatment Combination	pH (w/v)	EC (dsm ⁻¹)	O.C (%)	N ₂ O (Kg ha ⁻¹)	P ₂ O ₅ (Kg ha ⁻¹)	K ₂ O (kg ha ⁻¹)
T ₀ =L ₀ V ₀	7.10	0.28	0.60	207.17	16.00	115.98
T ₁ =L ₀ V ₁	7.14	0.40	0.66	209.48	16.61	123.05
T ₂ =L ₀ V ₂	7.29	0.44	0.73	211.49	17.51	127.83
T ₃ =L ₁ V ₀	7.11	0.46	0.68	213.48	19.00	131.11
T ₄ =L ₁ V ₁	7.26	0.43	0.66	217.19	19.59	127.57
T ₅ =L ₁ V ₂	7.40	0.42	0.70	219.22	20.21	134.57
T ₆ =L ₂ V ₀	7.13	0.50	0.69	222.17	21.10	139.85
T ₇ =L ₂ V ₁	7.16	0.48	0.75	225.20	22.87	144.27
T ₈ =L ₂ V ₂	7.11	0.52	0.84	229.48	24.03	147.87
Mean	7.19	0.43	0.70	217.20	19.65	132.45
F- test	NS	NS	NS	S	S	S
S. Em (±)	0.013	0.029	0.015	1.588	0.228	2.240
C.D. at 5%	0.028	0.062	0.032	3.367	0.483	4.749

NS: Non Significant, * significant at 5% and ** Significant at 1%

Chemical properties of soil at 0-15 cm depth.

Ph

The maximum pH (7.40) was found in T₅- [NPK @ 50% RDF + Vermicompost 100%] with (7.10) and the minimum value (7.11) pH was found in treatment T₈ (NPK @ 100% RDF + NPK @ 100% RDF). similar results have also been recorded by Bhattacharya *et al.* (2004) [4], Laxminarayan (2006) [10].

Electrical conductivity

(dSm⁻¹)

The maximum electrical conductivity (0.52) in depth 0 - 15 was found with T₈-[NPK 100% RDF + Vermicompost 100%] on followed by T₆- [NPK 100% RDF + Vermicompost 50%] with (0.50) and the minimum value (0.28) electrical

conductivity was found in treatment T₀ [NPK 50% RDF + Vermicompost 50 %] similar results have also been recorded by Aphale *et al.* (2005) [2].

Organic Carbon (%)

The maximum Carbon (0.84%) in depth 0 - 15 was found with T₈-[NPK 100% RDF + Vermicompost 100%] on followed by T₇- [NPK 100% RDF + Vermicompost 50 %] with (0.75 %) and the minimum value (0.60 %) Carbon was found in treatment T₀ (Control). similar results have also been recorded by Bhattacharya *et al.* (2004) [4].

Available nitrogen

The maximum available nitrogen (kg/ah) (229.15 kg/ha) in

depth 0 - 15 was found with T₈-[NPK 100% RDF + Vermicompost100%] on followed by T₇- [NPK 100% RDF + Vermicompost50 %] with (225.20kg ha⁻¹) and the minimum value (207.17) kg ha⁻¹) available nitrogen was found in treatment T₀ (Control). similar results have also been recorded by Iqbal *et al* (2013)^[6]. Singh (2003)^[19].

Available phosphorus

The maximum available phosphorus kg ha⁻¹ (24.03kg ha⁻¹) in depth 0 - 15 was found with T₈-[NPK 100% RDF + Vermicompost 100%] on followed by T₇- [NPK 100% RDF + Vermicompost 50 %] with (22.87 kg ha⁻¹) and the minimum value (16.00 kg ha⁻¹) available phosphorus was found in treatment T₀ (Control). similar results have also been recorded by, Iqbal *et al* (2013)^[6].

Available Potassium (kg/ha).

The maximum available potassium kg ha⁻¹ (147.87 kg ha⁻¹) in depth 0 - 15 was found in T₈-[NPK 100% RDF + Vermicompost100%] on followed by T₇- [NPK 100% RDF + Vermicompost 50%] with (144.27 kg ha⁻¹) and the minimum value (115.98 kg ha⁻¹) available potassium was found in treatment T₀ (Control.) similar results have also been recorded by Khatkar *et al.* (2009)^[8].

Conclusion

It was concluded from trial that the various levels of different sources in the experiment, the treatment T₈-L₂ V₂[@ 100%NPK+100%Vermicompost.] was found to be the best in increasing Chemical properties of Soil pH (7.19), EC (0.23 dSm⁻¹), Organic carbon (0.70%), N (209.25 Kg ha⁻¹), P (19.61 Kg ha⁻¹), K (132.45 Kg ha⁻¹), were found to be at par than any other treatment combinations. Since the result is based on one year experimental data. Further research may be initiated for the establishment of the above findings.

Acknowledgements

The authors are thankful to Advisor, Department of Soil Science & Agriculture chemistry, Sam Higginbottom Institute of Agriculture, Technology & Sciences (Deemed-to-be-University), Allahabad, U. P. for taking their keep interest and encouragement to carry the research work.

References

1. Arancon QN, Edwards CA. The utilization of vermicompost in Horticulture and Agriculture, 2009.
2. Aphale SL, Stokes TS, Black CR, Taylor IB, Roberts JA. The role of root to shoot signaling in coordinating responses to soil compaction. Proceedings of the 32nd Annual Meeting of the Plant Growth Regulation Society of America, 2005.
3. Black CA. (Ed) Method of soil analysis Vol. 1. Am. Soc Agron. Madison, Wisconsin, V.S.A, 1965.
4. Bhattacharya R, Vedprakash Kundu, Srivastava AK, Gupta HS. Effect of long term manuring on soil organic, bulk density and water retention characteristics under soyabean-wheat cropping sequence in north-western Himalayas journal of the Indian society of soil science. 2004; 52:238-242.
5. Edwards CA, Jeyaraaj R, Indira AJ. (Eds.) Vermitechnology in Human welfare. Rohini Achagam, Coimbatore, Tamil Nadu, India. 2009, 98-108.
6. Iqbal M, Khan AG, Hassan A, Islam KR. Tillage and Nitrogen Fertilization Impact on Irrigated Corn Yields, and Soil Chemical and Physical Properties Under Semi-arid Climate. J of Sustainable Water. Sc. & Mang. 2013; 1(3):90-98.
7. Jackson ML. Soil Chemical Analysis. Prentice Hall of India Private Ltd., New Delhi, 1958.
8. Khatkar Y, Dawson J, Rao ZKK, Dixit PM, Khatkar R. Effect of nitrogen, phosphorus and sulphur fertilization on growth and yield of maize (*zea mays*) Inter. J of Agri. Sc. 2009; 5(2):396-39.
9. Khanday ABS, Sharma U, Dubey, Bhardawaj SB. Effect of different fertilizer and irrigation management system on physico-chemical properties and yield of maize cropping sequence. Indian journal of Agronomy. 2012; 57(3):222-228.
10. Laxminarayan. Effect of integrated use of inorganic and organic manure on soil properties yield and nutrient uptake in ultisols to mizorum. Journal of the Indian society of soil Science. 2006; 54:120-123.
11. Lavelle P, Martin M. Small scale and large scale effects of endogeic earthworms on soil organic matter dynamics in soils of the humid tropics. Soil Bio and Biochem. 1992; 24:81-96.
12. Liu M, FHu X, Chen Q, Huang J, Jiao B, Zhang H Li. Organic amendments with reduced chemical fertilizer promotes soil microbial development and nutrient availability in a subtropical paddy field: the influence of quantity, type and application time of organic amendments. Appl Soil Ecol. 2009; 42:166-175.
13. Lourduraj CA, Yadav BK. Effect of vermicompost application on soil and plant growth. Verms and vermitechnology. A.P.H. Publ. Cor. 5, Ansari Road, Darya Ganj, NewDelhi. 2005, 81-96.
14. Melero M, Vanderlinden K, Ruiz JC, Madejon E. Long-term effect on soil biochemical status of a vertisol under conservation tillage system in semi-arid Mediterranean conditions. European journal of soil biology. 2008; 44(4):437-442.
15. Olsen SR, Cole CV, Watanhe FS, Dean LA. Estimation of available phosphorus in soil by interaction with sodium bicarbonate. U.S. Deptt. Agri. Circ. 1954, 939.
16. Prabha LM. Vermitech- A potential technology for the conversion of wastes into biofertilizer Ph.D. Thesis Department of Biochemistry, Kongunadu Arts and Science College, Coimbatore, Tamil Nadu, India. 2006, 79-86.
17. Rai Mangla. Effect of integrated nutrient management local and hybrid varieties of maize (*Zea mays* L.) yield. Hand Book of Agriculture. 2006, 872-886.
18. Ramasamy PK, Baskar K, Ignacimuthu S. Influence of vermicompost on kernel yield of Maize (*Zea mays* L.). Elixir Agriculture. 2011; 36(3):119-3121.
19. Singh RN, Sulatiya R, Ghata KRSA. Effect of higher application of Nitrogen and potassium over recommended level on growth yield and yield attribute of late sown winter maize (*Zea mays* L.) crop research, Hissar. 2003; 26(1):71-74.
20. Singh KP, Snman A, Singh PN, Srivastava TK. Improving quality of sugarcane-growing soils by organic amendments under subtropical climatic conditions of India. Biol Fertil Soils. 2007; 44:367-376.
21. Subbaiah BV, Asija CL. A rapid procedure for the estimation of available nitrogen in soil. Current Sci. 1956; 25:415-426.
22. Toth SJ, Prince AL. Estimation of Cation exchange capacity and exchangeable Ca, K and NA content of soil

by flame photometer technique. Soil. Sci. 1949; 67:439-445.

23. Walkley A, Black CA. Critical examination of rapid method for determination organic carbon in soil, effect on variation in digestion condition and of inorganic soil constituents. Soil. 1947, 632-651.
24. Wilcox LV. electrical conductivity, Amer. Water Works Assoc. J. 1950; 42:775-776.