



P-ISSN: 2349-8528

E-ISSN: 2321-4902

IJCS 2018; 6(2): 2998-3002

© 2018 IJCS

Received: 17-01-2018

Accepted: 18-02-2018

Prativa Sahu

ICAR- Indian Institute of Water
Management, Bhubaneswar,
Odisha, India

Ankita Sahu

ICAR-Central Institute for
Women in Agriculture,
Bhubaneswar, Odisha, India

Effect of pre-harvest sprays of forchlorfenuron and boron on fruit cracking and quality of pomegranate (*Punica granatum* L.) cv. Kandhari

Prativa Sahu and Ankita Sahu

Abstract

An investigation on pre-harvest spray of forchlorfenuron (CPPU) and boron along with some *in-situ* soil moisture conservation techniques was carried out on seven-year old plants of pomegranate cultivar Kandhari trained as four stems grown under rainfed conditions of Himachal Pradesh. Experimental plants were subjected to 11 treatments, viz., mulching, CBOC (crescent bund with open catchment pits), CPPU at 5 or 10 ppm, H₃BO₃ at 0.2 or 0.4%, mulching + CPPU at 5 ppm or H₃BO₃ at 0.2%, CBOC + CPPU at 5 ppm or H₃BO₃ at 0.2% and control. Growth parameters like plant height, plant spread, trunk girth and annual shoot growth were observed significantly higher under the treatment CBOC + H₃BO₃ at 0.2%. However, leaf area and leaf chlorophyll content were recorded significantly higher under the treatment CBOC + CPPU at 5 ppm. The extent of fruit cracking was reduced to the lowest level (2.8%) from 11.67% in control and highest fruit yield (26.8Kg/plant) was recorded when the plants were given foliar application of CPPU at 5 ppm under CBOC. The maximum fruit length (86.2 cm) and fruit breadth (88.3 cm), fruit weight (419.74 g), aril weight (279.21 g), aril percentage (66.52), juice content (67.93 ml/100 g), ascorbic acid content (16.84 mg/100g) were obtained with the treatment of CPPU at 5ppm +CBOC. The lowest titratable acidity (0.49%), highest TSS (15.210 B) and TSS/acid ratio (31.07) was recorded in fruits from plants under the treatment of H₃BO₃ at 0.2% with mulching. Total sugar, reducing sugar were recorded highest under the treatment H₃BO₃ at 0.2% + CBOC and lowest in control.

Keywords: Pre-harvest sprays, forchlorfenuron, boron, fruit cracking, *Punica granatum* L.

Introduction

Pomegranate (*Punica granatum* L.) is one of the most historic fruits and considered as a symbol of fertility and often quoted in the theological texts (Mars, 2000) [1]. Though pomegranate was originally adapted to Mediterranean climate, but it has also flourished well under hot dry summers with cool winters. The foot hills of Himachal Pradesh comprise sub-tropical sub-montane and valley areas of Shivalik hills hold tremendous scope for its cultivation. The plants suffer acute moisture stress leading to fruit cracking due to erratic and uneven rains under rainfed conditions. It therefore becomes imperative to conserve every drop of rain water in the field using *in-situ* moisture conservation techniques. Some practices like mulching of tree basins and soil working techniques like, crescent bund with open catchment pits (Sharma and Singh, 2010) have been found to be useful in many crops for conserving soil moisture.

N-(2-chloro-4-pyridinyl)-N-phenylurea (CPPU, forchlorfenuron) possesses strong cytokinin bioactivity and has been shown to stimulate the fruit growth in a number of fruit crops (Ahmed *et al.*, 2007; Kim *et al.*, 2006; Fang-XueZhi *et al.*, 2006; Supe and Marshal, 2008; Sharma and Belsare, 2009) [2, 4-7]. The most important physiological effects of boron in plants are structural role in cell walls, stimulation or inhibition of specific metabolic pathways, involvement in the movement of Ca into the plant, flower production and retention, pollen tube elongation and germination, and seed and fruit development, sugar transport and hormonal regulation (Alila *et al.*, 2005; Topno *et al.*, 2006; Kumar, 2006; Khalifa *et al.*, 2009) [9-11]. Therefore, the present study was undertaken to ascertain the effect on pre-harvest spray of forchlorfenuron (CPPU) and boron along with some *in-situ* soil moisture conservation techniques like mulching, crescent bund with open catchment pits in pomegranate cv. Kandhari with an objective to evaluate the morphological and physiological status, fruit cracking, yield potentials and quality of pomegranate under rainfed conditions.

Correspondence

Prativa Sahu

ICAR- Indian Institute of Water
Management, Bhubaneswar,
Odisha, India

Materials and Methods

The present investigation was carried out in the experimental orchard of the Department of Fruit Science, Dr Y.S. Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India during 2010-2011. For this study, 33 plants of 7-year-old pomegranate trained as four stem system and spaced 4 m x 4 m apart were selected. Soil at the experimental site was silty loam; having 6.81 pH, 1.76% organic carbon content, 8.00% permanent wilting point, 25.50% field capacity and 1.10 g cm⁻³ bulk density. Experimental plants were subjected to 11 treatments, viz. mulching (10 cm thick hey), CBOC (crescent bund with open catchment pits), CPPU (forchlorfenuron) at 5 or 10 ppm, H₃BO₃ at 0.2 or 0.4%, Mulching + CPPU at 5 ppm or H₃BO₃ at 0.2%, CBOC + CPPU at 5 ppm or H₃BO₃ at 0.2% and control (no treatment). These treatments were applied in mid-March (*in-situ* soil moisture conservation) and mid-May (CPPU & boron). All the treatments were replicated three times in a randomized block design.

For the preparation of CPPU solutions, the required quantity of its commercial formulation 'Sitofex' procured from M/S Degussa AG; Germany was directly dissolved in 12 litres of water. The required amount of H₃BO₃ was weighed on a digital balance and dissolved in warm water. In order to decrease surface tension and to facilitate absorption of droplets, 0.05 percent 'Break Thru' (surfactant) was added to each solution prior to use. Soil moisture (%) were taken with the help of AquaPro® soil moisture profiler at 15 day intervals during the growing season from 1st April to 15th July at 15, 30, 45 cm depths. The access tubes fitted 1 m away from the trunk.

The percentage increase in annual shoot growth was observed in the field before the commencement of the growth and after

the cessation of growth in autumn. Simultaneously, 20 fully expanded leaves were collected at random from the periphery of each plant in the month of August and their area (cm²) was measured with the help of LI-Cor 3100 leaf area meter. The number of the fruits cracked on each experimental plant were counted periodically during the entire fruiting season and expressed in percent. Fruit yield (kg/tree) was determined on the basis of total weight of fruits harvested from the each experimental plant. Russet formation on the skin of five randomly selected fruits from each replication was examined visually at the time of harvest and russet development on the fruits was determined on 10 point scale. The total soluble solids of the juice were determined with Erma-hand refractometer (0 to 320B range). Ascorbic acid content of the fruits was determined as per A.O.A.C. (1980) method. The data were subjected to statistical analysis as per the procedure outlined by Panse and Sukhatme.

Results and Discussion Soil moisture contents under the plant basins fluctuated greatly with dry spells and erratic rainfall cycles (1, 0, 0, 2.7, 48.2, 0, 116.6, 52.2, 280.7, 203.9 mm rainfall in 15th March, 1st April, 15th April, 1st May, 15th May, 1st June, 15th June, 1st July, 15th July, respectively during the course of investigation. However during all the observation dates, maximum soil moisture contents of 18.01% at 0-15 cm soil depth; 18.88% at 15-30 cm soil depth and 19.03% at 30-45 cm soil depth were recorded under the treatment CBOC + CPPU at 5 ppm. Soils under mulching also maintained more moisture level. The interaction effect of treatments with soil moisture levels during different period was also found to be significant.

Table 1: Effect of pre-harvest spray of forchlorfenuron and boron on annual shoot extension growth, average leaf area, Leaf chlorophyll content, fruit set and fruit cracking of pomegranate cv. Kandhari

Treatments	Annual shoot extension growth (cm)	Average leaf area (cm ²)	Leaf chlorophyll content (mg/g of fresh wt.)	Fruit set (%)	Fruit cracking (%)
T ₁ : Mulching with hay (10cm)	26.93	9.2	1.92	20.4	5.2
T ₂ : Crescent bund with open catchment pits (CBOC)	32.7	10.6	1.99	22.7	3.9
T ₃ : CPPU at 5 ppm	25.15	9.4	2.61	19.8	4.1
T ₄ : CPPU at 10 ppm	27.27	8.7	2.57	18.6	4.53
T ₅ : H ₃ BO ₃ at 0.2%	37.43	9.1	2.3	18.97	5.1
T ₆ : H ₃ BO ₃ at 0.4%	31.13	8.4	2.21	17.89	5.6
T ₇ : Mulching+ CPPU at 5 ppm	27.37	9.7	2.78	23.6	3.5
T ₈ : Mulching+ H ₃ BO ₃ at 0.2%	39.57	9.19	2.41	23.2	3.8
T ₉ : CBOC+ CPPU at 5 ppm	34.29	11.05	2.93	24	2.8
T ₁₀ : CBOC+ H ₃ BO ₃ at 0.2%	43.15	10.8	2.45	23.5	3.2
T ₁₁ : Control	17.75	7.07	1.41	13.33	11.67
^{CD} 0.05	1.817	0.55	0.05	0.55	0.28
SE ±	0.87	0.26	0.022	0.26	0.13

Highest shoot growth (43.15 cm) was recorded in plants under CBOC when sprayed with boric acid at 0.2%. The results can be attributed to the availability of more moisture under CBOC and mulching at all soil depths and role of boron in N metabolism, hormone movement action and cell division (Russel, 1957) [13]. The leaf area (11.05 cm²), accumulation of chlorophylls (2.93 mg/100 g fresh weight) was significantly higher under the treatment CBOC + CPPU at 5 ppm (Table 1). The present findings are in agreement with those of Iersel and Nemali, 2004, and Jyothi and Rajadhadh, 2004, who observed that higher soil moisture level favoured more leaf growth in Rangapur lime. The CPPU is an urea based cytokinin which induced the activity of invertase enzyme in conversion of lipids to glucose, fructose (Notodimedjo, 2000)

[16] and as solute concentration increases and OP becomes negative, thus resulting in more uptake of water, leading to better leaf expansion. Cytokinin helps in the retention of chlorophyll and stimulates the nutrient mobilization and thus might have resulted in the accumulation of more chlorophyll in treated leaves. In the study, the plants grown under the treatment of CBOC+ CPPU at 5 ppm (T₉), and CBOC+ H₃BO₃ at 0.2% (T₁₀) had higher fruit set. These results are in accordance with the findings of Farmahan and Sharma (2005), who observed increased fruit set due to *in-situ* moisture conservation in pomegranate. Boric acid helps in fruit set due to its effect on generative processes on pollen production, germination and pollen tube growth (Chaplin and Westwood, 1980). Fruit cracking was reduced to the lowest

level (2.8%) when the plants grown under CBOC and given foliar application of CPPU at 5 ppm (T9) or H₃BO₃ at 0.2% (T10). Fruit cracking is a physiological disorder, generally associated with differential rate of growth of fruit skin/rind and fruit aril. As CBOC maintained higher soil moisture level soil during the fruit development, which consequently might have decreased fruit cracking in this study. Kaulgud (2001) [18] reported that moisture stress was one of the reasons for fruit cracking in pomegranate. The growth regulator CPPU

might have stimulated overall growth of pomegranate fruits and thereby reduced fruit cracking. The findings are in agreement with those of Singh *et al.* (2003) [19], who observed reduced fruit cracking in pomegranate following the foliar application of boron. The highest fruit yield (26.8 kg/plant) was recorded under T9 (CBOC + CPPU at 5 ppm) over control (14.27 kg/plant).

Table 2: Effect of pre-harvest spray of forchlorfenuron and boron on russetting, fruit yield, fruit length, fruit dia and fruit L/D ratio of pomegranate cv. Kandhari

Treatments	Russetting (10 point scale)	Fruit yield (kg/tree)	Fruit length (mm)	Fruit diameter (mm)	Fruit L/D ratio
T ₁ : Mulching with hay (10cm)	2.66	21.2	81.2	83.4	0.97
T ₂ : Crescent bund with open catchment pits (CBOC)	2.33	22.9	82.1	84.2	0.97
T ₃ : CPPU at 5 ppm	1.67	22.1	84.5	86.6	0.98
T ₄ : CPPU at 10 ppm	2	20.6	84	86	0.97
T ₅ : H ₃ BO ₃ at 0.2%	4	21.9	83.7	85.7	0.98
T ₆ : H ₃ BO ₃ at 0.4%	4.33	20.2	83.1	85.3	0.97
T ₇ : Mulching+ CPPU at 5 ppm	2	24.9	85.3	87.4	0.98
T ₈ : Mulching+ H ₃ BO ₃ at 0.2%	3.33	23.3	83.9	85.9	0.98
T ₉ : CBOC+ CPPU at 5 ppm	1.33	26.8	86.2	88.3	0.98
T ₁₀ : CBOC+ H ₃ BO ₃ at 0.2%	3.66	24.02	84.1	86.2	0.97
T ₁₁ : Control	5	14.27	75.16	76.93	0.97
^{CD} 0.05	1.45	0.46	0.36	0.30	NS
SE ±	0.69	0.22	0.17	0.14	NS

Table 3. Effect of pre-harvest spray of forchlorfenuron and boron on fruit weight, aril weight, aril percentage, juice content and TSS of pomegranate cv. Kandhari

Treatments	Fruit weight (g)	Aril weight (g)	Aril percent age	Juice content (ml/100g)	TSS (° B)
T ₁ : Mulching with hay (10cm)	326.98	192.00	58.73	64	13.3
T ₂ : Crescent bund with open catchment pits (CBOC)	344.73	203.9	59.15	64.71	13.71
T ₃ : CPPU at 5 ppm	401.56	260.73	64.93	66.86	14.97
T ₄ : CPPU at 10 ppm	384.39	246.00	64	66.79	14.52
T ₅ : H ₃ BO ₃ at 0.2%	363.06	221.79	61.09	65	13.88
T ₆ : H ₃ BO ₃ at 0.4%	344.91	209.63	60.78	64.56	13.66
T ₇ : Mulching+ CPPU at 5 ppm	414.52	269.52	65.02	67.31	14.09
T ₈ : Mulching+ H ₃ BO ₃ at 0.2%	371.25	232.43	62.61	66.72	15.21
T ₉ : CBOC+ CPPU at 5 ppm	419.74	279.21	66.52	67.93	14.28
T ₁₀ : CBOC+ H ₃ BO ₃ at 0.2%	376.3	237.85	63.21	66.8	15.03
T ₁₁ : Control	287.96	154.98	53.82	56.73	12.8
^{CD} 0.05	8.85	1.13	0.05	0.038	0.02
SE ±	4.24	0.54	0.02	0.018	0.01

In the study, pre-harvest application of CPPU significantly influenced the fruit finish in respect of russet formation on skin (Table 2). It was observed that the least russet formation (1.33 on 10 point scale) occurred in fruits under treatment of CBOC+ CPPU at 5ppm (T₉), which was superior to control (5.00 points). Application of boric acid irrespective of the concentration was not effective in controlling the russetting on fruit skin. Maximum fruit length (86.2 mm), fruit diameter (88.3 mm), fruit weight (419.74 g), aril weight per fruit (279.21g), aril percentage (66.52), juice content (67.93 ml/100 g), ascorbic acid content (16.84

mg/100g) were obtained (Table 2 & 3) with the treatment of CPPU at 5ppm +CBOC, closely followed by the treatment T₇ (Mulching+ CPPU at 5 ppm). The CPPU is an urea based cytokinin (Ranjan *et al.*, 2003) [20], and its application at earlier stage of fruit development might have enhanced cell division of different fruit parts and consequently increased fruit size in the present study. Earlier, the synthetic cytokinin CPPU was reported to promote fruit development in apple (Costa *et al.* 2004) [21], grapes (Stern *et al.* 2002) [22] and kiwi fruit (Kim, 2006) [4].

Table 4: Effect of pre-harvest spray of forchlorfenuron and boron on acidity, TSS/acid ratio, ascorbic acid, total sugar and reducing sugar of pomegranate cv. Kandhari

Treatments	Acidity	TSS/Acid ratio	Ascorbic acid (mg/100g)	Total Sugar (%)	Reducing Sugar (%)
T ₁ :Mulching with hay (10cm)	0.68	19.58	14.64	11.38	9.92
T ₂ :Crescent bund with open catchment pits(CBOC)	0.63	21.79	15.32	11.74	10.33
T ₃ :CPPU at 5 ppm	0.53	28.25	16.06	11.66	10.19
T ₄ :CPPU at 10 ppm	0.55	26.49	15.21	11.69	10.45
T ₅ :H ₃ BO ₃ at 0.2%	0.60	23.17	16.34	11.71	10.46
T ₆ :H ₃ BO ₃ at 0.4%	0.65	21.05	15.91	11.61	10.23
T ₇ :Mulching+ CPPU at 5 ppm	0.59	24.31	16.59	11.67	10.52
T ₈ : Mulching+ H ₃ BO ₃ at 0.2%	0.49	31.07	16.65	11.83	10.63
T ₉ : CBOC+ CPPU at 5 ppm	0.56	25.55	16.84	11.76	10.55
T ₁₀ : CBOC+ H ₃ BO ₃ at 0.2%	0.51	29.35	16.70	11.95	10.67
T ₁₁ :Control	0.73	17.54	12.83	10.03	9.25
^{CD} 0.05	0.04	1.74	0.22	0.22	0.19

The lowest titratable acidity (0.49%), highest TSS (15.210 B) and TSS/acid ratio (31.07) was recorded in fruits from plants under the treatment of H₃BO₃ at 0.2% with mulching (Table 4). Total sugar, reducing sugar were recorded highest under the treatment H₃BO₃ at 0.2% + CBOC and lowest in control. These results are in conformity with those of Kano (2003) who reported increase in sugar content with the foliar application of CPPU at 20 ppm on Japanese pear trees. CPPU applied at 5 ppm increased the sucrose, glucose, fructose contents in kiwi Fang-XueZhi *et al.* (2006) [5].

From the ongoing study it can be concluded that growth parameters like plant height, plant spread, trunk girth and annual shoot growth were observed significantly higher under the treatment CBOC + H₃BO₃ at 0.2%. However, leaf area and leaf chlorophyll content were recorded significantly higher under the treatment CBOC + CPPU at 5 ppm. Physiological characteristics such as photosynthetic rate and transpiration rate were higher in the plant under crescent bund with open catchment pits. The extent of fruit cracking was reduced to the lowest level (2.8%) from 11.67% in control and highest fruit yield (26.8Kg/plant) was recorded when the plants were given foliar application of CPPU at 5 ppm under CBOC. The least russet formation (1.33 on 10 point scale) occurred in fruits from plants given the treatment of CPPU at 5ppm +CBOC. The maximum fruit length (86.2 cm) and fruit breadth (88.3 cm), fruit weight (419.74 g), aril weight (279.21 g), aril percentage (66.52), juice content (67.93 ml/100 g), ascorbic acid content (16.84 mg/100g) were obtained with the treatment of CPPU at 5ppm +CBOC. The lowest titratable acidity (0.49%), highest TSS (15.210 B) and TSS/acid ratio (31.07) was recorded in fruits from plants under the treatment of H₃BO₃ at 0.2% with mulching. Total sugar, reducing sugar were recorded highest under the treatment H₃BO₃ at 0.2% + CBOC and lowest in control. Hence, pre-harvest spray of forchlorfenuron and boron along with soil working technique-crescent bund with open catchment pit can be employed to conserve more soil moisture, reducing fruit cracking and increase production of pomegranate with improved fruit quality in rain fed conditions.

References

- Mars M. Pomegranate plant material: Genetic resources breeding, a review: Options Mediterranean series A, Seminaries Mediterranean, 2000; (42):55-62
- Ahmed FF, Aal, Aka MA. Effect of concentrations and date of spraying Sitofex on yield and quality of Le- Conte pear fruits. 8th –African-Crop-Science –Society-Conference, Egypt, 2007, 523-527
- Sharma N, Singh K. Effect of *in situ* moisture conservation on morphology, physiology and production of olives under rainfed conditions. Indian Journal of Horticulture 2010; 64(4):364-366
- Kim JG, Takami Y, Mizugami T, Beppu K, Fukuda T, Kata I. CPPU application on size and quality of hardy kiwifruit. Scientia Horticulturae 2006; 110(2): 219-222.
- Fang-XueZhi, Fei-XueQian, Ding-Ming, Yiao-XiaoHua, Xie-YiFu, Zhau-LiXiong. Effect of different concentrations of CPPU on growth and nutritional quality of *Actinidia deliciosa*. Acta Agricultural Universitatis Jiangxiensis. 2006; 28(2):217-221
- Supre VS, Marshal SK. Effect of plant growth substances on fruit quality of pomegranate. Asian Journal of Horticulture. 2008; 3(1):133-135
- Sharma N, Belsare C. Effect of plant bio-regulators and nutrients on fruit cracking and quality in pomegranate (*Punica granatum* L.) 'G-137' in Himachal Pradesh. Acta Horticulture. 2009;(890):347-352
- Alila P, Sanya D, Akali S.. References of papaya cv. Ranchi to micronutrient application. Horticultural Journal. 2005; 18(2):121-125
- Topno ATP, Ray RN, Bauri MK, Sengupta S. Effect of nutrients on fruit cracking and physico-chemical composition of litchi (*Litchi chinensis* Sonn.) cv. Purbi. Journal of Research Birsa Agricultural University 2006; 18(1):141-144
- Kumar A. Studies on the effect of soil and foliar application of zinc and boron on fruit set, retention, quality, cracking and yield of litchi cv. Dehradun. Ph.D. Thesis, Dr YS Parmar University of Horticulture and Forestry, Nauni, Solan HP. India 2006.
- Khalifa RKM, Omaira MH, Abd-El-Khair H.. Influence of foliar spraying with boron and calcium on productivity, fruit quality, nutritional status and controlling of blossom end rot disease of Anna apple trees. World Journal of Agricultural Sciences. 2009; 5(2):237-249
- Ranganna S. Handbook of analysis and quality control for fruits and vegetable production McGraw Hill Publishing Company Limited New Delhi 1995, 1-21
- Russel DA. Boron and soil fertility. In: The yearbook of Agriculture USDA, Washington, DC, 1957.
- Iersel MW, Nemali KS. Effect of soil moisture on growth parameters of longan. Hort Science. 2004; 39(6):1298-1301
- Jyothi Hadli, Rajadhav SB. Effect of soil moisture stress on growth and physiological attributes of different strains

- of Rangpur lime. Journal of Maharashtra Agricultural University. 2004; 29(3):263-266
16. Notodimedjo S. Effect of GA₃, NAA and CPPU on fruit retention, yield and quality of mango (cv.Arumanis) in East Java. Acta Horticulturae 2000; 509:587-600
 17. Chaplin MH, Westwood MN. Relationship of nutritional factors to fruit-set. Journal of Plant Nutrition 1980; 2:477-505
 18. Kaulgud SN. Pomegranate. In: Chadha, K.L. (ed.). Handbook of Horticulture. 2001, 297-304
 19. Singh DB, Sharma BD, Bhargava R. Effect of boron and GA₃ to control fruit cracking in pomegranate (*Punica granatum*). Current Agriculture 2003; 27(1/2): 125-127
 20. Ranjan A, Purohit SS, Prasad V. Plant Hormones: Action and Application, 2003, 245.
 21. Costa G, Bucchi F, Montefiori M, Bregoli AM, Grappadelli LC. Thinning activity and fruit quality of Gala and Fuji apple varieties as affected by cytokinins. Acta Horticulturae. 2004; (653):107-113
 22. Stern RA, Flaishman MA, Shargal A. Effect of the synthetic cytokinin CPPU on fruit size and yield of 'Spadona' pear. Acta Horticulturae. 2002; 596:797-80
 23. Kano Y. Effects of GA and CPPU treatments on cell size and types of sugars accumulated in Japanese pear fruit. Journal of Horticultural Science and Biotechnology. 2003; 78(3):331-334.