



P-ISSN: 2349-8528

E-ISSN: 2321-4902

IJCS 2018; 6(4): 162-165

© 2018 IJCS

Received: 17-05-2018

Accepted: 19-06-2018

**HS Patel**

Department of Silviculture and Agroforestry ASPEE College of Horticulture and Forestry Navsari Agricultural University Navsari, Gujarat, India

**MB Tandel**

Assistant Professor (Forestry), Department of Silviculture and Agroforestry ASPEE College of Horticulture and Forestry Navsari Agricultural University Navsari, Gujarat, India

**VM Prajapati**

Department of Silviculture and Agroforestry ASPEE College of Horticulture and Forestry Navsari Agricultural University Navsari, Gujarat, India

**MH Amlani**

Department of Silviculture and Agroforestry ASPEE College of Horticulture and Forestry Navsari Agricultural University Navsari, Gujarat, India

**DH Prajapati**

Department of Silviculture and Agroforestry ASPEE College of Horticulture and Forestry Navsari Agricultural University Navsari, Gujarat, India

**Correspondence**

**MB Tandel**

Assistant Professor (Forestry), Department of Silviculture and Agroforestry ASPEE College of Horticulture and Forestry Navsari Agricultural University Navsari, Gujarat, India

## International Journal of *Chemical Studies*

### Effect of different pre-sowing treatments on germination of Red sanders (*Pterocarpus santalinus* L. f.) in Poly house condition

**HS Patel, MB Tandel, VM Prajapati, MH Amlani and DH Prajapati**

#### Abstract

Red sanders (*Pterocarpus santalinus* L. f.) is an endangered and endemic tree species that occurs in Southern parts of Eastern Ghats of India. Heartwood of Red Sanders has high demand. The wood demand, restricted distribution, slow regeneration, illegal harvest, trade and habitat destruction and various limitations in propagation of Red Sanders by natural means are prolonged dormancy, low germinability and poor viability of seeds the present investigation was carried out to find out the best pre-sowing treatments in Red sanders (*Pterocarpus santalinus* L. f.). This experiment was comprised of eleven pre-sowing treatments viz., T<sub>1</sub> (Tap water for 4 days), T<sub>2</sub> (Tap water for 8 days), T<sub>3</sub> (Lukewarm water for 4 days), T<sub>4</sub> (Lukewarm water for 8 days), T<sub>5</sub> (GA<sub>3</sub> @ 250ppm for 1 day), T<sub>6</sub> (GA<sub>3</sub> @ 500ppm for 1 day), T<sub>7</sub> (10 % sulphuric acid for 1day), T<sub>8</sub> (10 % hydrochloric acid for 1day), T<sub>9</sub> (10 % nitric acid for 1day), T<sub>10</sub> (Separating seeds from pods with a sharp knife or scalpel and sown directly), T<sub>11</sub> (Control) and grown in Poly house condition with Completely Randomized Design (CRD) with three repetitions. Among different pre-sowing treatments, seeds pre-treated with T<sub>6</sub> (GA<sub>3</sub> @500ppm for 1 day) registered earlier sprouting (7.67) and higher germination percentage (70.00), collar diameter (5.34 mm), plant height (40.03 cm), number of leaves per plant (19.44), survival percentage (63.33 %), root length (30.43 cm), fresh (30.45 g/plant) and dry (23.48 g/plant) weight of plant. The next best pre-sowing treatment are T<sub>5</sub> (GA<sub>3</sub> @ 250ppm for 1 day) and T<sub>10</sub> (Separating seeds from pods with a sharp knife or scalpel and sown directly). The seeds which were no treated (T<sub>11</sub> : control) found poorest for all parameters under study.

**Keywords:** *Pterocarpus santalinus* L. f., Red Sanders, GA<sub>3</sub>, Poly house and germination traits

#### Introduction

Red Sanders (*Pterocarpus santalinus* L. f.) is an endangered and endemic tree species that occurs in Southern parts of Eastern Ghats of India. Being a commercially important tree of Indian forests commonly termed in the trade as “Red Sanders” and locally considered as the “Pride of India” and the “State Tree of Andhra Pradesh”. Due to prevailing dry and hot conditions in its natural habitat, the pollination ecology is vulnerable. As an endemic species, for its continuous survival, Red Sanders has devised cross pollination as the main system exhibiting xenogamous breeding though geitonogamy exists a bit (Rao and Raju, 2002) [14]. The matured reddish brown coloured winged pods are formed after eleven months of flowering and have one or two seeds (Dayanand, 1988) [3, 4]. Heartwood of Red Sanders has high demand in domestic as well as international market and the wavy grained wood is valued. It was found that seedlings raised from wavy grained trees were slow growing compared to that of straight grained trees. The seedlings had short internodes, smaller and greener leaves, petioles and more compact crown and concluded that these traits were distinct and can be used for segregating the seedlings. In view of its high timber price, overexploitation, restricted distribution and poor seed germination there is an urgent need for conservation of this plant species failing which the category of endangered may move to extinct.

Though, macro and micro propagation protocol have been developed, further refinement is required for mass propagation. The poor seed germination was reported by Kalimuthu and Lakshmanan (1995) [7]. Seed germination has been advocated as one of the most viable tools for the ex situ conservation of threatened germplasm (Rao, 2004) [13].

Considering the wood demand, restricted distribution, slow regeneration, illegal harvest, trade and habitat destruction, and various limitations in propagation of red sanders by natural means are prolonged dormancy, low germinability and poor viability of seeds.

## Materials and Methods

The pods of Red Sanders (*Pterocarpus santalinus* L. f.) were brought from Bengaluru, Karnataka and pods were sown directly into the polythene bag in the poly house. In one polythene bag one pod is sown at the depth of 2 cm. One pod of Red Sanders contains two seeds. The pods were treated with the following treatments *viz.*, T<sub>1</sub>: Pods of Red Sanders were kept in Tape water and allowed soaking for 4 days. Water was regularly changed once in 24 hours time period for the leach out of the phenolic compounds present in the pods; T<sub>2</sub>: Pods of Red Sanders were kept in Tape water and allowed soaking for 8 days. Water was regularly changed once in 24 hours time period for the leach out of the phenolic compounds present in the pods; T<sub>3</sub>: Pods of Red Sanders were kept in lukewarm water and allowed soaking for 4 days. Water was regularly changed once in 24 hours for leach out of the chemicals present in the pods; T<sub>4</sub>: Pods of Red Sanders were kept in lukewarm water and allowed soaking for 8 days. Water was regularly changed once in 24 hours for leach out of the chemicals present in the pods; T<sub>5</sub>: GA<sub>3</sub> @ 250 ppm for 24 hours; T<sub>6</sub>: GA<sub>3</sub> @ 500 ppm of solution for 24 hours; T<sub>7</sub>: 10 % H<sub>2</sub>SO<sub>4</sub> solution for 20 minutes and then after placed in water for 24 hours ; T<sub>8</sub>: 10 % HCl solution for 20 minutes and then after placed in water for 24 hours; T<sub>9</sub>: 10 % H<sub>2</sub>SO<sub>4</sub> solution for 20 minutes and then after placed in water for 24 hours ; T<sub>10</sub>: Seeds were separated from the pods with the help a sharp knife or scalpel and then sown directly into the poly bags. In each poly bag one seed were sown; T<sub>11</sub>: No pre-treatments were given to the pods of Red Sanders and directly sown in the poly bags in the Net house conditions. The observations on days to first sprout, germination percentage (%), plant height (cm), number of leaves per plants, collar diameter (mm), root length (cm), survival percentage (%), fresh weight of plant (g) and dry weight of plant (g) were recorded. Statistical analysis of the data of various characters studied in present investigation was carried out through the procedure of Completely Randomized Design (CRD). The appropriate standard error of mean (S. Em. ±) and the critical difference (C. D.) were calculated at 5 percent level of probability.

## Result and Discussion

It is evident from the data presented in Table – 1 that days to first sprout, germination percentage, number of leaves per plant, collar diameter, plant height, root length, fresh weight of plant, dry weight of plant and survival percentage were significantly influenced by different pre-sowing treatments in the present study.

### 1. Germination Parameters

The results of the study revealed that the minimum number of days to first sprout (7.67 days) and maximum germination percentage (70.00 %) were recorded in pods treated with GA<sub>3</sub> at 500 ppm for 1 day (T<sub>6</sub>) which was at par with Separating seeds from pods with a sharp knife or scalpel and sown directly (T<sub>10</sub> : 8.33 days and 66.67 %, respectively) and GA<sub>3</sub> at 250 ppm for 1 day (T<sub>5</sub> : 8.33 days and 63.33 %, respectively). The minimum germination percentage was recorded in treatment control (T<sub>11</sub> : 46.67 %). It might be due to the involvement of GA<sub>3</sub> in the activation of cytological

enzymes along with increase in cell wall plasticity and better water absorption. GA<sub>3</sub> acts directly on embryo relieving them from dormancy through promoting protein synthesis and elongation of coleoptiles and leaves and also helps in the production of ethylene. This ethylene invokes the synthesis of hydrolases, especially amylase, which favours the seed germination (Stewart and Freebairn, 1969) [21]. Another reason is that GA<sub>3</sub> stimulates seed germination by formation of α-amylase enzymes which converts insoluble starch into soluble sugars and it also initiates the radical growth by removing some metabolic blocks as suggested by Gillard and Walton (1973) [5]. The remarkable effect of GA<sub>3</sub> on seed germination especially in breaking seed dormancy has been well established by several workers; Dayanand and Lohidas (1988) [3, 4], Naidu and Mastan (2001) [11] and Naidu (2001) [11] in Red sanders; Shivanna *et al.* (2007) [20] in *Terminalia bellerica*, Banyal (2010) [1] in *Pinus gerardiana*, Das and Tah (2013) [2] in *Santalum album*, Kala *et al.* (2014) [6] in *Canarium strictum*, Lilabati and Sahoo (2015) [9] in *Embllica officinalis* and Sutteesh *et al.* (2016) [22] in *Santalum album* and Rout *et al.* (2016a) [15] in *Delonix regia*.

The results of the present study also revealed that the separating seeds from pods with a sharp knife or scalpel and sown directly increase the permeability of air and water through seed which favours the second most good seed germination and higher germination percentage which was statistically at par with GA<sub>3</sub> at 500 ppm for 1 day. Dayanand and Lohidas (1988) [3, 4] also recorded highest seed germination up to 83.50 per cent in the treatment of separating seeds with a sharp knife or scalpel before sowing the Red Sanders seed. More or less similar results were observed by Shivanna *et al.* (2007) [20] in *Terminalia bellerica* and Khan (2013) [8] in *Pongamia pinnata* L. and *Pterocarpus marsupium* Roxb.

### 2. Growth Parameters

Among the different pre-sowing treatments, GA<sub>3</sub> 500 ppm for 1 day (T<sub>6</sub>) gave maximum plant height (40.03 cm), number of leaves per plant (19.44), collar diameter (5.34 mm) and root length (30.43 cm). The increase in plant height with GA<sub>3</sub> treatment was due to the fact that this hormone increased osmotic uptake of nutrients, causing cell multiplication and cell elongation in the cambium tissue of the intermodal region and thus increased height of the plant (Shanmugavelu, 1966) [19] and collar diameter also increased due to gibberellic acid promoted cell division and cell elongation in the collar region (Sen *et al.*, 1990) [18]. The production of more number of leaves might be due to higher growth of seedlings and also due to activity of GA<sub>3</sub> at the apical meristem resulting in more synthesis of nucleoprotein responsible for increasing leaf initiation (Sen and Ghunti, 1976) [17]. The results are also in accordance with result of Masilamani and Dharmalingan (2002) [10] in Silver Oak (*Grevillea robusta* A. Cunn.), Radhakrishnan and Renganayaki (2008) [12] in Simaruba (*Simaruba glauca* Linn) and Rout *et al.* (2016b) [16] in *Delonix regia*.

### 3. Biomass parameters

For assessing the influence of different treatments on biomass the fresh and dry weight of plant were determined. The results of the present investigation significantly revealed that the maximum fresh (30.45 g/plant) and dry (23.48 g/plant) weight of plant were recorded in GA<sub>3</sub> at 500 ppm for 1 day. It might be due to overall growth of the seedling and increased rate of photosynthesis that lead to the overall assimilation and

redistribution of photosynthates within the plant and hence, resulted in higher fresh and dry weight of plant. Thus, increased growth is a consequence of increased dry matter accumulation. The results are in close conformity with the findings of Masilamani and Dharmalingan (2002)<sup>[10]</sup> in Silver Oak (*Grevillea robusta* A. Cunn.), Radhakrishnan and Renganayaki (2008)<sup>[12]</sup> in Simaruba (*Simaruba glauca* Linn) and Rout *et al.* (2016b)<sup>[16]</sup> in *Delonix regia*.

#### 4. Survival percentage

The results revealed the significant differences among the treatments on survival percentage of Red Sanders seedlings.

Among different growth regulators survival percentage (63.33 %) at the end of the experiment after sowing was significantly found to be the maximum when the seeds were treated with GA<sub>3</sub> at 500 ppm for 1 day. This might be due to the overall performance in relation to growth parameters which ultimately increased the survival percentage. The observation analogues to these findings were reported by Masilamani and Dharmalingan (2002)<sup>[10]</sup> in Silver Oak (*Grevillea robusta* A. Cunn.), Radhakrishnan and Renganayaki (2008)<sup>[12]</sup> in Simaruba (*Simaruba glauca* Linn) and Rout *et al.* (2016b)<sup>[16]</sup> in *Delonix regia*.

**Table 1:** Effect of various pre-treatments on germination and growth parameters of Red Sanders (*Pterocarpus santalinus* L. f.)

Treatments	Days to first sprout	Germination percentage	Plant height (cm)	Number of leaves per plant	Collar diameter (mm)	Root length (cm)	Fresh weight of plant (g/plant)	Dry weight of plant (g/plant)	Survival percentage (%)
T <sub>1</sub>	11.00	60.00	33.60	15.78	5.27	26.93	15.92	9.23	50.00
T <sub>2</sub>	10.33	60.00	34.27	16.00	5.30	27.27	18.78	12.05	53.33
T <sub>3</sub>	10.33	60.00	35.20	17.11	5.30	28.00	19.46	12.65	53.33
T <sub>4</sub>	9.67	63.33	36.33	17.22	5.31	28.10	24.07	17.33	56.67
T <sub>5</sub>	8.33	63.33	39.03	18.11	5.34	29.23	29.13	22.17	60.00
T <sub>6</sub>	7.67	70.00	40.03	19.44	5.34	30.43	30.45	23.48	63.33
T <sub>7</sub>	11.67	56.67	31.83	15.67	5.02	26.93	14.35	7.65	50.00
T <sub>8</sub>	13.00	56.67	32.30	16.11	5.05	27.03	16.07	9.34	50.00
T <sub>9</sub>	12.67	53.33	31.07	14.11	5.03	25.03	13.59	6.94	46.67
T <sub>10</sub>	8.33	66.67	37.33	18.11	5.35	29.17	27.19	20.25	56.67
T <sub>11</sub>	17.33	46.67	27.27	8.89	4.86	17.40	11.02	4.65	36.67
S. Em.	0.389	3.624	1.261	0.865	0.124	1.005	1.245	1.168	4.020
C.D. (5%)	1.14	10.63	3.70	2.54	0.36	2.95	3.65	3.43	11.79
C.V.%	6.16	10.51	6.35	9.34	4.13	6.48	10.78	15.27	13.28

#### Conclusion

From this investigation, it is concluded that among different pre-sowing treatments of Red Sanders (*Pterocarpus santalinus* L. f.), T<sub>6</sub> (GA<sub>3</sub> @ 500 ppm for 1 day) performed best with respect to days to first sprout, germination percentage, collar diameter, plant height, number of leaves per plant, survival percentage, root length, fresh and dry weight of plant. The next best pre-sowing treatment were T<sub>5</sub> (GA<sub>3</sub> @ 250 ppm for 1 day) and T<sub>10</sub> (Separating seeds from pods with a sharp knife or scalpel and sown directly). The seeds which were no treated (T<sub>11</sub>: Control) found poorest for all parameters under study. The further study is required to know the effect of GA<sub>3</sub> on plants at field conditions and to know the growth performance of those plants.

#### Reference

- Banyal R. Pre-sowing treatments and germination behaviour of *Pinus gerardiana* Wall- An endangered bio-resource of dry temperate region. J Tree Sci. 2010; 29(1, 2):21-26.
- Das SC, Tah J. Effect of GA<sub>3</sub> on seed germination of Sandal (*Santalum album* L.). Int. J. Curr. Sci. 2013; 8:79-84.
- Dayanand T. Study on development and maturity of pod in Red Sanders (*Pterocarpus santalinus*). Indian Journal of Forestry. 1988; 11(3):207-208.
- Dayanand T, Lohidas T. Effect of different treatments on pod germination of Red sanders (*Pterocarpus santalinus* L. f.). Indian J Forestry. 1988; 11(1):87-88.
- Gillard DF, Walton DC. Germination of *Phaseolus vulgaris* IV. Patterns of Protein Synthesis in Excised Axes. Plant Physiology. 1973; 51:1147-1149.
- Kala Antonyamy, Raja Prakasam, Soosairaj Sebastin, Balaguru Balakrishnan. Reproductive biology and seed

germination of tropical evergreen tree *Canarium strictum* Roxb. Journal of Biodiversity and Environmental Sciences. 2014; 5(6):24-32.

- Kalimuthu K, Lakshmanan KK. Effect of different treatments on pod germination of *Pterocarpus* species. Indian Forester. 1995; 18(2):104-106.
- Khan MR. Studies on seed germination of some useful plants with medicinal and other potentials. Golden Research Thoughts, ISSN 2231-5063, 2013; 3(5):1-3.
- Lilabati L, Sahoo UK. Effect of pre-treatments on seed germination and seedling vigour of *Embliba officinalis* Gaertn. Global Journal of Advance Research. 2015; 2(10):1520-1526.
- Masilamani P, Dharmalingan C. Enhancement of germination and seedling growth in Silver Oak (*Grevillea robusta* A. Cunn.) seeds. Indian J Forestry. 2002; 25(1):30-33.
- Naidu CV, Mastan M. Seed pre-treatment methods to improve germination in *Pterocarpus santalinus* L.f. Indian J. Forestry. 2001; 24(3):342-343.
- Radhakrishnan P, Renganayaki PR. Effect of plant growth regulators on seed germination and seedling growth of stored Simaruba (*Simaruba glauca* Linn) seeds. Indian Forester. 2008; 134(7):947-950.
- Rao KN. Plant genetic resources: Advancing conservation and use through biotechnology. African Journal of Biotechnology. 2004; 3:136-145.
- Rao SP, Raju AJS. Pollination ecology of the Red sanders *Pterocarpus santalinus* (Fabaceae). An endemic and endangered tree species. Current Science. 2002; 8(3):1144-1148.
- Rout S, Sashikala B, Neelam Khare. Effect of GA<sub>3</sub> on seed germination of *Delonix regia*. Research Journal of Recent Sciences. 2016a; 5:1-3.

16. Rout S, Sashikala B, Neelam Khare. Impact of GA<sub>3</sub> seed pre-treatment on seedling growth in *Delonix regia*. Research Journal of Recent Science. 2016b; 5(11):47-49.
17. Sen SK, Ghunti P. Effect of pre-sowing seed treatment on the germination and seedling growth in papaya. Orissa J. Hort. 1976; 4:38-43.
18. Sen SK, Hore IK, Bandhopadhyay A. Pre-sowing seed treatment and its role in germination, seedling growth and longevity of papaya. Orissa J Agril. Res. 1990; 2(3, 4):160-164.
19. Shanmugavelu KG. Studies on the effect of plant growth regulator on the seedling of some tree plant species. South Indian Hort. 1966; 14:24-25.
20. Shivanna H, Balachandra HC, Suresh NL. Effect of pre-sowing treatment on germination of *Terminalia bellerica* (Gaertn.). Karnataka J Agric. Sci. 2007; 20(2):442-443.
21. Stewart ER, Freebairn HT. Ethylene, seed germination and epinasty. Plant Physiology. 1969; 44:955-958.
22. Suthesh VK, Jijeesh CM, Divya TP. Evaluation of organic and inorganic pre-treatments for better seed germination and seedling vigour in *Santalum album* L. Plant Archives. 2016; 16(1):143-150.