



P-ISSN: 2349-8528

E-ISSN: 2321-4902

IJCS 2018; 6(4): 749-751

© 2018 IJCS

Received: 19-05-2018

Accepted: 20-06-2018

Harikesh Jat

Research Scholar Department of
Agronomy, Rajasthan College of
Agriculture, MPUAT, Udaipur,
Rajasthan, India

MK Kaushik

Professor Department of
Agronomy, Rajasthan College of
Agriculture, MPUAT, Udaipur,
Rajasthan, India

JL Choudhary

Professor and Head
Department of Animal
Production, Rajasthan College of
Agriculture, MPUAT, Udaipur,
Rajasthan, India

Hemraj Jat

Research Scholar
Department of Agricultural
Chemistry and Soil Science,
Rajasthan College of Agriculture,
MPUAT, Udaipur, India

NS Solanki

Professor Department of
Agronomy, Rajasthan College of
Agriculture, MPUAT, Udaipur,
Rajasthan, India

LN Dashora

Professor Department of
Agronomy, Rajasthan College of
Agriculture, MPUAT, Udaipur,
Rajasthan, India

Irrigation and Nitrogen Management in Fodder Oat (*Avena sativa* L.) in the agro-climatic zone IV 'a' of Rajasthan

Harikesh Jat, MK Kaushik, JL Choudhary, Hemraj Jat, NS Solanki and LN Dashora

Abstract

A field experiment entitled "Irrigation and Nitrogen Management in Fodder Oat (*Avena sativa* L.)" was conducted during *rabi* 2013-14 at Instructional Farm, Rajasthan College of Agriculture, Udaipur. The experiment comprised combinations of four irrigations and three nitrogen levels. Thus, 12 treatments; all were evaluated in split-plot design with three replications keeping irrigation in main and nitrogen in sub plots. The results indicated that application of irrigations produced significantly higher crop growth rate and dry fodder yield as well as available soil nitrogen after harvest of fodder oat crop. Similarly, addition of 110 kg N ha⁻¹ significantly enhanced the crop growth rate and dry fodder yield as well as available soil nitrogen after harvest of oat crop. Application of irrigation and nitrogen did not significantly affect relative growth rate of fodder oat (*Avena sativa* L.) during year of experiment.

Keywords: Irrigation, Nitrogen, CGR, RGR, Yield and Oat

Introduction

Oat (*Avena sativa* L.) is most important winter cereal forage crop of the north-western and central zone of the country due to congenial climate grown during winter season. Its popularity among the farmers as a fodder crop in the agro-climatic zone IV 'a' of Rajasthan (India) is increasing because of excellent growth habit, quick regrowth/regeneration and high nutritive value. The agronomic requirements such as irrigation and nitrogen fertilization may vary within the region. The identification of the irrigation and nitrogen levels for fodder production and nutritional requirement especially of irrigation and nitrogen are important for getting higher nutritive fodder yield under varying environments.

Materials and Methods

The field experiment was conducted at Instructional Farm, Rajasthan College of Agriculture, Udaipur during *rabi* 2013-14 to assess appropriate irrigation and nitrogen level for increasing yield of fodder oat and to assess available nutrient (nitrogen) status of soil. The experiment consisted of 12 treatment combinations, comprising four irrigation levels (Two-20 & 60 DAS, Three-20, 40 & 60 DAS, Four-20, 40, 60 & 80 DAS and Five-20, 40, 60, 80 & 100 DAS) and three nitrogen levels (70, 90 and 110 kg N ha⁻¹) tested in split-plot design with three replications. The experimental soil was clay loam in texture, slightly alkaline in reaction pH, medium in available nitrogen (295.3 kg ha⁻¹), low in available phosphorus (16.60 kg ha⁻¹) and high in potassium (275.70 kg ha⁻¹). The crop was sown on 26 Oct. 2013 with recommended seed rate of 100 kg ha⁻¹. Total rainfall (18.4 mm) was recorded during crop season. The data pertaining to growth parameters, CGR and RGR between 30-60 and between 60-90 DAS and dry fodder yield of the crop at both first and second cuttings and available soil nitrogen after harvest of the crop were evaluated.

Results and Discussion**Irrigation levels**

It is evident from the data given in Table 1 that the four irrigations gave significantly higher crop growth rate (74.1 and 34.2 g m⁻² day⁻¹) between 30-60 and 60-90 DAS over two irrigations but three and four irrigations were found statistically at par with it while, between 60-90 DAS only five irrigations were statistically at par.

Correspondence**Harikesh Jat**

Research Scholar Department of
Agronomy, Rajasthan College of
Agriculture, MPUAT, Udaipur,
Rajasthan, India

Different irrigations did not significantly affect on relative growth rate during experimental year. Highest dry fodder yield was recorded with the application of three irrigations (8426 kg ha⁻¹) followed by four irrigations (8416 kg ha⁻¹) at first cutting whereas, at second cutting five irrigation produced significantly higher dry fodder yield (11156 kg ha⁻¹) over rest of the treatments. It might be due to dry fodder yield being the function of several yield components depends on complementary interaction between vegetative and reproductive growth of crop; was optimized in fodder oat by irrigation, because irrigation maintained most optimum soil

moisture for growth: yield attributes and yield performance. Two irrigations recorded maximum available nitrogen content in soil (276.94 kg ha⁻¹) which was significantly higher over five irrigations but statistically at par with three and four irrigations. Available N content of soil was decrease due to increased irrigations (Table 1). This is due to nitrogen being an essential plant nutrient for luxuriant and lush growth; plant takes it in abundant quantity. The results of present investigation are in accordance with Kumar *et al.* (2012) [2] and Kumar *et al.* (2013) [3].

Table 1: Effect of irrigation and nitrogen on CGR, RGR, dry fodder yield and available soil nitrogen

Treatments	CGR (g m ⁻² day ⁻¹)		RGR (g g ⁻¹ day ⁻¹)		Dry fodder yield (kg ha ⁻¹)		Soil available nutrient Nitrogen (kg ha ⁻¹)
	Between 30 – 60 DAS	Between 60 – 90 DAS	Between 30 – 60 DAS	Between 60 – 90 DAS	First cutting (60 DAS)	Second cutting (120 DAS)	
Irrigation levels							
Two	48.9	27.5	0.0348	0.1870	7213	7098	276.94
Three	69.4	29.9	0.0427	0.1898	8426	8252	274.85
Four	74.1	34.2	0.0443	0.1940	8416	9706	272.96
Five	67.6	32.1	67.6	0.0422	8405	11156	230.99
S.Em.±	4.1	1.2	4.1	0.0024	272	254	6.20
C.D. (P=0.05)	14.0	4.1	NS	NS	941	881	21.44
Nitrogen levels (kg ha⁻¹)							
70	58.6	27.7	0.0406	0.1872	7405	8328	244.27
90	66.3	31.1	0.0412	0.1911	8229	9206	267.24
110	70.1	34.1	0.0412	0.1941	8709	9624	280.29
S.Em.±	2.0	1.0	0.0008	0.0026	207	211	4.23
C.D. (P=0.05)	6.0	2.9	NS	NS	621	635	12.69

Nitrogen Levels

Application of 110 kg N ha⁻¹ was recorded significantly superior crop growth rate over 70 and 90 kg N ha⁻¹. The per cent increase in crop growth rate by 110 kg N ha⁻¹ was 16.6 and 5.7, 23.1 and 9.6 over 70 and 90 kg N ha⁻¹, between 30-60 and 60-90 DAS, respectively. Relative growth rate of fodder oat did not differ significantly due to nitrogen application. It was observed that the dry fodder yield was increased significantly with addition of nitrogen levels and the maximum was recorded with the application of 110 kg N ha⁻¹ (8709 and 9624 kg ha⁻¹) during both first and second cuttings (Table 1). The higher dry fodder yield with increasing levels of nitrogen might be attributing to the cumulative effect of yield attributing characters viz., crop growth rate and relative growth rate which ultimately increased the fodder yield. The beneficial effect of nitrogen on the CGR may be due to increased cell division, cell enlargement and chlorophyll synthesis ultimately positively effect observed on the fodder yield. The maximum available soil nitrogen (280.29 kg ha⁻¹) was recorded with 110 kg N ha⁻¹ which was significantly higher over 70 kg and 90 kg N ha⁻¹ after harvest of crop. On the basis of analysis soil available nitrogen varied from 244.27 to 280.29 kg ha⁻¹. However, the available status of nitrogen in soil after harvest of crop was observed with increasing fertility levels might be due to buildup of nutrients in soil as result of addition of nitrogenous fertilizer into the soil. The results corroborate the findings of Devi *et al.* (2010) [1], Malakar *et al.* (2009) [4], Patel *et al.* (2011) [5] and Sheoran *et al.* (2010) [6].

References

1. Devi U, Singh KP, Meena S, Kumar S, Kumar S. Effect of nitrogen levels, organic manures and *Azotobacter* inoculation on nutrient uptake of multi-cut oats. Forage Research. 2010; 36:9-14.
2. Kumar N, Kumar S, Arya RK. Effect of cultivars and plant geometry on dry fodder production of barley under various moisture regimes on raised bed planting. Forage Research. 2012; 38:174-176.
3. Kumar S, Dhindwal AS, Arya RK. Dry matter and straw yield in wheat as influenced by preceding crops, planting methods and irrigation levels, Forage Research. 2013; 39:88-92.
4. Malakar B, Mondal S, Bandopadhyay P, Kundu CK. Response of forage oat (var. OS-6) to nitrogen and phosphate fertilizers in the new alluvial zone of West Bengal. Journal of Crop and Weed. 2009; 5:36-38.
5. Patel TU, Arvadia MK, Malik PK, Patel DD, Patel PS. Productivity of oat (*Avena sativa* L.) under different cutting management and split application of nitrogen. Indian Journal of Agronomy. 2011; 56:164-167.
6. Sheoran RS, Joshi UN. Agronomic evaluation of oat varieties for growth, forage yield and quality with varying levels of nitrogen under semi-arid conditions. Forage Research. 2010; 36:32-33