International Journal of Chemical Studies

P-ISSN: 2349–8528 E-ISSN: 2321–4902 IJCS 2019; 7(1): 127-129 © 2019 IJCS Received: 04-11-2018 Accepted: 08-12-2018

Asma Siddiqua

Department of Floriculture & Landscape Architecture, Horticultural Collage and Research Institute, Anantharajupeta, Dr. Y.S.R. Horticultural University, Andhra Pradesh, India

K Swarajya Lakshmi

Department of Floriculture & Landscape Architecture, Horticultural Collage and Research Institute, Anantharajupeta, Dr. Y.S.R. Horticultural University, Andhra Pradesh, India

R Nagaraju

Department of Floriculture & Landscape Architecture, Horticultural Collage and Research Institute, Anantharajupeta, Dr. Y.S.R. Horticultural University, Andhra Pradesh, India

D Srinivasa Reddy

Department of Floriculture & Landscape Architecture, Horticultural Collage and Research Institute, Anantharajupeta, Dr. Y.S.R. Horticultural University, Andhra Pradesh, India

Correspondence

Asma Siddiqua Department of Floriculture & Landscape Architecture, Horticultural Collage and Research Institute, Anantharajupeta, Dr. Y.S.R. Horticultural University, Andhra Pradesh, India

Vase life of spray chrysanthemum cultivars (*Dendrathema grandiflora* Tzvelev.) cultivated in polyhouse conditions

Asma Siddiqua, K Swarajya Lakshmi, R Nagaraju and D Srinivasa Reddy

Abstract

A study was conducted to examine the efficacy of vase solutions on postharvest life of spray chrysanthemum cultivars Terry, White Dolly, Yellow Spoon, Red Stone, Star Pink, Bronze Spoon, Paper White, Kelvin Victory and Indiana. Vase life and flower quality were significantly influenced by vase solutions. The maximum vase life (7.50 days) was observed in S₂ and its interaction S₂ C₄ recorded maximum vase life (9.50 days). C₉ recorded maximum fresh weight of flower (104.0 g) and as far as interaction is concerned, S₁C₉ recorded maximum fresh weight of flower (106.7 g). The minimum water loss (4.19 g) was recorded on day 4-5 in S₁, as far as interaction is concerned S₁ C₈ recorded minimum water uptake (8.30 days) was recorded on day 0-1 in S₁ and interaction S₁ C₄ recorded maximum water uptake (11.81 g).

Keywords: spray chrysanthemum cultivars, vase life, distilled water and 8HQC

Introduction

World trade of fresh cut flower is increasing, day by day. Cut flowers make up about one-third of the value of the global ornamental horticulture market. Fresh flowers lose their freshness and quality both during travel and also during and after arrangements due to flower specific short vase life. Such deficiencies can be ameliorated through application of nutrient additives to vase water. Use of preservative solutions has been known for many years to increase the vase life of cut flowers. Different methods like use of holding, pulsing and bud opening solution, growth regulators, gamma irradiation, precooling, cold storage, packaging etc. are already in practice in the flower trade to ensure the garden fresh quality of flowers to consumers (Singh *et al.*, 2001)^[7]. Influence of different holding solutions on chrysanthemum (*Dendrathema grandiflora* Tzvelve.) have been reported earlier (Koframek and Halevy, 1972; Marousky, 1969, 1971; Talukdar *et al.*, 2004)^[3, 5, 6, 8]. Informations on this subject are scattered. There is need to develop appropriate crop specific postharvest technology, suitable to specific agro-climatic zone, to avoid loses at the growers, florists and consumers levels. Keeping this in view, attempts were made to find out the best holding solutions for commercial exploitation to increase the vase life.

Materials and Methods

The experiment was conducted in post harvest technology laboratory at Horticultural college and research institute, located at Anantharajupeta, Kadapa district, Andhra Pradesh during the period October 2015 – March 2016. The experimental material consisted of nine spray cultivars of chrysanthemum such as Terry – anemone, White Dolly – anemone, Yellow Spoon – spoon, Red Stone – decorative, Star Pink – decorative, Bronze Spoon – spoon, Paper White – decorative, Kelvin Victory – Pompon and Indiana – pompon, was collected from the crop raised under naturally ventilated polyhouse with uniform standard cultural practices. The crop was raised under naturally ventilated polyhouse with uniform standard cultural practices. The stems were harvested with help of sharp secateurs at 8:00 am in the morning when 50 per cent of flowers were about ³/₄ th open. The stems lengths of all flowers were uniformly maintained i.e 35 cm. The cut stems were kept in two different solutions S₁ – distilled water and S₂ – sucrose (2 %) + 8HQC (500 ppm). The experiment was laid out in 2 – factorial CRD with three replications. Observations like vase life, fresh weight of flower, water uptake and transpiration loss of water.

Results and Discussion

Post harvest characteristics of cut chrysanthemum cultivars were significantly affected by two vase life solutions as well as cultivars and their interaction.

Vase life

The maximum mean vase life was observed in (7.50 days) S_2

and (8.50 days) C_4 : Red Stone. it's The interaction effects showed that interaction S_2 C_4 recorded maximum vase life (9.50 days) followed by in S_2 C_6 (8.75 days) while minimum vase life (4.00 days) in S_1 C_3 (Table 1). The variations in vase life may be due to the difference in accumulation of carbohydrates due to varied leaf production and sensitivity of cultivars to ethylene and in turn variations in these aspects might be due to genetical makeup of genotypes as reported by Vetrivel and Jawaharlal (2014)^[9].

Name of the cultivar	Distilled water (days) (S1)	Sucrose (2%) + 8HQC (500ppm) (days) (S ₂)	Mean
C ₁ : Terry	5.00	7.00	6.00
C ₂ : White Dolly	6.25	8.25	7.25
C ₃ : Yellow Spoon	4.00	7.00	5.50
C4: Red Stone	7.50	9.50	8.50
C ₅ : Star Pink	5.25	5.50	5.37
C6: Bronze Spoon	5.50	8.75	7.12
C7: Paper White	5.00	7.75	6.37
C8: Kelvin Victory	4.50	6.75	5.62
C9: Indiana	5.50	7.00	6.25
Mean	5.38	7.50	
	S.Em±	CD	
S	0.11	0.31	
С	0.23	0.67	
S x C	0.33	0.95	

Table 1: Vase life of spray chrysanthemum cultivars

Fresh weight of cut stems (g)

The cultivar, C₉ (Indiana) recorded maximum fresh weight of flower (104.0 g) followed by C₅ (Star.25 Pink) (80.08 g) and as far as interaction is concerned, S₁ C₉ recorded maximum fresh weight of flower (106.7 g) followed by S₂C₉ (101.2 g) which is statistically on par with S₁C₉ whereas minimum weight

(45.47 g) was recorded in $S_1 C_2$ on second day. (Table 2). The increase in fresh weight can happen only when the rate of water absorption is greater than transpiration rate (Rogers., 1973). Similar findings were reported by Baskaran *et al.* (2009)^[1] in chrysanthemum and Kumar *et al.* (2013)^[4] in gerbera.

Table 2: Fresh weight (g) changes during the period of vase life studies in spray chrysanthemum cultivars

Nome of the Cultivere	1 day			2 day		3 day				4 day		5 day			
Name of the Cultivars	S 1	S ₂	Mean	S1	S ₂	Mean	S 1	S ₂	Mean	S 1	S2	Mean	S 1	S ₂	Mean
C ₁ : Terry	65.87	72.77	69.32	66.45	75.05	70.51	64.39	72.55	68.57	54.47	70.19	62.33	52.25	67.50	59.87
C ₂ : White dolly	43.92	65.34	54.63	45.47	67.24	56.36	43.75	64.51	54.13	41.45	63.73	52.59	39.75	61.10	50.42
C ₃ :Yellow Spoon	65.12	58.26	61.69	67.30	59.87	63.58	65.62	58.79	62.21	63.37	57.09	60.23	60.75	54.87	57.81
C4: Red Stone	72.27	75.46	73.87	73.75	77.47	75.61	71.47	75.02	73.25	69.80	73.08	71.44	67.00	71.30	69.15
C5: Star Pink	76.85	79.66	78.25	78.85	81.31	80.08	77.30	78.55	77.92	75.12	76.94	76.03	73.00	74.27	73.63
C ₆ :Bronze Spoon	75.77	77.86	76.81	76.90	78.98	77.94	74.97	76.32	75.64	73.00	74.05	73.52	71.00	71.85	71.42
C9: Paper white	105.8	99.40	102.6	106.7	101.2	104.0	102.3	97.90	100.1	99.67	96.80	98.23	96.50	95.02	95.76
C8:Kelvin Victory	78.80	61.94	70.37	0.17	63.46	71.81	77.92	58.37	68.15	76.42	55.85	66.13	73.00	52.63	62.78
C 9: Indiana	55.25	52.07	53.66	56.17	53.57	54.87	54.37	51.30	52.83	52.10	49.47	50.78	46.50	48.25	47.37
Mean	71.08	71.42		72.42	73.13		70.24	70.39		67.26	68.58		64.41	66.30	
	$S.Em\pm$	CD	S.E	èm±	CD		S.Em±		CD	S.Em	۱±	CD	S.Em	1±	CD
S	0.27	0.77	7 0.	26	0.74		0.24		0.70	0.22	2	0.63	0.23		0.66
С	0.57	1.63	3 0.	55	1.57		0.52		1.48	0.47	7	1.35	0.49		1.41
SxC	0.81	2.31	l 0.	78	2.23		0.74		1.85	0.67	7	1.91	0.70)	1.99

Transpiration loss of water (g)

The minimum water loss (4.19 g) was recorded on throughout the vase life period upto day 4-5 in S_1 , as far as interaction is concerned $S_1 C_8$ recorded minimum water loss (3.20 g) while maximum average water loss (8.27 g) was recorded in S_2 on day 0-1 and its interaction $S_2 C_1$ showed maximum water loss (9.49 g). (Table 3). The water loss due to decline in uptake of water coupled with transpiration leads to water deficit, which ultimately reduces turgidity in cut flowers as stated by Halevy and Mayak (1981)^[2] and Baskaran *et al.* (2009)^[1].

\mathbf{T}	Table 3: Transp	iration loss of water	(g) (TLW) during the	period of vase	life studies in s	prav chrysanthemun	n cultivars
--------------	-----------------	-----------------------	----------	--------------	----------------	-------------------	--------------------	-------------

Norma of the Carltinana	Day 0-1			Day 1-2			Day 2-3				Day	3-4		Day 4-5		
Name of the Cultivars	S ₁	S_2	Mean	S_1	S_2	Mean	S ₁	S_2	Mean	S ₁	S_2	Mean	S_1	S_2	Mean	
C ₁ : Terry	11.89	9.49	10.69	8.29	7.85	8.07	4.57	6.80	5.68	5.53	5.37	5.45	5.18	5.22	5.20	
C ₂ : White dolly	8.67	8.45	8.56	10.50	7.84	9.17	4.58	6.06	5.32	6.52	5.77	6.15	5.77	5.63	5.70	
C ₃ : Yellow Spoon	54.63	9.93	7.59	6.17	9.08	7.62	6.63	8.39	7.51	5.09	7.15	6.12	4.32	6.07	5.20	
C4: Red Stone	5.24	6.58	7.07	5.14	5.65	5.40	4.66	4.87	4.76	6.54	4.57	5.55	5.28	3.52	4.40	
C5: Star Pink	7.56	8.08	5.66	1.67	7.90	4.78	5.38	7.90	6.64	5.22	5.27	5.25	3.65	5.17	4.41	
C ₆ : Bronze Spoon	3.24	9.75	8.38	10.01	8.21	9.11	9.62	7.81	8.72	3.88	6.54	5.21	3.32	6.49	4.90	
C ₉ : Paper white	7.01	8.55	7.38	6.28	7.04	6.66	2.63	6.18	4.41	4.20	6.08	5.14	3.56	5.34	4.45	
C ₈ : Kelvin Victory	6.21	7.26	5.11	5.71	6.15	5.93	3.76	4.25	4.21	4.85	3.90	4.37	3.20	3.55	3.37	
C 9: Indiana	2.96	6.30	5.30	4.05	5.24	4.64	4.13	4.87	4.50	3.61	4.59	4.10	3.45	3.55	3.50	
Mean	6.34	8.27		6.42	7.22		5.11	6.35		5.05	5.47		4.19	4.95		
	S.Em	±	CD	S.Em	±	CD	S.En	۱±	CD	S.En	ι±	CD	S.E	m±	CD	
S	0.05		0.16	0.05		0.15	0.04	1	0.11	0.04	4	0.11	0.	03	0.10	
Ĉ	0.12		0.24	0.11		0.33	0.0	3	0.24	0.08	8	0.24	0.0	08	0.22	
SxC	0.17		0.48	0.16		0.47	0.12	2	0.34	0.12	2	0.34	0.	11	0.32	

Water uptake (g)

The maximum water uptake (8.30 days) was recorded on day 0-1 in S_1 and interaction S_1 C₄ recorded maximum water uptake (11.81 g) followed by interaction S_1 C₅ (11.60 g) while minimum water uptake (3.11 g) was recorded in S_1 on day 4-5

as far as interaction is concerned, $S_1 C_7$ showed minimum water uptake (0.80 g). (Table 4). The increased water uptake maintains turgidity, freshness of flowers and thus enhances vase life owing to improved water balance and post harvest physiology as observed by Kumar *et al.* (2013)^[4] in gerbera.

Table 4: Water uptake (g) changes during the period of vase life studies in spray chrysanthemum cultivars

Nome of the Cultivore	Day 0-1			Day 1-2			Day 2-3]	Day 3	-4	Day 4-5		
Name of the Cultivars	S ₁	S_2	Mean	S ₁	S_2	Mean	S ₁	S_2	Mean	S ₁	S_2	Mean	S ₁	S_2	Mean
C ₁ : Terry	10.50	11.08	10.79	8.71	4.91	6.81	6.50	6.90	6.70	5.26	4.94	5.10	5.14	3.73	4.44
C ₂ : White dolly	5.26	3.67	4.47	5.68	7.82	6.75	5.19	8.81	7.00	4.73	8.22	6.47	3.84	7.55	5.70
C ₃ : Yellow Spoon	4.59	5.16	4.88	3.66	6.43	.04	3.07	5.37	4.22	2.17	4.93	3.55	1.36	4.78	3.07
C4: Red Stone	11.81	8.49	10.15	6.70	7.40	7.05	5.61	9.22	7.41	4.20	8.58	6.39	3.48	7.48	5.48
C5: Star Pink	11.60	9.73	10.66	8.28	8.08	8.18	6.27	5.37	5.82	5.56	3.19	4.37	4.61	2.43	3.52
C ₆ : Bronze Spoon	10.32	5.77	8.05	8.23	6.08	7.15	4.18	6.56	5.37	4.25	6.09	5.17	3.43	5.60	4.52
C ₉ : Paper white	7.54	8.50	8.02	5.34	10.48	7.91	3.79	7.46	5.62	1.99	6.78	4.38	0.80	5.53	3.16
C ₈ : Kelvin Victory	8.28	10.27	9.27	6.42	12.82	9.62	3.35	7.90	5.62	3.24	7.09	5.16	2.53	6.23	4.38
C 9: Indiana	4.79	7.47	6.13	3.67	11.27	7.47	3.11	4.75	3.93	2.51	3.10	2.80	2.82	2.61	2.71
Mean	8.30	7.79		6.30	8.37		4.56	6.93		3.77	5.88		3.11	5.10	
	S.Em	±	CD	S.En	n±	CD	S.En	۱±	± CD		۱±	CD	S.Em±		CD
S	0.02		0.07	0.02		0.07	0.0	1	0.04	0.01	1	0.04	0.04		0.12
С	0.05		0.16	0.0	5	0.15	0.03	3	0.09	0.03	3	0.10	0.09)	0.25
SxC	0.08		0.22	0.0	7	0.22	0.04	1	0.13	0.05	5	0.14	0.12	2	0.36

References

- 1. Baskaran V, Janakiram T, Jayanthi R. Evaluation of post harvest quality of some cultivars of chrysanthemum. Journal of Ornamental Horticulture. 2009; 12(1):59-61.
- 2. Halevy AH, Mayak S. Senescence and post harvest physiology of cut flowers, Part-II. Hort. Rev. 1981; 3:59-143.
- 3. Koframek AM, Halevy AM. Conditions for opening cut chrysanthemum flower buds. Journal of American Society for horiticultural Science. 1972; 97(5):578-584.
- 4. Kumar R, Ahmed N, Sharma OC, Mahendiran G, Lal S. Screening of gerbera (*Gerbera jamesonii*) cultivars for quality, vase life and stem bending. Progressive Horticulture. 2013; 45(2):317-320.
- Marousky FJ. Influence of various commercial floral preservatives and 8 hydroxyquinoline citratenplus sucrose on development and lasting ability of flower buds of several chrysanthemum cultivars. Proceedings of Florida State./wrticultural Society, 1969; 82:381-385.
- 6. Marousky FJ. Handling and opening bud cut chrysanthemum flowers with 8hydroxyquinoline citrate

and sucrose. Marketing Research Report, United States Development of Agriculture. 1971; 90S:14.

- 7. Singh K, Arora JS, Bhattacharjee SK. Postharvest management of cut flowers. All India Co-ordinated Research Project on Floriculture, Indian Council of Agricultural Research, New Delhi, Technical Bulletin, 2001, 10.
- Talukdar MC, Sharma BJ, Das S, Mahanta SJ. Eff~ct of pulsing on post harvest quality' of chrysanthemum (*Dendrathema grandiflora*, Tzvelve.) under different storage conditions. Journal of Ornamental Horticulture. 2004; 7(2):133-140.
- 9. Vetrivel T, Jawaharlal M. Evaluation of chrysanthemum (*Dendranthema grandiflora* Tzvelev) varieties for yield and quality under subtropical hills. Trends in Biosciences. 2014; 7(14):1812-1815.