



P-ISSN: 2349-8528

E-ISSN: 2321-4902

www.chemijournal.com

IJCS 2022; 10(1): 85-89

© 2022 IJCS

Received: 13-11-2021

Accepted: 15-12-2021

Raiyani VN

Ph.D. Scholar, Department of
Agronomy, BACA, Anand
Agricultural University, Anand,
Gujarat, India

Dr. Mevada KD

Professor and Head (Agronomy),
Gujarat Organic Agricultural
University, Anand, Gujarat,
India

Vaghela GM

SRA, Gujarat Organic
Agricultural University, Anand,
Gujarat, India

Patil YG

Ph.D. Scholar, Department of
Agronomy, BACA, Anand
Agricultural University, Anand,
Gujarat, India

Corresponding Author:**Dr. Mevada KD**

Professor and Head (Agronomy),
Gujarat Organic Agricultural
University, Anand, Gujarat,
India

Influence of potash management on yield and economics of *rabi* maize (*Zea mays* L.)

Raiyani VN, Dr. Mevada KD, Vaghela GM and Patil YG

Abstract

A field experiment was conducted at College Agronomy Farm, B. A. College of Agriculture, Anand Agricultural University, Anand, Gujarat, India during *rabi*-summer seasons of the years 2019-20 and 2020-21 to study the influence of potash management on yield and economics of *rabi* maize (*Zea mays* L.) with different potash management practices *viz*: Control (Without potash fertilizers), 30 kg K₂O /ha through MOP or SOP or Schoenite with and without KMB @ 1 lit/ha, 45 kg K₂O /ha through MOP or SOP or Schoenite with and without KMB @ 1 lit/ha and Foliar spray of Schoenite @ 1.5% at 30 and 60 DAS. The soil of the experimental plot was loamy sand in texture, marginal in organic carbon, and medium in available phosphorus and potash. Results revealed that 45 kg K₂O /ha through MOP or SOP or Schoenite along with KMB @ 1 lit/ha recorded significantly higher growth and yield attributes *viz*: plant height at 60 DAS and at harvest, cob length and cob girth. They also produced significantly higher grain and stover yield over rest of the treatments. However, higher net monetary returns of ₹ 101643/ha and BCR of 4.39 was fetched under 45 kg K₂O/ha through MOP + KMB @ 1 lit/ha.

Keywords: potash, maize, MoP, SoP, Schoenite, KMB

1. Introduction

Being C₄ plant, maize has the highest genetic yield potential amongst the cereals owing to its better dry matter accumulation efficiency in a unit area and time, which is why, it often refers to as “Miracle Crop”, “King of Grain Crops”, “Backbone of America” and “Queen of Cereals”, as together with rice and wheat, it provides approximately 30% of the food calories to more than 4.5 billion people in 94 developing countries and the demand for maize in these countries is assumed to double by 2050 (Johnston *et al.*, 2011) [6]. Worldwide it occupies an area of about 184 million ha covering 160 countries providing around 36 per cent towards the global food grain production. In India maize has been emerged as the third most important cereal crop, after rice and wheat, occupying an area of 9.60 million ha with the production of 27.15 million tones, having average productivity of about 2.8 tones/ha, whereas in Gujarat, 0.44 million ha area is covered with a production of 0.68 million tones having productivity of 1659 kg/ha (Anon., 2020) [1], which is quite below the national and world average. One of the major factors responsible for lower yield is imbalance fertilization, particularly potassium. Maize has a high production potential as an exhaustive crop for potassium fertilizer. Kusro *et al.*, (2014) [7] reported that more than 400 kg K₂O/ha of potassium is taken up by maize crop under intensive cropping system.

Among potassium sources, muriate of potash (MOP) is the only fertilizer that totally imported in India. Among common potassic fertilizers, sulphate of potash, is mostly used by the majority of growers only because of its low salt index, non-hygroscopic and chlorine free K-fertilizer compared with muriate of potash, but the cost is expensive. Indigenously, Central Salt and Marine Chemicals Research Institute, Bhavnagar, Gujarat and some private agri-input agencies have developed the extraction process of sulphate of potash (K₂SO₄) and schoenite (K₂SO₄, MgSO₄) from sea bittern. Among them, indigenously produced double salts of K, *i.e.*, potassium schoenite (molar mass 415 g/mol) is appropriate for use in sandy and alluvial soils providing a readily available supply of Potassium, Magnesium and Sulfur to growing plants in an ideal ratio (Rathore *et al.*, 2014) [14]. Likewise, inoculation of maize plants with *Bacillus mucilaginosus*, *Azotobacter chroococcum* and *Rhizobium* resulted in significant higher mobilization of potassium from waste mica, which in turn act as a source of potassium for plant growth. *Bacillus mucilaginosus* and *Bacillus edaphicus* are the most commonly used Potassium solubilising bacteria (KSB) used as biofertilizer. Potassium rock is solubilised by

KSB through production and secretion of organic acids (Han and Lee, 2005) [3]. However, information pertaining to potassium management in maize through different rates and sources of potassium is unknown. With this background present study was conducted to know the effect of different management practices of K on yield and economics of maize crop.

Materials and Methods

A field experiment was conducted at Agronomy Farm, Anand Agricultural University, Anand during the years 2019-20 and 2020-21 to find out influence of different potash management practices on yield and economics of *rabi* maize (*Zea mays* L.). The experimental site was loamy sand in texture, alkaline in nature (8.23 pH) with low soluble salts (0.18 dS/m), organic carbon (0.36%) and available nitrogen (142 (kg/ha)), medium in available phosphorus (36 kg/ha) and available potassium (227 kg/ha). Fourteen potassium management treatments comprising of Control (Without potash fertilizers), 30 kg K₂O /ha through MOP or SOP or Schoenite with and without KMB @ 1 lit/ha, 45 kg K₂O /ha through MOP or SOP or Schoenite with and without KMB @ 1 lit/ha and Foliar spray of Schoenite @ 1.5% at 30 & 60 DAS were studied under Randomized Block Design (RBD). The maize variety GAYMH 3 (Gujarat Anand Yellow Maize Hybrid) was taken for the experiment. Recommended dose of nutrients i.e., 150:60:00 NPK kg/ ha were applied through fertilizers uniformly in the furrows, wherein, 50% of recommended dose of nitrogen (RDN) and 100% recommended dose of phosphorus were applied as basal, whereas remaining 50% RDN was applied in two equal splits at 30 and 60 DAS. The nitrogen was applied through urea and phosphorus was applied through DAP. Potash fertilizer was applied as per treatment. Potash

mobilizing bacteria (KMB) treatment was applied to the soil after sowing with irrigation as per treatment. Foliar spray of Schoenite @ 1.5% was applied at 30 and 60 DAS. Maize crop was sown at 60 cm x 20 cm spacing in experimental plot. Shelling percentage was calculated using following formula.

$$\text{Shelling percentage} = \frac{\text{Weight of grain}}{\text{Weight of cob with shells}} \times 100$$

The harvest index for each treatment was worked out by using formula given by Donald and Hamblin, 1976.

$$\text{HI (\%)} = \frac{\text{Economic yield (kg/ ha)}}{\text{Economic yield (kg/ ha)} + \text{Biological yield (kg/ ha)}} \times 100$$

The benefit: cost ratio was calculated on the basis of formula given below.

$$\text{BCR} = \frac{\text{Total income (₹/ha)}}{\text{Total expenditure (₹/ha)}} \times 100$$

Results and Discussion

The data pertaining to various growth parameter, yield attributes, yield and economics of maize as influenced by different potash management practices with their statistical inference are presented and discussed as under:

A. Growth parameters

The mean data pertaining to periodical plant height of maize measured periodically at 30, 60 DAS and at harvest as influence by different potash management practices.

Table 1: Periodical plant height (cm) of maize as influenced by different treatments of potash (pooled over two years)

Treatments	At 30 DAS	At 60 DAS	At harvest
T ₁ : Control	61.25	136.89 ^e	159.77 ^e
T ₂ : 30 kg/ha K ₂ O through MOP	64.81	145.64 ^{efg}	164.17 ^{de}
T ₃ : 30 kg/ha K ₂ O through SOP	63.44	142.44 ^{fg}	161.33 ^e
T ₄ : 30 kg/ha K ₂ O through Schoenite	62.46	144.74 ^{efg}	162.83 ^{de}
T ₅ : 30 kg/ha K ₂ O through MOP + KMB @ 1 lit/ha	70.63	163.12 ^{abcd}	186.50 ^{abc}
T ₆ : 30 kg/ha K ₂ O through MOP SOP + KMB @ 1 lit/ha	69.33	160.89 ^{abcde}	182.33 ^{abcd}
T ₇ : 30 kg/ha K ₂ O through Schoenite + KMB @ 1 lit/ha	71.09	165.07 ^{abc}	187.17 ^{abc}
T ₈ : 45 kg/ha K ₂ O through MOP	67.67	156.58 ^{abcdef}	172.00 ^{bcde}
T ₉ : 45 kg/ha K ₂ O through SOP	66.89	149.50 ^{defg}	167.83 ^{cde}
T ₁₀ : 45 kg/ha K ₂ O through Schoenite	69.83	153.88 ^{bcdef}	169.50 ^{cde}
T ₁₁ : 45 kg/ha K ₂ O through MOP + KMB @ 1 lit/ha	73.69	170.64 ^a	196.33 ^a
T ₁₂ : 45 kg/ha K ₂ O through SOP + KMB @ 1 lit/ha	71.57	167.08 ^{ab}	190.17 ^{ab}
T ₁₃ : 45 kg/ha K ₂ O through Schoenite + KMB @ 1 lit/ha	73.00	170.45 ^a	192.83 ^a
T ₁₄ : Foliar spray of Schoenite @ 1.5% at 30 & 60 DAS	66.08	147.53 ^{defg}	166.02 ^{de}
S.Em. ±	1.92	5.08	6.06
C. D. at 5%	NS	Sig.	Sig.
C.V. %	6.92	8.02	8.43

The plant height (Table 1) at 30 DAS did not exert any significant effect in different potash management treatments. However, at later stages, significantly higher plant height of 170.64 cm and 196.33 cm were recorded under the application of 45 kg K₂O/ ha through MOP + KMB @ 1 lit/ha at 60 DAS and at harvest, respectively. This treatment exhibited at par relations under the treatments T₅, T₆, T₇, T₈, T₉, T₁₀, T₁₂, T₁₃ and T₁₄ at 60 DAS and with treatments T₁₃, T₁₂, T₅, T₆ and T₇ at harvest. Potassium is most pertinent nutrient that has the potential to enhance cell membranes and the generation of turgor in plants. It is also the major cation for the maintenance of cation-anion balances (Rajarithnam,

2006) [13]. Application of potassium at various rates and sources along with KMB might improve the K efficiency and availability to maize. Han *et al.* (2006) [4] reported that KMB plays an important role in mobilizing potassium to the crop. Potassium activates many enzymes and plays an important role in the maintenance of electrical potential genetics. This might be resulted into longer plants at 45 kg K₂O/ha along with KMB @ 1 lit/ha. Similar results were reported by Shinde *et al.* (2011) [16] and Jadhav *et al.* (2012) [5].

B. Yield attributes

Data pertaining to yield attributes of maize as influenced by different potash management on pooled basis are presented in Table 2.

Results revealed that though number of rows per cob, seed index and shelling percentage of maize did not differ remarkably due to different potash management practices on pooled basis, significant differences on cob length, cob girth and harvest index were observed. During pooled analysis significantly higher cob length (25.92 cm) was recorded under application of 45 kg K₂O/ha through MOP + KMB @ 1 lit/ha, but it was comparable with T₁₃ and T₁₂ treatments. Same trend was observed for cob girth and being at par for the treatments T₂, T₃, T₄, T₁ and T₁₄ significantly higher cob girth (19.51 cm) was reported under the same treatment. Harvest index was also observed significantly higher under application of 45 kg

K₂O/ha through MOP + KMB @ 1 lit/ha. The significant increase in cob length and cob girth under the treatment might be the result of adequately higher mineralization of potassium at higher dose (45kg/ha), irrespective of the sources, in presence of KMB which might cause cell elongation, root development, enhance accumulation of photosynthates from source to sink and ultimately increase the cob length and cob girth. The results were in conformity with Pathak *et al.* (2002)^[11], Louraduraj (2006)^[28] and Zakir *et al.* (2019)^[17].

In pooled analysis, significantly higher harvest index of maize (42.81%) was recorded under the same treatment of 45 kg K₂O/ha through MOP + KMB @ 1 lit/ha (T₁₁). However, it was found to be substantially at par with all the treatments except T₁, T₂, T₃ and T₄. As harvest index is a ratio of economic yield to the total biological yield, the results are on calculation basis.

Table 2: Influence of different potash management practices on yield attributes of maize (pooled over two years)

Treatments	Cob length (cm)	Cob girth (cm)	Number of rows per cob	Seed Index (g)	Harvest index (%)	Shelling Percentage
T ₁ : Control	17.87 ^f	15.10 ^d	15.04	19.28	39.20 ^e	73.44
T ₂ : 30 kg/ha K ₂ O through MOP	21.17 ^{cde}	15.82 ^{cd}	15.52	20.15	41.53 ^{bcd}	74.32
T ₃ : 30 kg/ha K ₂ O through SOP	19.73 ^{def}	15.55 ^{cd}	15.31	19.62	41.26 ^{cd}	73.60
T ₄ : 30 kg/ha K ₂ O through Schoenite	19.23 ^{ef}	14.94 ^d	15.60	19.59	41.06 ^d	74.19
T ₅ : 30 kg/ha K ₂ O through MOP + KMB @ 1 lit/ha	22.67 ^{bc}	18.10 ^{ab}	16.65	21.52	42.48 ^{ab}	76.28
T ₆ : 30 kg/ha K ₂ O through MOP SOP + KMB @ 1 lit/ha	22.24 ^{bcd}	17.70 ^{abc}	16.28	21.26	42.27 ^{abc}	76.20
T ₇ : 30 kg/ha K ₂ O through Schoenite + KMB @ 1 lit/ha	23.17 ^{bc}	18.15 ^{ab}	16.81	21.83	42.52 ^{ab}	76.56
T ₈ : 45 kg/ha K ₂ O through MOP	21.06 ^{cde}	17.05 ^{bcd}	15.85	21.06	41.86 ^{abcd}	75.31
T ₉ : 45 kg/ha K ₂ O through SOP	20.50 ^{cdef}	16.57 ^{bcd}	15.77	20.90	41.76 ^{abcd}	74.99
T ₁₀ : 45 kg/ha K ₂ O through Schoenite	21.83 ^{cde}	16.83 ^{bcd}	15.79	20.62	41.78 ^{abcd}	75.08
T ₁₁ : 45 kg/ha K ₂ O through MOP + KMB @ 1 lit/ha	25.92 ^a	19.51 ^a	17.76	23.24	42.81 ^a	78.15
T ₁₂ : 45 kg/ha K ₂ O through SOP + KMB @ 1 lit/ha	23.39 ^{abc}	18.43 ^{ab}	17.51	22.19	42.61 ^{ab}	77.05
T ₁₃ : 45 kg/ha K ₂ O through Schoenite + KMB @ 1 lit/ha	24.74 ^{ab}	18.62 ^{ab}	17.62	22.72	42.66 ^a	77.13
T ₁₄ : Foliar spray of Schoenite @ 1.5% at 30 & 60 DAS	19.70 ^{def}	15.55 ^{cd}	15.66	20.15	41.75 ^{abcd}	74.49
S.Em. ±	0.68	0.46	0.85	2.11	0.46	0.46
C. D. at 5%	Sig.	Sig.	NS	NS	Sig.	NS
C.V. %	10.02	9.88	6.97	7.21	1.91	6.85
C.V. %	10.02	9.88	6.97	7.21	1.91	6.85

C. Grain and Stover Yields

Data pertaining to grain and stover yield of maize as influenced by different potash management practices are presented in Table 3. Data showed that different potash treatments significantly influenced the grain yield of maize during both the years (2019-20 and 2020-21) as well as in pooled analysis. Among all the treatments tested, treatment with 45 kg K₂O/ha through MOP + KMB @ 1 lit/ha had produced apparently higher grain yield of 5859 kg/ha and 5796 kg/ha during the year 2019-20 and 2020-21, respectively. However, it remained comparable with treatments T₁₀, T₁₂, T₁₃, T₅, T₆, T₁₄, T₇, T₈ and T₉ and treatments T₁₀, T₁₂, T₁₃, T₅, T₆, T₁₄, T₇, T₈, T₂ and T₉, during the year 2019-20 and 2020-21, respectively.

Results of pooled analysis followed the same trend and revealed that significantly higher grain yield of 5828 kg/ha was recorded under the application of 45 kg K₂O/ha through MOP + KMB @ 1 lit/ha (T₁₁), which was closely related to the treatments T₁₂, T₁₃, T₅, T₆ and T₇. Treatment T₁₁ produced 59% higher grain yield compared to control (T₁) during the pooled analysis. The results also indicated that treatment with the same rate (45 kg/ha) with SOP and Schoenite as the sources of potash also responded alike to MOP along with KMB. Similarly, all the sources of potash along with KMB at 30 kg/ha were also found analogous with T₁₁. It was also

further pointed out that treatments with KMB had significant influence on yield irrespective of rates (30 or 45 kg/ha) and sources (MoP/SoP/Schoenite) might be due to higher mineralization accrued to synergistic impact of KMB on the efficacy of potash.

Grain yield is a consequential effect of cumulative impact of all the growth and yield attributes. This impact was reflected in higher growth and yield attributes like plant height, cob length, cob girth, number of rows per cob and seed index obtained under the same treatment T₁₁. These results are in conformity with findings of Pathak *et al.* (2002)^[11], Han *et al.* (2006)^[4], Sharma (2011)^[15] and Patil *et al.* (2017)^[12] on maize.

The results of pooled analysis showed that significantly higher stover yield (7683 kg/ha) of maize recorded under application of 45 kg K₂O/ha through Schoenite + KMB @ 1 lit/ha (T₁₃). However, it was closely related to treatments T₁₁, T₁₂, T₅, T₆ and T₇. An increase over control was reported to the tune of 38.88, 37.11 and 37.26% under T₁₁ during the year 2019-20, 2020-21 and in pooled analysis, respectively. An increase in stover yield under the treatments is mainly attributed to growth attributing parameter like plant height. The results are in close conformity with those of Pandey and Avasthi (2014)^[10], Maske *et al.* (2015)^[9] and Patil *et al.* (2017)^[12].

Table 3: Influence of different potash management on yields of maize

Treatments	Grain yield (kg/ha)			Stover yield (kg/ha)		
	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled
T ₁ : Control	3665 ^d	3623 ^d	3644 ^f	5644 ^e	5551 ^c	5597 ^f
T ₂ : 30 kg/ha K ₂ O through MOP	4828 ^{bc}	4822 ^{abc}	4825 ^{de}	6807 ^{bcd}	6750 ^{ab}	6779 ^{de}
T ₃ : 30 kg/ha K ₂ O through SOP	4622 ^c	4641 ^{bc}	4631 ^e	6549 ^{cd}	6568 ^{ab}	6559 ^e
T ₄ : 30 kg/ha K ₂ O through Schoenite	4541 ^c	4508 ^c	4525 ^e	6491 ^d	6436 ^{bc}	6463 ^e
T ₅ : 30 kg/ha K ₂ O through MOP + KMB @ 1 lit/ha	5554 ^{ab}	5507 ^{ab}	5530 ^{abc}	7533 ^{ab}	7435 ^{ab}	7484 ^{abc}
T ₆ : 30 kg/ha K ₂ O through MOP SOP + KMB @ 1 lit/ha	5406 ^{abc}	5350 ^{abc}	5378 ^{abcd}	7385 ^{abc}	7278 ^{ab}	7332 ^{abcd}
T ₇ : 30 kg/ha K ₂ O through Schoenite + KMB @ 1 lit/ha	5629 ^{ab}	5523 ^{ab}	5576 ^{abc}	7608 ^{ab}	7451 ^{ab}	7530 ^{abc}
T ₈ : 45 kg/ha K ₂ O through MOP	5058 ^{abc}	5022 ^{abc}	5040 ^{bcd}	7037 ^{abcd}	6950 ^{ab}	6994 ^{bcd}
T ₉ : 45 kg/ha K ₂ O through SOP	4977 ^{abc}	4959 ^{abc}	4968 ^{cde}	6956 ^{abcd}	6887 ^{ab}	6921 ^{cde}
T ₁₀ : 45 kg/ha K ₂ O through Schoenite	4975 ^{abc}	4973 ^{abc}	4974 ^{cde}	6954 ^{abcd}	6901 ^{ab}	6928 ^{cde}
T ₁₁ : 45 kg/ha K ₂ O through MOP + KMB @ 1 lit/ha	5859 ^a	5796 ^a	5828 ^a	7838 ^a	7391 ^a	7615 ^{ab}
T ₁₂ : 45 kg/ha K ₂ O through SOP + KMB @ 1 lit/ha	5649 ^{ab}	5624 ^{ab}	5637 ^{ab}	7628 ^{ab}	7552 ^a	7590 ^{ab}
T ₁₃ : 45 kg/ha K ₂ O through Schoenite + KMB @ 1 lit/ha	5714 ^{ab}	5683 ^a	5698 ^a	7755 ^a	7611 ^a	7683 ^a
T ₁₄ : Foliar spray of Schoenite @ 1.5% at 30 & 60 DAS	4965 ^{abc}	4973 ^{abc}	4969 ^{cde}	6944 ^{abcd}	6901 ^{ab}	6923 ^{cde}
S.Em. ±	268	297	260	265	307	203
C. D. at 5%	Sig.	Sig.	Sig.	Sig.	Sig.	Sig.
C.V. %	9.11	10.14	9.64	6.48	7.40	7.07

Table 4: Economics of maize as influenced by various potash management practices (Av. Of two years)

Treatment	Yields (kg/ha)		Fixed cost (₹/ha)	Treatment cost (₹/ha)	Total cost of cultivation (₹/ha)	Gross realization (₹/ha)	Net realization (₹/ha)	BCR
	Grain	Stover						
T ₁ : Control	3644	5597	28663	00	28663	83032	54369	2.90
T ₂ : 30 kg/ha K ₂ O through MOP	4825	6779	28663	800	29463	109310	79847	3.71
T ₃ : 30 kg/ha K ₂ O through SOP	4631	6585	28663	1920	30583	104994	74411	3.43
T ₄ : 30 kg/ha K ₂ O through Schoenite	4525	6478	28663	3640	32303	102634	70331	3.18
T ₅ : 30 kg/ha K ₂ O through MOP + KMB @ 1 lit/ha	5530	7484	28663	920	29583	124997	95414	4.23
T ₆ : 30 kg/ha K ₂ O through MOP SOP + KMB @ 1 lit/ha	5378	7332	28663	2040	30703	121615	90912	3.96
T ₇ : 30 kg/ha K ₂ O through Schoenite + KMB @ 1 lit/ha	5576	7530	28663	3760	32423	126020	93597	3.89
T ₈ : 45 kg/ha K ₂ O through MOP	5040	6994	28663	1200	29863	114094	84231	3.82
T ₉ : 45 kg/ha K ₂ O through SOP	4968	6921	28663	2880	31543	112491	80948	3.57
T ₁₀ : 45 kg/ha K ₂ O through Schoenite	4974	6928	28663	5488	34151	112626	78475	3.30
T ₁₁ : 45 kg/ha K ₂ O through MOP + KMB @ 1 lit/ha	5828	7781	28663	1320	29983	131626	101643	4.39
T ₁₂ : 45 kg/ha K ₂ O through SOP + KMB @ 1 lit/ha	5637	7590	28663	3000	31663	127376	95713	4.02
T ₁₃ : 45 kg/ha K ₂ O through Schoenite + KMB @ 1 lit/ha	5698	7652	28663	5608	34271	128735	94464	3.76
T ₁₄ : Foliar spray of Schoenite @ 1.5% at 30 & 60 DAS	4969	6923	28663	776	29439	112514	83075	3.82

Selling price: Seed: ₹ 21.25/ kg, Stover: ₹ 1.0 / kg

D. On Economics

Economics as influenced by different potassium management practices for maize crop comprised of cost of cultivation, gross income, net realization and B:C ratio obtained on hectare basis during the course of investigation are presented in Table 4.

An appraisal of data presented in Table 4 revealed that maximum net realization of ₹ 101643/ ha was fetched under application of 45 kg K₂O /ha through MOP + KMB @ 1 lit/ha (T₁₁) followed by treatment 45 kg K₂O /ha through SOP + KMB @ 1 lit/ha (T₁₂) of ₹ 95713 / ha. The lowest net realization of ₹ 54369/ ha was realized under treatment observed in control treatment(T₁).

In case of benefit cost ratio, the highest benefit cost ratio of 4.39 was recorded under application of 45 kg K₂O /ha through MOP + KMB @ 1 lit/ha (T₁₁) followed by T₅ i.e. 30 kg K₂O /ha through SOP + KMB @ 1 lit/ha (4.23) and T₁₂ i.e. 45 kg K₂O /ha through SOP + KMB @ 1 lit/ha (4.02).

Conclusion

In light of the two years' field experiment, it can be concluded that higher growth, yield attributes, yield and economics of *rabi* maize could be achieved with an application of 45 kg K₂O /ha either through MOP/SOP/Schoenite along with Potash Mobilizing Bacteria (KMB) 1 lit/ha, however, it was found comparable with 30 kg K₂O /ha either through

MOP/SOP/Schoenite along with Potash Mobilizing Bacteria (KMB) 1 lit/ha.

References

- Anonymous. Ministry of Agriculture, Directorate of economics and statistics. 2020. Website: <http://www.agricoop.nic.in>
- Donald CM, Hamblin J. The biological yield and harvest index of cereals as agronomic and plant breeding criteria. *Advance in Agronomy*. 1976;28:361-405.
- Han HS, Lee KD. Phosphate and potassium solubilizing bacteria effect on mineral uptake, soil availability and growth of eggplant. *Research Journal of Agriculture and Biological Sciences*. 2005;1(2):176-180.
- Han HS, Supanjani, Lee KD. Effect of co-inoculation with phosphate and potassium solubilizing bacteria on mineral uptake and growth of pepper and cucumber. *Plant, Soil and Environment*. 2006;52(3):130-136.
- Jadhav KL, Bhilare RL, Kunjir NT. Influence of integrated nutrient management on growth and yield of maize. *Journal of Agricultural Research and Technology*. 2012;37(2):344-346.
- Johnston AM, Khurana HS, Majumdar K, Satyanarayana T. Site specific nutrient management – Concept, current research and future challenges in Indian agriculture.

- Journal of the Indian Society of Soil Science. 2011;57(1):1-10.
7. Kusro PS, Singh DP, Paikra MS, Kumar D, Viashwavidyalaya IGK. American International Journal of Research in Formal, Applied and Natural sciences. 2014;5(1):51-53.
 8. Jagajjanani G, HariPriya A. Formulation and evaluation of sweet corn milk (*Zea Mays*) based mayonnaise. Int. J Agric. Food Sci. 2021;3(1):33-38. DOI: 10.33545/2664844X.2021.v3.i1a.49
 9. Maske NM, Pawar SB, Munde GR, Patange MJ. Integrated nutrient management and irrigation schedules on growth and yield of *rabi* maize. Bioinfolet. 2015;12(3):622-623.
 10. Pandey KK, Awasthi A. Integrated nutrient management in the maize (*Zea mays* L.) yield and soil properties. International Journal of Agricultural Sciences. 2014;10(1):244-246.
 11. Pathak SK, Singh SB, Singh SN. Effect of integrated nutrient management on growth, yield and economic in maize (*Zea mays* L.) - wheat (*Triticum aestivum* L.) cropping system. Indian Journal of Agronomy. 2002;47(3):325-332.
 12. Patil S, Basavaraja PK, Sheshadri T. Effect of Different Sources and Levels of K on Maize (*Zea mays* L.) yield, nutrient content and uptake by maize crop in low K soils of Eastern Dry Zone of Karnataka, India. International Journal of Current Microbiology and Applied Science, 2017;6(8):577-587.
 13. Rajarathinam S. Effect of higher application of nitrogen and potassium over recommended level on growth, yield and yield attributes of late sown winter maize (*Zea mays* L.). Crop Research. 2006;26(1):71-74.
 14. Rathore SS, Chaudhary DR, Bhatt BP. Schoenite and potassium sulphate: Indigenous potassic fertilizer for rainfed groundnut. Indian journal of traditional knowledge. 2014;13(1):222-226.
 15. Sharma JP. Economy in fertilizer use through organic manures in growing maize. Indian Journal of Agronomy. 2011;28:154-155.
 16. Shinde PD, Jadhav AS, Shaikh AA. Effect of integrated nutrient management and row spacing on growth and yield of composite maize (*Zea mays* L.). International Journal of Science and Research. 2011;4(2):65-70.
 17. Zakir M, Singh D, Yadav S. Effect of different levels of potassium on yield and yield attributes of *kharif* maize (*Zea mays* L.). International Journal of Current Microbiology and Applied Sciences. 2019;8(1):2054-2060.
 18. Anonymous Food and agriculture organization (FAOSTAT). 2007. Website: <http://www.fao.org/faostat/data>.
 19. Cochran WG, Cox GM. Experimental designs. Wiley New York, USA. 1967.
 20. Department of Fertilizer, Government of India. 2016. <https://fert.nic.in/publication-reports/annual-report>
 21. Jackson ML. Soil chemical analysis. Prentice hall of India Pvt. Ltd. 1973.
 22. Meena HN, Yadav RS, Bhaduri D. Effects of potassium application on growth of peanut (*Arachis hypogaea* L.) and ionic alteration under saline irrigation. Indian Journal of Agronomy. 2018;63(1):95-103.
 23. Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. Circulation from USDA. 1954, 939.
 24. Ombase KC. Site Specific Nutrient Management in *rabi* maize and their residual effect on summer green gram. Ph. D. thesis submitted to the Anand Agricultural University, 2018.
 25. Panwar JDS, Singh O. Response of *Azospirillum* and *Bacillus* on growth and yield of wheat under field conditions. Indian Journal of Plant Physiology. 2000;5(1):108-110.
 26. Piper CS. Soil and Plant Analysis. Hans Publishers, Bombay. 1966, 137-153.
 27. Walkey A, Black IA. Estimation of organic carbon by the chromic acid titration method. Soil Science. 1934;37:29-38.
 28. Louraduraj AC. Identification of optimum quantity of vermicompost for maize under different levels of fertilization. Journal of Ecobiology. 2006;18:23-27.