



P-ISSN: 2349-8528

E-ISSN: 2321-4902

[www.chemijournal.com](http://www.chemijournal.com)

IJCS 2023; 11(4): 28-30

© 2023 IJCS

Received: 03-06-2023

Accepted: 07-07-2023

**Garima Kaushik Parashar**

Research Scholar, Department of Agronomy, School of Agricultural Sciences, Shri Guru Ram Rai University, Dehradun, Uttarakhand, India

**Vidhi Chamoli**

Research Scholar, Plant Pathology Department, School of Agricultural Sciences, Shri Guru Ram Rai University, Dehradun, Uttarakhand, India

**Shakshi Dubey**

Assistant Professor, Department of Agriculture, GRDIMIT RAJPUR ROAD Dehradun, Uttarakhand, India

**Shivani Khatri**

Assistant Professor, Department of Agriculture, GRDIMIT Rajpur Road, Dehradun, Uttarakhand, India

**Corresponding Author:****Garima Kaushik Parashar**

Research Scholar, Department of Agronomy, School of Agricultural Sciences, Shri Guru Ram Rai University, Dehradun, Uttarakhand India

## Effect of different organic manure with Beejamrit inoculation on growth of summer moong (*Vigna radiata* L.)

**Garima Kaushik Parashar, Vidhi Chamoli, Shakshi Dubey and Shivani Khatri**

### Abstract

The present study aimed to evaluate the Effect of different organic manure with Beejamrit inoculation on growth of summer moong (*Vigna radiata* L.) conducted during summer season 2021 at SGRRU Pathri Bagh Dehradun. The experiment was laid out into Randomized block design with 3 replication and 12 treatments namely T<sub>0</sub> Control, T<sub>1</sub> 100% N Through FYM+ Beejamrit inoculation, T<sub>2</sub> 100% N through Vermicompost + Inoculation, T<sub>3</sub> 100% N through Shivansh Khad + Beejamrit inoculation, T<sub>4</sub> 100% N through Neem cake+ Inoculation, T<sub>5</sub> 50% N through FYM +50% Vermicompost+ Beejamrit inoculation, T<sub>6</sub> 50% N through FYM +50% Shivansh Khad + Inoculation, T<sub>7</sub> 50% N through FYM +50% Neem cake+ Beejamrit inoculation, T<sub>8</sub> 50% N through Vermicompost + 50% Shivansh Khad + Inoculation, T<sub>9</sub> 50% N through Vermicompost + 50% Neem cake+ Beejamrit inoculation, T<sub>10</sub> 25% FYM + 25% Vermicompost + 25% Shivansh Khad +25% Neem cake+ Inoculation, T<sub>11</sub> 25% FYM + 25% Vermicompost +25% Shivansh Khad +25% Neem cake+ Beejamrit inoculation. The result revealed that application of T<sub>5</sub> 50% N through FYM +50% Vermicompost+ Beejamrit inoculation recorded significantly higher plant height (41.53 cm), number of branches per plant (11.46) and dry weight (8.65 g) as compare to rest of the treatment except control.

**Keywords:** Beejamrit inoculation, Shivansh Khad, organic manure, vermicompost, neem cake

### Introduction

In Asia, where 90% of the world's production currently takes place, the mungbean (*Vigna radiata* (L.) Wilczek) is one of the most significant food legumes. Total area protected at a yield of 694 kg/ha, the area under pulses in India was around 23 Mha, producing a total of 25.23 Mt. Mungbean cultivation covered 2.93 Mha in India in 2017-18, and 2.01 Mt were produced there (DASE 2018) [1]. According to Meena and Yadav (2015) [3] and Meena *et al.* 2020a [4], it produces on average just 436 kg/ha. Because pulse crops have a biological nitrogen fixation (BNF) mechanism built into them, they can meet 80–90% of their nitrogen (N) needs. Most pulse crops can therefore be supplied with a beginning dose of 15–25 kg N/ha. A few years ago, micronutrients such as zinc (Zn) and iron (Fe), pulse crop yield has increased significantly in many areas (Meena and Lal 2015) [3]. Organic sources that are useful for enhancing soil characteristics have been found to supply nutrients for a long time without negatively impacting the soil. Organic farming will produce high-quality food without having a negative impact on the environment or soil health (Yadav *et al.*, 2013) [5]. Application of organic manure maintains soil fertility while preserving the soil ecosystem through optimal soil biological activity (Meena *et al.* 2017) [6].

Compost and farmyard manure (FYM) are the most significant and popular bulky organic manures. Since FYM includes 0.5-1.5, 0.4-0.8, and 0.5-1.9 percent of N, P, and K, respectively, it feeds N, P, and K to the plant in an accessible form through mineralization.

All of the plant nutrients are abundant in vermicompost. When compared to FYM, it has a larger nutritional content. Both nitrogen fixation and P solubilization are aided by it. It is a rich source of nutrients including nitrogen (1.6%), phosphorus (0.54%), potassium (0.80%), calcium (0.44%), magnesium (0.15%), sulphur (0.45%), iron (175.2 ppm), zinc (24.43 ppm), organic carbon (9.5-17.98%), vitamins, and growth hormones that promote microbial population and plant growth.

A low-cost sustainable method called Shivansh Khad lessens reliance on damaging chemical fertilizers. It is prepared with weeds, dry plant material, wheat straw, rice husks, tree trimmings, and animal manure. Composting is an 18-day process that turns agricultural waste into rich, black compost from raw ingredients.

Organic non-edible oil cake is sometimes referred to as neem cake. Neem cake contains 5.2% N, 1.0% P, and 1.4% K. It also serves as a nematocide. It reduces nitrification by acting as a nitrogen inhibitor. Long-term availability of nitrogen is provided to the soil through it. (2012). Katyayan.

### Materials and Methods

The field experiment was conducted during summer season of 2020 at Shri Guru Ram Rai University Dehradun, Uttarakhand at the Crop Research Centre of School of Agricultural Sciences. The experiment laid out in Randomized Block Design consisting of 12 treatments T<sub>0</sub> Control, T<sub>1</sub> 100% N Through FYM+ Beejamrit inoculation, T<sub>2</sub> 100% N through Vermicompost + Inoculation, T<sub>3</sub> 100% N through Shivansh Khad + Beejamrit inoculation, T<sub>4</sub> 100% N through Neem cake+ Inoculation, T<sub>5</sub> 50% N through FYM +50% Vermicompost+ Beejamrit inoculation, T<sub>6</sub> 50% N through FYM +50% Shivansh khad+ Inoculation, T<sub>7</sub> 50% N through FYM +50% Neem cake+ Beejamrit inoculation, T<sub>8</sub> 50% N through Vermicompost + 50% Shivansh khad+ Inoculation, T<sub>9</sub> 50% N through Vermicompost + 50% Neem cake+ Beejamrit inoculation, T<sub>10</sub> 25% FYM + 25% Vermicompost + 25% Shivansh khad +25% Neem cake+ Inoculation, T<sub>11</sub> 25% FYM + 25% Vermicompost +25% Shivansh Khad + 25% Neem cake+ Beejamrit inoculation. The city is located at 25.26 degree N latitude and 81.68 degree east longitude and 410 m above sea level. The experiment site was Sandy loam in texture with Ph-6.5, organic carbon (0.42%), available N (3.52%), available Phosphorus (7.1%), available Potassium (18.1%). Between germination to harvesting several plant growth parameters

were recorded. In growth parameter plant height (cm), dry weight (g/m<sup>2</sup>) and number of branches per plant. The data recorded for different characteristics were subjected to statistical analysis by adopted the procedure of Gomez & Gomez (1984) [2].

## Result and Discussion

### Growth Parameters

#### Plant Height

At 60DAS maximum plant height (41.53cm) was recorded with T<sub>5</sub>. 50% N through FYM +50% Vermicompost+ Beejamrit inoculation which was significantly superior over all other treatments and statistically at par with treatment of T<sub>11</sub>- 25% FYM + 25% Vermicompost +25% Shivansh Khad + 25% Neem cake+ Beejamrit inoculation (39.17 cm) and T<sub>2</sub>- 100% N through Vermicompost + Inoculation (36.79 cm).

#### Number of Branches per plant

At 60 DAS maximum number of branches per plant (11.46) was recorded with application of T<sub>5</sub>. 50% N through FYM +50% Vermicompost+ Beejamrit inoculation which was significantly superior over all other treatments and statistically at par with treatment of T<sub>11</sub>- 25% FYM + 25% Vermicompost +25% Shivansh Khad +25% Neem cake+ Beejamrit inoculation (10.43) and T<sub>2</sub>- 100% N through Vermicompost + Inoculation (9.56).

#### Dry Weight

At 60 DAS, maximum dry weight (8.65gm) was recorded with application of T<sub>5</sub>. 50% N through FYM +50% Vermicompost+ Beejamrit inoculation which was significantly superior over all other treatments and statistically at par with treatment of T<sub>11</sub>- 25% FYM + 25% Vermicompost +25% Shivansh Khad +25% Neem cake+ Beejamrit inoculation (7.83g) and T<sub>2</sub>- 100% N through Vermicompost + Inoculation (6.88 g).

**Table 1:** Show the treatment of plant height, No. of branches per plant and dry weight of plant (G)

Treatments	Plant height (cm)	Number of branches per plant	Dry weight of plant (g)
Control	29.013	5.500	4.480
100% N Through FYM+ Beejamrit inoculation	35.757	8.417	6.373
100% N through Vermicompost + Inoculation	36.790	9.567	6.887
100% N through Shivansh Khad + Beejamrit inoculation	30.357	5.863	4.690
100% N through Neem cake+ Inoculation	33.157	6.620	5.963
50% N through FYM +50% Vermicompost+ Beejamrit inoculation	41.537	11.460	8.650
50% N through FYM +50% Shivansh Khad + Inoculation	31.643	6.043	5.077
50% N through FYM +50% Neemcake+ Beejamrit inoculation	32.763	6.520	5.597
50% N through Vermicompost + 50% Shivansh Khad + Inoculation	34.327	7.427	6.127
50% N through Vermicompost + 50% Neem cake+ Beejamrit Inoculation	32.043	6.253	5.470
25% FYM + 25% Vermicompost + 25% Shivansh Khad +25%Neem cake+ Inoculation	30.617	6.180	5.200
25% FYM + 25% Vermicompost +25% Shivansh Khad +25% Neem cake+ Beejamrit inoculation	39.170	10.433	7.837
C.D.	1.588	0.578	0.282
SE(m)	0.538	0.196	0.095
SE(d)	0.761	0.277	0.135
C.V.	2.746	4.510	2.740

### Conclusion

We concluded that all the treatments of different organic manures with beejamrit inoculation on growth of summer moon bean showed significant positive influence over control treatment in respect of growth parameters, treatment T<sub>5</sub>. 50% N through FYM +50% Vermicompost+ Beejamrit inoculation

T<sub>11</sub>- 25% FYM + 25% Vermicompost +25% Shivansh Khad +25% Neem cake+ Beejamrit inoculation and T<sub>2</sub>- 100% N through Vermicompost + Inoculation perform comparatively better. Therefore, from the investigation T<sub>5</sub>. 50% N through FYM +50% Vermicompost+ Beejamrit inoculation are

recommended as the best organic manure with beejamrit inoculation in summer moongbean.

### References

1. DASE. Directorate of Agriculture Statistics and Economics, Govt. of India, New Delhi; c2018.
2. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Wiley & sons; c1984.
3. Meena RS, Yadav RS. Yield and profitability of groundnut (*Arachis hypogaea* L.) as influenced by sowing dates and nutrient levels with different varieties. Legume Research. 2015;38(6):791-797.
4. Meena RS, Lal R, Yadav GS. Long term impacts of topsoil depth and amendments on soil physical and hydrological properties of an Alfisol in central Ohio, USA. Geoderma. 2020a;363:1141164 <https://doi.org/10.1016/j.geoderma.2019.114164>.
5. Yadav SK, Subhash Babu, Yadav MK, Kalyan Singh, Yadav GS, Suresh PA. review of organic farming for sustainable agriculture in Northern India. International J Agronomy; c2013. p. 1-8.
6. Meena RS, Gogoi N, Kumar S. Alarming issues on agricultural crop production and environmental stresses. Journal of Cleaner Production. 2017;142:3357-3359.
7. Anasuyamma B, Singh S, Asirinaidu B, Abhigna K. Effect of organic manures and Inorganic fertilizers on the growth and yield of Black gram (*Vigna mungo* L.). The Pharma Innovation Journal. 2022;11(4):1214-1218.
8. Khan PMI, Verma R, Dawson J, Singh M. Effect of panchagavya and organic manures on growth and yield of blackgram (*Vigna mungo* L.); c2022.
9. Yadav JK, Sharma M, Yadav RN, Yadav SK, Yadav S. Effect of different organic manures on growth and yield of chickpea (*Cicer arietinum* L.). Journal of Pharmacognosy and Phytochemistry. 2017;6(5):1857-1860.