



P-ISSN: 2349-8528
 E-ISSN: 2321-4902
www.chemijournal.com
 IJCS 2024; 12(3): 68-72
 © 2024 IJCS
 Received: 04-05-2023
 Accepted: 11-06-2024

Dr. Anupam Vibha
 Sarvoday High School,
 Darbhanga, Bihar, India

Kinetic studies of oxidation of maltose with Bi^v oxidizing agent

Anupam Vibha


Abstract

The present research work has explained the effect of concentration of maltose and effect of the concentration of Bi^v. The rates of reactions were studied at three different temperatures at 298 K, 308 K and 318 K of disaccharide sucrose. The research work has confirmed that the rate of reaction was first order with respect to the oxidant in the whole range of the concentration of bismuthate ion.

Keywords: Disaccharide, maltose, bismuth, rate of reaction, order of reaction etc

Introduction

The sugar D-(+)-maltose¹ is a glucoside consisting of two glucose monomers that are connected by an α-1,4 glycosidic bond to form maltose, 4-O-α-Dglucopyranosyl-D-glucose (C₁₂H₂₂O₁₁), of molecular weight of 342.3 g.



The most common oxidation states of bismuth are: III and V. Due to the inert pair effect, bismuth in V oxidation state is less stable and hence in this state, element has tendency to convert readily into III or elemental form. So, in this state, bismuth acts as strong oxidizing agent.

Experimental

Kinetic procedure

The requisite quantities of various components of the reaction mixture were taken in glass stoppered Erlenmeyer flasks, which were then suspended in a water bath thermostatic at the desired temperature ± 0.1 °C. The reaction was initiated by adding the requisite solution of the temperature pre-equilibrated formic acid into the reaction mixture and the time of initiation was recorded when half of the contents from the pipette were released. Aliquot samples (5 to 10 cm³) were withdrawn at different intervals of time and then quenched in an ice-cold KI (10%) solution; the liberated iodine was titrated against thiosulphate solution using starch as an indicator. Measurements in triplicate without any interference from the ingredients of the reaction mixture were in agreement to within $\pm 5\%$.

Initial rates were computed employing plane mirror method². Pseudo-first order plots were also made wherever reaction conditions permitted. Since the kinetics is not affected by the light, no further precautions were taken to exclude light from the reaction mixtures.

Corresponding Author:
Dr. Anupam Vibha
 Sarvoday High School,
 Darbhanga, Bihar, India

Kinetics of maltose with Bi^V


The kinetics of oxidation of maltose with Bi (V) in phosphoric acid medium was studied at three different temperatures say 298 K, 308 K, 318 K. The order with respect to oxidant is one. It is complex one with respect to maltose. Rate law may be expressed as:

$$\text{Rate law} = -\frac{d[\text{Bi}^{\text{V}}]}{dt}$$

$$-\frac{d[\text{Bi}^{\text{V}}]}{dt} = \frac{kK[\text{Bi}^{\text{V}}][\text{Malt}]}{1 + K[\text{Malt}]} \cdot \frac{1}{1 + K_p[\text{H}^+]}$$

$$k^l = \frac{kK[\text{Malt}]}{1 + K[\text{Malt}]} \cdot \frac{1}{1 + K_p[\text{H}^+]}$$

Where, [Bi^V] is the total gross concentration of all phosphate



The oxidation product of formaldehyde and formic acid were detected by spot test ^[4]. The results are recorded in following table.

Table 1: Stoichiometric results of oxidation of maltose Bi (V) in phosphoric acid medium (3.0M) at temp.-298 K

| [Bi ^V] $\times 10^3$ mol dm ⁻³ taken | [Malt] $\times 10^3$ mol dm ⁻³ taken | [Bi ^V] $\times 10^3$ mol dm ⁻³ (Measured) | $\frac{\Delta[\text{Bi}^{\text{V}}]}{\Delta[\text{Malt}]}$ |
|---|---|--|--|
| 1.25 | 1.0 | 1.13 | 1.13 |
| 2.25 | 1.0 | 1.08 | 1.08 |
| 2.25 | 1.5 | 1.7 | 1.133 |
| 2.25 | 1.5 | 1.61 | 1.07 |
| 3.375 | 1.5 | 1.60 | 1.06 |
| 3.375 | 2.0 | 2.31 | 1.15 |
| 3.375 | 2.0 | 2.23 | 1.15 |
| 4.5 | 2.0 | 2.23 | 1.15 |
| 5.625 | 2.0 | 2.25 | 1.124 |
| 5.625 | 2.5 | 2.81 | 1.124 |
| 4.5 | 2.5 | 2.84 | 1.136 |

Bi (V) Dependence

The concentration of Bi (V) was varied from 1.5×10^{-3} to 9.8×10^{-3} mol dm⁻³ at fixed concentration of maltose 5.0×10^{-3} mol dm⁻³. The initial rates (I_R) were computed by plane mirror method⁵. These results are recorded in following table.

Table 2: [Malt] = 5.0×10^{-3} mol dm⁻³, Temp. = 308 K [H₃PO₄] = 3.0 mol dm⁻³, Aliquot = 5 mL

| [Bi ^V] $\times 10^{-3}$ mol dm ⁻³ | [Malt] $\times 10^{-3}$ mol dm ⁻³ | [I _R] $\times 10^7$ sec ⁻¹ |
|--|--|---|
| 1.5 | 5.0 | 4.1 |
| 2.5 | 5.0 | 6.34 |
| 3.3 | 5.0 | 7.90 |
| 4.16 | 5.0 | 10.1 |
| 5.4 | 5.0 | 13.0 |
| 7.35 | 5.0 | 17.3 |
| 8.2 | 5.0 | 19.4 |
| 9.8 | 5.0 | 23.2 |

species of Bi (V) and [Malt] is the equilibrium concentration of maltose.

Stoichiometry

The stoichiometry of the reaction was determined by taking an excess of Bi (V) concentration over that of maltose (Malt) at fixed concentration of phosphoric acid (3.0M). The reaction mixtures were kept in a thermostated water bath at desired temperature and the excess of Bi (V) was determined iodometrically ^[3] after ensuring the completion of the reaction after twelve hours. The stoichiometry of the reaction was found to be 1:1 i.e., one mole of Bi (V) reacted with one mole of maltose.

The stoichiometric ratio, which is greater than this ratio account for decomposition of Bi (V), therefore more Bi (V) is consumed than required by maltose. The stoichiometry indicates the formation of these products as the end products and can be represented as follows:

The plot of initial rates (I_R) vs Bi (V) yielded straight line passing through the origin indicating first order with respect to Bi (V) oxidant.




Fig 1: A plot of IR vs [Bi^V] At Fixed concentration of maltose at 308K

Maltose Dependence

The concentration of maltose was varied in the range of 2.0×10^{-3} mol dm⁻³ to 8.0×10^{-3} mol dm⁻³ at fixed concentration of Bi (V) i.e. 1.63×10^{-3} mol dm⁻³ and also at fixed concentration of phosphoric acid 3.0 mol dm⁻³ at 298 K. The results are in following tables.

Table 3: [Bi^V] = 1.63×10^{-3} mol dm⁻³, Temp. = 298 K [H₃PO₄] = 3.0 mol dm⁻³, Aliquot = 5 mL

| [Malt] $\times 10^3$ mol dm ⁻³ | [Bi ^V] $\times 10^{-3}$ mol dm ⁻³ | [I _R] $\times 10^7$ Sec ⁻¹ | K ₀ $\times 10^4$ Sec ⁻¹ |
|---|--|---|--|
| 2.0 | 1.63 | 1.17 | 4.01 |
| 3.0 | 1.63 | 1.60 | 6.34 |
| 4.0 | 1.63 | 1.84 | 7.9 |
| 5.0 | 1.63 | 2.02 | 10.2 |
| 6.0 | 1.63 | 2.13 | 12.9 |
| 8.0 | 1.63 | 2.18 | 17.4 |

The concentration of maltose was varied in the range of $1.0 \times 10^3 \text{ mol dm}^{-3}$ to $7.0 \times 10^3 \text{ mol dm}^{-3}$ at fixed concentration

of Bi (V) i.e. $1.63 \times 10^{-3} \text{ mol dm}^{-3}$ and also at 308 K. The results are in following tables.

Table 4: $[\text{Bi}^{\text{V}}] = 1.63 \times 10^{-3} \text{ mol dm}^{-3}$, Temp.= 308 K $[\text{H}_3\text{PO}_4] = 3.0 \text{ mol dm}^{-3}$, Aliquot= 5mL

| $[\text{Malt}] \times 10^3 \text{ mol dm}^{-3}$ | $[\text{Bi}^{\text{V}}] \times 10^{-3} \text{ mol dm}^{-3}$ | $[\text{I}_R] \times 10^7 \text{ mol dm}^{-3} \text{Sec}^{-1}$ | $\text{K}_0 \times 10^4 \text{ Sec}^{-1}$ |
|---|---|--|---|
| 1.0 | 1.63 | 1.68 | 1.68 |
| 2.0 | 1.63 | 2.68 | 1.34 |
| 3.0 | 1.63 | 3.34 | 1.12 |
| 4.0 | 1.63 | 4.33 | 1.08 |
| 5.0 | 1.63 | 5.02 | 1.01 |
| 6.0 | 1.63 | 5.52 | 0.92 |
| 7.0 | 1.63 | 6.02 | 0.88 |

The concentration of maltose was varied in the range of $2.0 \times 10^3 \text{ mol dm}^{-3}$ to $20.0 \times 10^3 \text{ mol dm}^{-3}$ at fixed concentration

of Bi (V) i.e. $1.63 \times 10^{-3} \text{ mol dm}^{-3}$ and also at 318 K. The results are in following tables.

Table 5: $[\text{Bi}^{\text{V}}] = 1.63 \times 10^{-3} \text{ mol dm}^{-3}$, Temp.= 318 K $[\text{H}_3\text{PO}_4] = 3.0 \text{ mol dm}^{-3}$, Aliquot= 5mL

| $[\text{Malt}] \times 10^3 \text{ mol dm}^{-3}$ | $[\text{Bi}^{\text{V}}] \times 10^{-3} \text{ mol dm}^{-3}$ | $[\text{I}_R] \times 10^7 \text{ Sec}^{-1}$ | $\text{K}_0 \times 10^4 \text{ Sec}^{-1}$ |
|---|---|---|---|
| 2.0 | 1.63 | 4.67 | 2.33 |
| 4.0 | 1.63 | 10.68 | 2.61 |
| 6.0 | 1.63 | 20.13 | 3.31 |
| 10.0 | 1.63 | 24.01 | 2.42 |
| 15.0 | 1.63 | 32.02 | 2.11 |
| 20.0 | 1.63 | 40.13 | 2.02 |

The initial rates were calculated by plane mirror method [5]. The plot of initial rate (I_R) vs concentration of maltose yielded a curve line. Thus, the order with respect to sugar seemed to be complex one.

Ionic strength dependence

Investigations were carried out at different ionic strength in the range of 1.4 mol dm^{-3} to 2.6 mol dm^{-3} . The reaction rates increase with increasing ionic strength. The following table shows the results.

Table 6: Ionic Strength Dependence $[\text{Bi}^{\text{V}}] = 1.6 \times 10^{-3} \text{ mol dm}^{-3}$, Temp.= 308 K $[\text{Malt}] = 5.0 \times 10^{-3} \text{ mol dm}^{-3}$, Aliquot=5mL $\text{H}_3\text{PO}_4 = 3.0 \text{ mol dm}^{-3}$

| 1 mol dm^{-3} | $[\text{I}_R] \times 10^7 \text{ Sec}^{-1}$ |
|-------------------------|---|
| 1.4 | 5.36 |
| 1.6 | 5.76 |
| 1.8 | 6.01 |
| 2.0 | 6.41 |
| 2.4 | 7.02 |
| 2.6 | 7.36 |

H^+ ION Dependence

The H^+ ion concentration was varied from 0.5 to 2.0 mol dm^{-3}

by adding $[\text{HClO}_4]$ at constant ionic strength. The reaction rate decreases with increasing H^+ ion concentration.


Table 7: H^+ Ion Dependence, $[\text{Bi}^{\text{V}}] = 1.6 \times 10^{-3} \text{ mol dm}^{-3}$, Temp.= 308 K $[\text{Malt}] = 5.0 \times 10^{-3} \text{ mol dm}^{-3}$, Aliquot=5mL $\text{H}_3\text{PO}_4 = 3.0 \text{ mol dm}^{-3}$

| $[\text{HClO}_4] 1 \text{ mol dm}^{-3}$ | $[\text{I}_R] \times 10^7 \text{ Sec}^{-1}$ |
|---|---|
| 0.50 | 9.22 |
| 0.90 | 7.52 |
| 1.00 | 7.13 |
| 1.25 | 6.02 |
| 1.50 | 5.33 |
| 1.75 | 5.02 |
| 2.00 | 4.85 |

Results and Discussion

The oxidation of maltose by sodium bismuthate was carried out, which gave two equivalents of formaldehyde and ten

equivalents of formic acid in acidic medium ($\text{pH} = 3$). The titration curves suggested the formation of following oxidative products as:



Effect of Bismuthate

The plot of initial rates (I_R) vs Bi (V) yielded straight line passing through the origin indicating first order with respect to Bi (V) oxidant as observed in table:-2. It was observed that the rate law equation with respect to bismuthate ion may be written as:

$$\text{Rate} \propto [\text{Bismuthate}] \quad (1)$$

Effect of Concentration of Maltose

The initial rates were calculated by plane mirror method¹. The plot of initial rate (I_R) vs concentration of maltose was yielded a curve line. Thus, the order with respect to maltose seemed to be complex one.

The concentration of maltose was varied in the range of 1.0×10^3 mol dm⁻³ to 20.0×10^3 mol dm⁻³ at fixed concentration of Bi (V) i.e. 1.63×10^{-3} mol dm⁻³ and also at fixed concentration of phosphoric acid 3.0 mol dm⁻³. The investigations were carried out at three different temperatures i.e. 298 K, 308 K, 318 K. The results are shown in Tables:-3, 4 and 5 and their respective graphs.

The graph at 298 K shows that at lower concentration of the maltose, rate linearly varies with the concentration of the sugar exhibiting first order say about 4.0×10^3 mol dm⁻³ concentration of the maltose. However, as concentration exceeds this value, rate became independent or more clearly it remains constant.

The graph at 308 K remains almost same as at 298 K. it shows that at lower concentration of the maltose say about 6.0×10^3 mol dm⁻³ concentration of the maltose, rate linearly varies with the concentration of the maltose exhibiting first order. However, as concentration exceeds this value, rate became constant.

The graph at 318 K was found to be more complex. The plot of initial rate (I_R) vs concentration of maltose was yielded a curve lines and different from the graphs obtained at 308 K and 318 K. Thus, the order with respect to maltose seemed to be complex one.

Hence, the rate law equation with respect to maltose is difficult to express. The overall rate may be pseudo first order say first order with respect to the [bismuthate] and remains independent from [Maltose].

Effect of Ionic Strength

Investigations were showed reaction rates increases with increasing ionic strength shown in table 6.


The variation of rate of reaction with ionic strength indicates that formation of ionic intermediate, which is stabilized by ionic medium, during the course of reaction. Since ionic intermediate is stabilized by ionic medium, which decreases the activation energy. Hence rate of reaction increases with increase of ionic strength.

Effect of $[H^+]$

The concentration of H^+ ion was varied from 0.5 to 2.0 mol dm⁻³ by adding $[HClO_4]$ at constant ionic strength. The reaction rate decreases with increasing H^+ ion concentration. The observation at 308 K has been given in table 7. It is obvious that the rate of reaction decreases with increasing H^+ ion concentration i.e rate decreases with decreasing pH.

Reaction Mechanism

Considering all these observations and also accounting for the experimental findings a probable reaction mechanism can be proposed as follows:



By using the Eq.1, Eq.2 and Eq.3, the rate law equation may be derived as:

$$-\frac{d[Bi^V]}{dt} = \frac{kK[Bi^V][Malt]}{1+K[Malt]} \cdot \frac{1}{1+K_P[H^+]} \quad (4)$$

$$\text{or, } \frac{-\frac{d[Bi^V]}{dt}}{[Bi^V]} = \frac{kK[Malt]}{1+K[Malt]} \cdot \frac{1}{1+K_P[H^+]}$$

$$\text{or, } K^I = \frac{kK[Malt]}{1+K[Malt]} \cdot \frac{1}{1+K_P[H^+]} \quad (5)$$

Where, K^I = observed first order rate constant,
 $[Malt]$ = Equilibrium concentration of maltose
 $[Bi^V]$

= total gross concn. of all phosphato species of Bi^V

$$\frac{1}{K^I} = \left\{ \frac{1+K[Malt]}{kK[Malt]} \right\} \cdot \{1 + K_P[H^+]\} \quad (6)$$

$$\text{or, } \frac{1}{K^I} = \left\{ \frac{1}{kK[Malt]} + \frac{1}{k} \right\} \cdot \{1 + K_P[H^+]\}$$

$$\text{or, } \frac{1}{K^I} = \left\{ \frac{1+K_P[H^+]}{kK[Malt]} + \frac{1+K_P[H^+]}{k} \right\} \quad (7)$$

A plot of $1/K^I$ vs $1/[maltose]$ at constant $[H^+]$ ion concentration was constructed from the equation. A straight line with non-zero intercept was obtained as below:

Taking different concentrations of maltose at three different temperatures 298 K, 308 K and 318 K, the value of 'k' was calculated to be ~95, ~65 and ~20 at 298 K, 308 K and 318 K, respectively.

A plot of $1/[k^I]$ vs $1/[maltose]$ at different concentration of $[H^+]$ ion yield a straight line as below:


The plot of intercept $\frac{1+K_P[H^+]}{k}$ and gradient $\frac{1+K_P[H^+]}{kK}$ against $[H^+]$ were further constructed.

The value of k and K_P were calculated from the ratio of intercept and gradient as $4.3 \text{ dm}^3 \text{ mol}^{-1}$ and $1.4 \text{ dm}^3 \text{ mol}^{-1}$.

Further, K^I was calculated from the values of k, K and K_P by the help of respective equations.

The mode of electron transfer is through an intermediate to be formed in two equivalent steps taking oxidative cleavage.

The oxidation of maltose with sodium bismuthate in phosphoric acid medium takes place via the formation of bismuthic acid followed by the formation of cyclic intermediate, which readily decomposes to form carbonyl compounds. It may be shown as:



Oxidation kinetics of maltose has been studied by some researchers [6, 7] and it has been observed that oxidation of sugar by sodium busmuthate takes place almost similar to that of lead tetraacetate [8-10]. Such oxidative cleavage had been also observed [11]. It is remarkable that the side reactions are also observed [12, 13]. The oxidation products of maltose have also been found as formaldehyde and formic acid by other workers [14].

Conclusion

Although, it was observed that rate of reaction was first order with respect to concentration of the maltose only upto the lower range of the concentration of maltose. However, at higher concentration of maltose, the rate reaction became independent of the concentration. Thus, the rate of reaction was observed as complex with respect to the concentration of the maltose.

Further, the effect of ionic strength on the rate of reaction was also studied. It was found that rate of reaction increases with increase of ionic strength, which indicates that the formation of ionic intermediate. As ionic intermediate is stabilized by the ionic medium so, activation energy decreases. It increases the rate of reaction.

Furthermore, the effect of acidic medium i.e. the concentration of H^+ ion (i.e pH of the reaction mixture) on the rate of reaction was also studied. It was found that rate of reaction decreases with increase of pH of the reaction mixture.

References

1. Preedy VR. Dietary sugars: Chemistry, analysis, function and effects. Royal Society of Chemistry; c2012.
2. Latshaw M. Journal of the American Chemical Society. 1925;47:793.
3. Cseke LJ, Kirakosyan A, Kaufman PB, Warber S, Duke JA, Brielmann HL. Natural Products from Plants. CRC Press; c2016.
4. Rosa M, Prado C, Podazza G, Interdonato R, González JA, Hilal M, *et al.* Soluble sugars. Plant Signaling & Behavior. 2009;4(5):388-393.
5. Hu FB. Resolved. Obesity Reviews. 2013;14(7):606-19.
6. Burstein GT, Wright GA. Nature. 1969;221:169.
7. Cardillo G, Orena M, Sandri S. Synthesis of aldehydes and ketones from allylic and benzylic halides. Tetrahedron Letters. 2009;50(34):3985-3989.
8. Angyal SJ, James K. Oxidation of carbohydrate acetals: a new synthesis of ketoses. Australian Journal of Chemistry. 2005;24(6):1219.
9. Angyal SJ, Evans ME. Oxidation of carbohydrates with LTA. Australian Journal of Chemistry. 2004;25(8):1513.
10. Ramakrishanan PS. Kinetics and mechanism of sugar. Asian Journal of Chemistry. 2000;12(4):1096.
11. Trutneva AA, Nurgatin VV, Shernin GP. Oxidation of 2,2,4,4',6,6'-hexanitro-3-methyldiphenyl sulphide by chromic acid under acidic conditions. Transactions of Kazan Chemical-Technological Institute. 2003;50:76.
12. Meenakshisundaram S, Amutha M. Studies on the oxidation of some sugar. Journal of Chemical Research (S). 1999;2.
13. Banerji KK. Kinetics of the oxidation of diol. Journal of the Chemical Society, Perkin Transactions 2. 2000;2(12):2065. Erdal G, Esengün K, Erdal H, Gündüz O: Energy use and economical analysis of sugar beet production in Tokat province of Turkey. Energy 2007, 32: 35-41.