



ISSN: 2321-4902

Volume 1 Issue 4

Online Available at [www.chemijournal.com](http://www.chemijournal.com)

## International Journal of Chemical Studies

### Thermokinetic Parameter of Co(II), Ni(II), Cu(II) and Zn(II) of Schiff base Complexes

S. R. Kelode

1. Department of Engg. Chemistry, Jagadambha College of Engineering and Technology Yavatmal  
[Email: [sandeep\\_kelode@yahoo.co.in](mailto:sandeep_kelode@yahoo.co.in) Mob No. 9371804248]

Metal complexes Co(II), Ni(II), Cu(II) and Zn(II) of newly synthesized Schiff base 2-hydroxy-5-chloro acetophenone 2-imino-4-phenyl thiazole. It should be prepared from 2-hydroxy-5-chloro acetophenone and 2-amino-4-phenyl thiazole have been synthesized and characterized on the basis of elemental analysis, Infrared, <sup>1</sup>H NMR, molar conductance and magnetic susceptibilities analysis. The Schiff base acts as a monobasic bidentate ligand commonly coordinates through the oxygen atom of phenolic OH group and the nitrogen atom of azomethine group, which is confirmed by IR spectral data. All the metal complexes have studies thermal properties and their thermokinetic parameter.

*Keyword:* Schiff base, Magnetic, Thermal studies

#### 1. Introduction

The condensation of primary amines with aldehydes and ketones give imines. Imines that contain an aryl group bound to the nitrogen or to the carbon atom are called Schiff bases. The Schiff bases are widely used ligands due to their facile synthesis, significant versatility and good solubility in common solvents. Schiff bases derived from aldehydes or ketones with hydrazones and aroyl hydrazines have been widely used as ligand for the synthesis of transition metal complexes<sup>[1]</sup>. Schiff bases or their metal complexes have many applications in different fields<sup>[2]</sup> Hydrazones, Heteroaroyl hydrazones ligands and their metal complexes are biologically active. It should be heteroaroyl Schiff base forms stable metal complexes with transition metal ions and inner transition metal ions due to complexing ability of ligand through keto-enol tautomerism and availability of other donor sites in the ligand i.e. isonicotinoyl

hydrazide is one of the drug in chemotherapy of tuberculosis<sup>[3]</sup>. Due to its biological potency, pharmacological properties and synthetic flexibility of Schiff base derived from isonicotinic acid hydrazide<sup>[4,5]</sup>.

The aim of the present research paper to synthesize various transition metal complexes of Schiff base condensed from 2-hydroxy-5-chloroacetophenone and 2-amino-4-phenyl thiazole.

#### 2. Materials and Methods


All the chemicals were of A. R. grade and used as received. 2-hydroxy-5-chloro- acetophenone (HCA) and 2-amino-4-phenylthiazole was prepared by known methods<sup>[6-9]</sup>. The solvents were purified by standard methods<sup>[10]</sup>.

##### 2.1. Synthesis of 2-amino-4-phenylthiazole

The synthesis of 2-amino-4-phenylthiazole prepared by known method<sup>[7-9]</sup>. The product was

filtered and crystallized from 70% ethanol, after several minutes the golden coloured product of 2-

amino-4-phenylthiazole was separated out. Yield: 75%; m.p.: 148–150 °C.




Scheme 1. Synthesis of 2-amino-4-phenylthiazole

## 2.2 Synthesis of 2-hydroxy-5-chloro acetophenone 2-imino-4-phenyl thiazole [HCAIPT]

A solution of 2-hydroxy-5-chloro acetophenone (0.02 M) in 25ml of ethanol was added to an ethanolic solution (25 ml) of 2-amino-4-phenylthiazole(0.02 M) and the reaction mixture was refluxed on a water bath for 4 h. After

cooling a pale yellow coloured crystalline solid was separated out. It was filtered and washed with ethanol, crystallized from DMF and dried under reduced pressure at ambient temperature. The purity of ligand was checked by elemental analysis and m.p. It was also characterized by IR and <sup>1</sup>H NMR spectral studies.

Yield: 35%; m.p. 290 °C.



Scheme 2. Synthesis of HCAIPT

**Table 1:** Analytical data of the Ligands.

Sr. No	Ligand	Molecular Formula	Formula Weight	Color and nature	Elemental Analysis				
					C% found (Cal.)	H% Found (Cal.)	N% Found (Cal.)	Cl% Found (Cal.)	S% Found (Cal.)
1.	HCAIPT	C <sub>17</sub> H <sub>12</sub> N <sub>2</sub> OSCl	327.6	Yellow Crystalline	61.38 (62.27)	03.10 (03.66)	08.24 (08.54)	10.21 (10.83)	09.13 (09.79)

## 2.3 Preparation of complexes

All the metal complexes were prepared in a similar way by following method. To a hot solution of ligand HCAIPT (0.02 M) in 25 ml of ethanol, a suspension of respective metal salts was added drop wise with constant stirring. The reaction mixture was refluxed on a water bath for 4-6 h. The precipitated complexes were filtered, washed with ethanol followed by ether and dried over fused calcium chloride. Yield: 45-50% The complexes are soluble in DMSO and DMF but insoluble in water and common organic solvents. The metal chloride content of complexes were analyzed (Table 2) by standard methods<sup>[11]</sup>. The <sup>1</sup>H NMR spectra of ligand was obtained from

RSIC Chandigarh. IR spectra of the compounds were recorded on Perkin Elmer 842 spectrophotometer in the region 400-4000 cm<sup>-1</sup>, Carbon, Hydrogen and Nitrogen analysis were carried out at RSIC, Punjab University, Chandigarh. The molar conductance of the complexes at 10<sup>-3</sup> M dilution in DMF were determined using equiptronic digital conductivity meter EQ-660 with a cell constant 1.00 cm<sup>-1</sup> at room temperature. The magnetic moment measurement were made on a Gouy balance at room temperature using [HgCo(SCN)<sub>4</sub>] as the calibrant. The molecular weights of the complexes were determined by Rast method. The thermogravimetric analysis were performed on

laboratory set up apparatus in air atmosphere at 10 °C min<sup>-1</sup> heating rate. The molecular weights of the complexes were determined by Rast method.

**Table 2:** Analytical data and molar conductance of the compounds

Compounds	Colour	Mol.wt.	Analysis % Found (calc.)					$\mu_{\text{eff}}$ B.M.	$\Lambda_M$ ( $\Omega^{-1}$ cm <sup>2</sup> mol <sup>-1</sup> )	
			M	C	H	N	Cl			S
[Co(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> H <sub>2</sub> O	Brown	783.1	7.22 (7.52)	51.86 (52.10)	3.35 (3.57)	6.96 (7.15)	8.80 (9.06)	7.84 (8.20)	4.38	5.9
[Ni(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> H <sub>2</sub> O	Green	782.9	7.20 (7.39)	51.78 (52.11)	3.38 (3.58)	6.86 (7.15)	8.90 (9.06)	7.79 (8.19)	3.1	8.4
[Cu(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> H <sub>2</sub> O	Brown	787.7	7.20 (7.98)	51.60 (51.79)	3.15 (3.55)	6.79 (7.11)	8.82 (9.01)	7.95 (8.15)	1.90	7.2
[Zn(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> H <sub>2</sub> O	Reddish	807.6	7.80 (7.91)	50.11 (50.52)	3.48 (3.71)	6.12 (6.93)	8.30 (8.79)	7.54 (7.94)	Dia	8.4

### 3. Result and Discussion

The Schiff base ligand HCAIPT and its complexes have been characterized on the basis of <sup>1</sup>H NMR, IR spectral data, elemental analysis, molar conductance, magnetic susceptibility measurements and thermogravimetric analysis data. All these values and analytical data is consistent with proposed molecular formula of ligand. All the compounds are coloured solid and stable in air. They are insoluble in water but soluble in coordinating solvents like DMF and DMSO. The molar conductance values in

DMF(10<sup>-3</sup>M) solution at room temperature (Table2) shows all the complexes are non electrolytes.

The <sup>1</sup>H NMR spectra of ligand HCAIPT shows signals at  $\delta$  11.26, (1H, s phenolic OH), 7.41, 7.40, 7.39 and 7.38 (4H, m, phenyl)  $\delta$  6.51, 6.50, and 6.58 (3H, s Phenyl), 6.62 (1H s thiophene), and 2.16(3H, s, methyl)<sup>[12-15]</sup>. IR spectra of ligand and metal complexes summarized in table 3. As per observation  $\nu(\text{C}=\text{N})$  peaks at 1618cm<sup>-1</sup> and absence of C=O peak at around 1700–1730 cm<sup>-1</sup> indicates the Schiff base formation<sup>[16-19]</sup>.

**Table 3:** IR spectra of ligand and metal complexes.

Compound	$\nu(\text{O-H})$ hydrogen bonded	$\nu(\text{C}=\text{N})$ Imine	$\nu(\text{C-O})$ phenolic	$\nu(\text{C-S})$	$\nu(\text{M-O})$	$\nu(\text{M-N})$
HCAIPT	3109	1618	1514	1122	-	-
[Co(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> H <sub>2</sub> O	-	1606	1504	1098	470	430
[Ni(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> H <sub>2</sub> O	-	1580	1465	1090	468	422
[Cu(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> H <sub>2</sub> O	-	1602	1504	1110	509	410
[Zn(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> H <sub>2</sub> O	-	1503	1448	1106	470	425

#### 3.1 Thermogravimetric Parameter

Thermogravimetric<sup>[18]</sup> study indicates all the complexes are stable up to 60-70 °C. All the complexes shows half decomposition temperature (Table 4). The Thermal activation energy, Frequency factor, Entropy change and Free Energy change was calculated by Freeman-Carroll<sup>[19]</sup>, Horowitz-metzger<sup>[20]</sup> and Broido<sup>[21]</sup> method.

#### 4. Conclusions

Studies of magnetic moment and electronic spectral data shows characterization and structural changes in metal complexes. Thermal studies concluded conformed structure of metal complexes from the study of water loss in metal complexes and calculate other parameter like frequency factor, entropy change, free energy change and half decomposition temperature.

**Table 4:** Thermal decomposition data of HCAIPT and its complexes.

Compounds	Half Decomposition Temp.(°C)	Activation Energy (kJ mole <sup>-1</sup> )			Frequency Factor Z (sec <sup>-1</sup> )	Entropy Change -ΔS (J mol <sup>-1</sup> K <sup>-1</sup> )	Free Energy Change ΔF (kJ mol <sup>-1</sup> )
		B*	H-M**	F-C***			
HCAIPT	267.51	3.28	4.41	4.53	87.20	210.75	115.66
[Co(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] H <sub>2</sub> O	404.21	5.28	8.61	6.84	138.87	210.62	148.51
[Ni(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] H <sub>2</sub> O	386.11	6.76	8.35	6.42	134.47	210.56	146.42
[Cu(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] H <sub>2</sub> O	423.31	6.54	8.46	7.41	150.68	210.49	155.20
[Zn(L) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> ] 2H <sub>2</sub> O	710.41	11.21	18.52	11.22	222.32	209.36	217.59

\* Broido, \*\* Horowitz-Metzger and \*\*\* Freeman-Carroll

## 5. References

- Jayaramudu M, Reddy KH. Indian J. Chem.1999; 38A:1173.
- Ramesh M, Chandrashekar KB, Hussain RK. Indian J Chem 2000; 39A:1337.
- Sharma RP, Kothari AK, Sharma NK. Indian J Derm Vener Lepr 1995; 61:26.
- Jahagirdar JA, Patil BG, Havinale BR. Indian J Chem1990; 29A:924.
- Hawlder MBH, Begum MS. Indian J Chem 2004; 43A:2352.
- Ballhausen CJ, Gray HB. Inorg Chem 1962; 1:11.
- Furniss BS, Hannaford AJ, Smith PWG, Tatchell AR. Vogel's practical organic chem. Edn 5, Logman Scientific Technical, John Wiley and Sons, 1989.
- Vogel AI. A Text book of quantitative inorganic chemistry. Edn 3, ELBS London, 1961.
- Maurya MR, Gopinathan C. Indian J Chem 1996; 35A:701.
- Jayaramadu M, Reddy KH. Indian J Chem, 1999; 38A:1173.
- Sastry PS, Rao TR. Proc. Indian Acad Sci 1995; 107(1):25.
- Mahajan RK, Patial VP. J Indian Council Chem 2001; 18(1):4.
- Maurya MR, Antony DC, Gopinathan, S, Gopinathan C, Bull Chem Soc Jpn 1995; 68:554.
- Viroopakshapp J, Vithalrao D, J Indian Chem Soc 1996; 73:531.
- Figgis BN. "Introduction to ligand field", weley Estern Ltd 1976.
- Sacconi L. J Ame Chem Soc 1954; 76:3400.
- Panda AK, Dash DC, Mishra P, Mohnaty H. J Indian Chem Soc 1996; 35A:324.
- Jürgen S, Frank S, Lentz A, Silvia R, J Chem Soc Perkin Trans 2 1998; 75.
- Mallikarjun K. E. J Chem 2004; 1(2):105.
- Horowitz H, Metzger G Anal Chem 1963; 35:1464.
- Broido A. J Polym Sci 1964; A2:1761.