



P-ISSN 2349-8528
 E-ISSN 2321-4902
 IJCS 2015; 3(2): 83-87
 © 2015 JEZS
 Received: 20-06-2015
 Accepted: 21-07-2015

Shubha Ratna Shakya
 Department of Zoology, Amrit
 Science Campus, Tribhuvan
 University Kathmandu, Nepal.

Medicinal uses of ginger (*Zingiber officinale* Roscoe) improves growth and enhances immunity in aquaculture

Shubha Ratna Shakya

Abstract

The medicinal plants are of great use in pharmaceutical, cosmetic, agricultural and food industry. The efficacy of some herbal products is beyond doubt, the most recent examples being *Silybum marianum* (Linn.) Gaertn (Silymarin), *Artemisia annua* Linn. (Artemisinin) and *Taxus baccata* Linn. (taxol). Randomized, controlled trials have proved the efficacy of some established remedies, for instance *Zingiber officinale* Roscoe, commonly known as ginger. Ginger contains natural organic materials beneficial to health and enhances resistance to infectious diseases by increasing non-specific and specific immune mechanisms. The rhizome of ginger has shown to be effective in the control of a range of bacterial, viral, fungal and parasitic diseases in humans, poultry and aquaculture owing to its antimicrobial, antioxidant, growth promoter and as immunostimulant properties to health. Hence, this review focuses on the use of ginger as growth promoter, antimicrobial agent, and antioxidant and as immunostimulant in aquaculture.

Keywords: Ginger, *Zingiber officinale*, gingerols, antioxidant, aquaculture.

1. Introduction

The world trend to improve food security and to use natural products will drive the chemically synthesized antibiotics and growth promoters out of use. Aquaculture is therefore an emerging industrial sector which requires continued research with scientific technical development and innovations (Ibrahim *et al.*, 2010) [24]. Extensive use of antibiotics in aquaculture leads to the emergence of antibiotic-resistant bacteria and generation of toxicants, which may cause risks to the environment (Esiobu *et al.*, 2002) [13], and immunosuppression in the host (Panigrahi & Azad, 2007) [38]. There are a large number of feed additives available to improve fish growth performance. Some of these additives used in feed mill are chemical products, especially hormones and antibiotics, which may cause unfavorable side effects. To alleviate these problems, increasing attention is being given to the use of natural alternative feed additives such as ginger for disease-control strategies in aquaculture. Ginger enhances resistance to infectious disease by increasing non-specific and specific immune mechanisms (Harikrishnan *et al.*, 2011) [21]. Ginger contains natural organic materials that facilitate growth, anti-stress, environmentally friendly and antimicrobial properties in fish (Maqsood *et al.*, 2011) [33].

Ginger as a natural antibiotic is the earliest known medicinal plant. It has shown to be effective in treating diseases in humans, poultry and aquaculture owing to its antimicrobial, antioxidant, growth promoter and immunostimulant properties. An optimized dose of ginger is recommended in the diet. Ginger (*Zingiber officinale* (L.) Roscoe) has been used as a spice for over 2000 years (Bartley & Jacobs, 2000) [6]. It is also called "The Great Medicament" in Ayurvedic medicines (Tan & Vanitha, 2004) [44] and is generally considered as a safe herbal medicine (Weidner and Sigwart, 2000) [46].

Ginger (*Zingiber officinale* Roscoe) is a creeping perennial underground rhizome belonging to family Zingiberaceae (Sharma, *et al.*, 2010) [41]. Nepal is the third biggest producer of ginger in the world (FAO, 2012). In the first year, a green, erect reed like stem about 60 cm high grows from this rhizome. The plant has narrow; lanceolate to linear-lanceolate, 15-30 cm long leaves which die of each year. The odour and taste are characteristic, aromatic and pungent. Ginger valued as a spice has been used through ages in almost all systems of medicine against many maladies. The plant is indigenous to Southeast Asia and is cultivated in a number of countries including Nepal. The smell and taste of the drug are typical and aromatic. The medicinal part

Correspondence:
Shubha Ratna Shakya
 Department of Zoology, Amrit
 Science Campus, Tribhuvan
 University Kathmandu, Nepal.

of the herb is dried rhizomes. It is now recognized as a drug of choice for nausea and vomiting. It has also been found useful in pregnancy related morning sickness. In rheumatoid arthritis and osteoarthritis it is used as a natural pain reliever and an anti-inflammatory agent. It is also useful in curing ulcer and preventing heart attack and stroke. A number of active constituents and medicinal properties have been reported during the last decade. Thus, the present article provides a comprehensive account of important medicinal properties of this versatile herb.

1.1. Nutrient Composition

Fresh ginger contains 80.9% moisture, 2.3% protein, 0.9% fat, 1.2% minerals, 2.4% fibre and 12.3% carbohydrates. The minerals present in ginger are iron, calcium and phosphorous. It also contains vitamins such as thiamine, riboflavin, niacin and vitamin C. The composition varies with the type, variety, agronomic conditions, curing methods, drying and storage conditions (Govindarajan, 1982; Gugnani & Ezenwanze, 1985) [15, 17].

1.2. Chemical composition

Ginger is antiplatelet, antibacterial, antifungal, antiviral, antiworm, anti-inflammatory and has anti-oxidative activity. It affects gastrointestinal and cardiovascular systems, antilipidemic and antihyperglycemic, anti-tumour properties and are known to be effective as an immuno-modulatory agent in human and animals, including fish (Nya and Austin, 2009; Apines-Amar *et al.*, 2012 and Talpure *et al.*, 2013) [37, 3, 43]. Gingerols are the major active components in the fresh rhizome (Hoffman, 2007). The volatile oil components consists mainly of sesquiterpene hydrocarbons, predominantly zingiberene (35%), curcumene (18%) and farnesene (10%) (Govindarajan, 1982) [15]. Non-volatile pungent compounds include gingerols, shogaols, paradols and zingerone. Paradol is similar to gingerol and is formed from hydrogenation of shogoal (phenylalkanones).

Ginger contains fats, waxes, carbohydrates, vitamins and minerals. Ginger rhizomes also contain a potent proteolytic enzyme called zingibain. The pungent taste of ginger is due to nonvolatile phenylpropanoid-derived compounds, particularly gingerols and shogaols. Supplementing ginger in fish diets may enhance disease resistance by reinforcing host innate immune functions that are necessary for protection against infectious diseases. Ginger may play diverse biological roles in anti-oxidative, anti-inflammatory, hypolipidemic, anti-carcinogenic, anti-nausea, antithrombotic, cardiovascular, and antibacterial processes (Kikuzaki & Nakatan, 1993; Grzannar *et al.*, 2005; Kadnur & Goyal, 2005; Stoilova *et al.*, 2007; Nicoll & Henein 2009) [30, 16, 26, 42, 35].

1.3. Ginger as growth promoter

Ginger extracts have been reported to enhance the growth of teleosts. For instance, rainbow trouts (*Oncorhynchus mykiss*) that were fed ginger had significant increases in growth, feed conversion, and protein efficiency. Supplementing diets with acetone extract from ginger was reported to enhance the growth of tilapia (*Oreochromis mossambicus*) (Immanuel *et al.*, 2009) [25]. The administration of ginger can produce significantly higher weight gain and specific growth rates in *Penaeus monodon* post larvae (Venkataramalingam *et al.*, 2007) [45]. The efficiency of feed proportionately increased with the increased percentage of ginger. Moreover, digestive enzyme activity significantly increased with ginger enrichment (Venkataramalingam *et al.*, 2007) [45].

1.4. Ginger as antioxidant agent

Ginger is a strong antioxidant substance and may either mitigate or prevent generation of free radicals (Haksar *et al.* 2006; Kim *et al.* 2007) [20, 31]. It is considered to be a safe herbal medicine with only a few insignificant side effects (Ali *et al.*, 2007) [2]. All major active ingredients of Ginger, such as zingerone, gingerdiol, zingiberene, gingerols and shogaols, are known to possess anti-oxidant activities (Chrubasik *et al.*, 2005) [9]. This antioxidant activity in ginger is due to the presence of polyphenol compounds (6-gingerol and its derivatives) (Chen, *et al.*, 1986; Herrmann, 1994) [8]. The total phenolic content in the alcoholic extract of the dried rhizome of ginger is 870.1 mg/g dry extract (Stoilova *et al.*, 2007) [42]. Antioxidant property of ginger is extremely significant as it can be used as a preventive agent against a number of diseases in aquaculture.

1.5. Ginger as antimicrobial agent

Ginger has some antifungal properties as well (Nielsen & Rios; 2000) [36]. Ginger inhibits *Aspergillus* sp, a fungus known for the production of aflatoxin, a carcinogen (Nielsen & Rios, 2000) [36]. Fresh ginger juice showed inhibitory action against *Aspergillus niger*, *Saccharomyces cerevisiae*, *Mycoderma* sp. And *Lactobacillus acidophilus* (Kapoor, 1999) [27]. It provides protection against invading microorganisms, including bacteria such as *E. coli* and *Staphylococcus aureus* (a common cause of skin infections) and fungi, including *Candida albicans*. The benzene extract of *Zingiber officinale* rhizome showed highest antibacterial activity against drug resistant *P. aeruginosa* isolated from wound and pus samples. Melvin *et al.*, 2009 also reported that the ginger extract exhibited maximum antimicrobial activity against *P. aeruginosa*. Ginger was also found to be protective against DNA damage induced by H₂O₂ and enhanced health (Khaki, *et al.*, 2009; Robertsen, 1999) [29, 39]. The rhizome of ginger (*Zingiber officinale*) has been reported to possess a broad-spectrum of prophylactic and therapeutic activities (Ernst and Pittler, 2000) [12].

Ginger is effective in the control of a range of bacterial, viral, fungal and parasitic diseases (Agrawal *et al.*, 2001; Martins *et al.*, 2001; Endo *et al.*, 1990) [1, 34, 11]. Cultured fish suffer from a wide variety of bacterial, viral, parasitic and fungal diseases (Austin & Austin, 2007) [4]. The application of ginger in aquaculture is an innovative approach to enhance health of fish and to prevent diseases. Numerous evidences suggest that many dietary factors may be used alone or in combination with traditional chemo- therapeutic agents to prevent or treat diseases. Ginger has an excellent antimicrobial activity against various gram positive and gram negative bacteria and fungi. In vitro studies have shown that active constituents of ginger inhibit multiplication of colon bacteria. These bacteria ferment undigested carbohydrates causing flatulence. This can be counteracted with ginger (Gupta & Ravishankar, 2005) [18]. The extract of ginger inhibits the growth of *Escherichia coli*, *Proteus vulgaris*, *Staphylococcus aureus*, *Streptococcus pyogenes* and *Salmonella* (Ernst and Pittler, 2000; White, 2007) [12, 47]. This plant can be used as one of the best medicinal plants is controlling pathogenic bacteria.

1.6. Ginger as an immunostimulant

In aquaculture, the application of dietary medicinal herbs as immunostimulants can elevate the innate defense mechanisms of fish against pathogens during periods of stress, such as, intensive farming practices, grading, sea transfer, vaccination and reproduction. The excess use of antibiotics in the

management of disease in aquaculture has resulted in serious health and environmental problems. Consequently, the need of safe and effective alternatives to antibiotics is required. In this context, immunostimulants have attracted significant attention. Ginger as a natural antibiotic is one of the most effective natural immunostimulants. The powdered ginger rhizome is able to enhance non-specific immune response in rainbow trouts (Haghighi and Rohan, 2013) ^[19]. Non-specific immunity plays an especially important role in the defense of fish and is the sole immunological mechanism by which invertebrates protect themselves from diseases (Secombes and Olivier, 1997) ^[40]. Non-specific defense mechanism plays an important role in all stages of fish infection. Fish particularly depend mostly on these non-specific mechanisms than mammals do (Avtalion, 1981) ^[5]. Ginger essential oil showed improvement in humoral and cell mediated immune response in immune-suppressed mice (Carrasco, *et al.*, 2009) ^[7]. However, future studies might look into the dose-response, determination of optimal dose, treatment duration, and its use in large scales in fish farms. In general, this study suggests that ginger can be applied as an alternative diet and a supplement to boost immune system for rainbow trout.

Ginger is effective as an immunomodulatory agent in animals and fish and helps to reduce the losses caused by diseases in aquaculture (Nya and Austin, 2009; Ali *et al.*, 2008; Zhou *et al.*, 2006; Tan and Vanitha, 2004) ^[37, 48, 44]. Rainbow trout on powdered ginger rhizome diet for 12 weeks showed increased haematocrit, haemoglobin, erythrocyte, MCH, MCHC, WBC values and neutrophils percentage in comparison to the control group ($p < 0.05$). De Pedro *et al.*, (2005) ^[10] indicated that total and differential leukocyte counts are important indices of non-specific defense activities in fish. Also, they are centrally involved in phagocytic and immune responses to bacterial, viral and parasitic challenges (Houston, 1990) ^[23].

1.7. Traditional use

Ginger is known as *Sunthi* in Ayurveda and description of the plant appears in the old text like *Charaka*, *Sushruta*, *Vagbhata* and *Chakra-dutta* (Malhotra & Singh, 2003) ^[32]. The use of drug is mentioned in form of *Trikatu*, a famous Ayurvedic remedy for the treatment of digestive disorders. In *Ashtanga Hridaya*, the plant has been used in *Rasna Saptak Quath* (a decoction based on seven medicinal herbs), and a traditional remedy of arthritis. Pharmacologically, the drug in Ayurveda has been described as appetizer. It is also indicated in ointment form for local application in pains.

1.8. Phytochemistry

Ginger is a rich source of volatile oil. Zingiberol, zingiberene, phellandrene and linalool are important constituents of the oil. They account for the aroma of the drug. The pungency of the ginger is due to gingerols and shogaols. Investigations have shown gingerol and shogaols to be mutagenic. In addition, ginger contains a special group of compounds called diarylheptanoids including gingerenone (Kikuzaki *et al.*, 1991) ^[28]. The standardization of the drug is based on presence of pungent principles of the plant.

2. Conclusion

Medicinal herbs are rich source of synthetic and herbal drugs. They contain a wide range of chemical compounds commonly referred to as phytochemicals. Ginger is a hot herb today and number of studies has shown it to be a useful medicinal agent. Its potential as an effective anti-inflammatory and anti-emetic agent cannot be ruled out. *Zingiber officinale*, ginger, is an

important plant with several medicinal, ethno-medicinal and nutritional values used in traditional medicine. Ginger is consumed worldwide as a spice and flavoring agent and is attributed to have many medicinal properties such as cardio-protective, anti-inflammatory, anti-microbial, anti-oxidant, anti-cancer properties, etc. The ginger is used as growth promoter, antimicrobial agent, antioxidant and as immunostimulant in aquaculture. Gingerol, the active constituent of ginger has been isolated and studied for pharmacological and toxic effects. Large-scale clinical studies are required to justify ginger as suitable phytopharmaceutical drug although initial data seems to be promising.

3. Acknowledgements

The Author is highly grateful to Prof. Dr. Shyam Narayan Labh, President of Nepal Aquaculture Society and Professor of Zoology, T. U., Nepal for his cooperation and help while writing this review article.

4. References

1. Agrawal M, Walia S, Dhingra S, Khambay BPS. Insect growth inhibition antifeedant and antifungal activity of compounds isolated derived from *Zingiber officinale* Roscoe, ginger rhizome. *Pest Manag Sci.* 2001; 57:289-300.
2. Ali BH, Blunden G, Tanira MO, Nemmar A. Some phytochemical, pharmacological and toxicological properties of ginger *Zingiber officinalis* Roscoe: a review of recent research. *Food Chem Tox.* 2007; 46:409-20.
3. Apines-Amar MJS, Amar EC, Faisan JP. Jr., Pakingking, R. V. Jr., & Satoh, S. Dietary onion and ginger enhance growth, hemato-immunological responses, and disease resistance in brown marbled grouper, *Epinephelus fuscoguttatus*. *AACL Bioflux.* 2012; 5(4):231-239
4. Austin B, Austin DA. *Bacterial Fish Pathogens In: Disease in Farmed and Wild Fish*, 4th Ed. Springer-Praxis, Godalming. 2007, 57-63.
5. Avtalion RR. Environmental control of the immune response in fish. In: Clem LW, editor. *CRC critical reviews in environmental control*. London: CRC Press, 1981, 163-88.
6. Bartley J, Jacobs A. Effects of drying on flavour compounds in Australian-grown ginger (*Zingiberofficinale*). *Journal of the Science of Food and Agriculture.* 2000; 80:209-215.
7. Carrasco FR, Schmidt G, Romero AL, Sartoretto JL, Caparroz-Assef SM, Bersani-Amado C A *et al.* Immunomodulatory activity of *Zingiber officinale* Roscoe, *Salvia officinalis* L. and *Syzygiumaromaticum* L. essential oils: evidence for humor- and cell-mediated responses. *J Pharm Pharmaco.* 2009; 1(7):961-967.
8. Chen Ch, Kuo M, Wu Ch, Ho Ch. Pungent compounds of ginger (*Zingiber officinale* (L) Rosc) extracted by liquid carbon dioxide. *Journal of Agricultural and Food Chemistry.* 1986; 34:477-480.
9. Chrubasik S, Pittler MH, Roufogalis BD. *Zingiberis rhizoma*: a comprehensive review on the ginger effect and efficacy profiles. *Phytomedicine.* 2005; 12:684-701.
10. Pedro De N, Guijarro AI, Lopez-Patino MA, Martinez-Alvarez MJ, Delgado R. Daily and seasonal variations in hematological and blood biochemical parameters in the tench, *TincatincaLinnaeus*, 1758. *Aquacult Res.* 2005; 36:1185-1196.
11. Endo K, Kanno E, Oshima Y. Structures of antifungal diarylheptenons, gingerones A, B, C and

- isogingerenones, isolated from the rhizomes of *Zingiber officinale*, *Phytochemistry*. 1990; 29:797-799.
12. Ernst E, Pittler MH, Efficacy of ginger for nausea and vomiting: A systematic review of randomized clinical trials. *Br. J Anaesth*. 2000; 84:367-371.
 13. Esiobu N, Armenta L, Ike J, Antibiotic resistance in soil and water environments. In: *J Environ. Health Res*. 2002; 12:133-144.
 14. Food and agricultural Organization of United Nations: Economic and social Department: The statistical Division, 2012.
 15. Govindarajan V. Ginger-chemistry technology and quality evaluation: Part-I CRC. *Critical Reviews in Food Science and Nutrition*. 1982; 17:1-96.
 16. Grzannar R, Lindmark L, Frondoza GG. Ginger -An herbal medicinal product with broad anti-inflammatory actions. *J Med Food*. 2005; 8:125-32.
 17. Gughani HC, Ezenwanze EC. Antibacterial activity of extracts of ginger (*Zingiber officinale*) and African oil bean seed (*Pentaclethra macrophylla*). *J Commun Dis*. 1985; 17:233.
 18. Gupta S, Ravishankar S. A comparison of the antimicrobial activity of garlic, ginger, carrot, and turmeric pastes against *Escherichia coli* O157:H7 in laboratory buffer and ground beef. *Food borne Pathogen Dis*. 2005; 2(4):330-40.
 19. Haghighi M, Rohani MS. The effects of powdered ginger (*Zingiber officinale*) on the haematological and immunological parameters of rainbow trout *Oncorhynchus mykiss*. *Journal of medicinal Plant and Herbal therapy research*. 2013; 1:8-12.
 20. Haksar A, Sharma A, Chawla R, Kumar R, Arora R, Singh S *et al*. *Zingiber officinale* exhibits behavioral radio protection against radiation-induced CTA in a gender-specific manner. *Pharmacology Biochemistry and behavior*. 2006; 84(2):179-88.
 21. Harikrishnan R, Balasundaram C, Heo MS. Impact of plant products on innate and adaptive immune system of cultured finfish and shellfish. *Aquaculture*. 2011; 317:1-15.
 22. Hoffman T. Antimicrobial activity of some medicinal plants from India. *Hawaii Medical Journal*. 2007; 66:326-327.
 23. Houston AH. Blood and Circulation. In: *Methods for Fish Biology*, Schreck. C.B and P.B. Moyle (Eds.). American Fisheries Society, USA. ISBN: 0-913235-58-X, 1990, 273-334.
 24. Ibrahim MD, Fathi M, Mesalhy S, Abd El-Aty AM. Effect of dietary supplementation of insulin and vitamin C on the growth, haematology, innate immunity, and resistance of Nile tilapia (*Oreochromis niloticus*). *Fish Shellfish Immunol*. 2010; 29:241- 6.
 25. Immanuel G, Uma RP, Iyapparaj P, Citarasu T, Peter SM, Babu MM. Dietary medicinal plant extracts improve growth, immune activity and survival of tilapia *Oreochromismossambicus*. *J Fish Biol*. 2009; 74:1462-75.
 26. Kadnur SV, Goyal RK. Reneficial effects of *Zingiber officinale* Roscoe on fructose induced hyperlipidemia and hyperinsulinemia in rats. *Indian J ExpBiol*. 2005; 43:1161-4.
 27. Kapoor A. Antifungal activities of fresh juice and aqueous extracts of turmeric and ginger (*Zingiber officinale*). *J Phytological Res*. 1999; 10:59.
 28. Kikuzaki H, Kobayashi H, Nakatani N. Constituents of Zingiberaceae, Diarylheptanoids from rhizomes of *Zingiber officinale*, *Phytochemistry*. 1991; 30:3947-3952.
 29. Khaki A, Fathiazad F, Nouri M, Khaki A, Chelar C, Ozanci *et al*. The Effects of Ginger on Spermatogenesis and Sperm parameters of Rat. *Iranian J Reprod Med*. 2009; 7:7-12.
 30. Kikuzaki H, Nakatani N. Antioxidant effect of some ginger constituents. *J Food Sci*. 1993; 58:1407-10.
 31. Kim JK, Kim Y, Na KM, Surh YJ, Kim TY. [6]-Gingerol prevents UVB-induced ROS production and COX-2 expression in vitro and in vivo. *Free Radic Res*. 2007; 41:603-614.
 32. Malhotra S, Singh PS. Medicinal properties of Ginger (*Zingiber officinale*), *Natural product Radiance*. 2003, 2(6).
 33. Maqsood S, Singh P, Samoon MH, Munir K. Emerging role of immunostimulants in combating the disease outbreak in aquaculture. *International Aquatic Research*. 2011; 3:147-163.
 34. Martins AP, Salgueiro L, Goncalves MJ, Proenca da, Cunha A, Vila R *et al*. Essential oil composition and antimicrobial activity of three Zingiberaceae from S. Tome Principe. *Planta Medicine*. 2001; 67:580-584.
 35. Nicoll R, Henein MY. Ginger (*Zingiber officinale* Roscoe): a hot remedy for cardiovascular disease? *Int. J Cardiol*. 2009; 131:408-409.
 36. Nielsen PV, Rios R. Inhibition of fungal growth on bread by volatile compounds from spices and herbs and mustard essential oil. *Inter J Food Microbiol*. 2000; 60:219-229.
 37. Nya EJ, Austin B. Use of dietary ginger, *Zingiber officinale* Roscoe as an immunostimulant to control *Aeromonas hydrophila* infections in rainbow trout, *Oncorhynchusmykiss* (walbaum). *Journal of Fish Diseases*. 2009; 32:971-977.
 38. Panigrahi A, Azad IS. Microbial intervention for better fish health in aquaculture: the Indian scenario. *Fish Physiol. Biochem*. 2007; 33:429-40.
 39. Robertsen B. Modulation of the non-specific defence of fish by structurally conserved microbial polymers. *Fish Shellfish Immunol*. 1999; 9:269-290.
 40. Secombes CJ, Olivier G. *Furunculosis*. Academic Press, New York, 1997, 269-296.
 41. Sharma S, Vijayvergia R, Singh T. Evaluation of antimicrobial efficacy of some medicinal plants. *J Chem. Pharm. Res*. 2010; 2(1):121-124.
 42. Stoilova I, Krastanov A, Stoyanova A, Denev P, Gargova S. Antioxidant activity of a ginger extract (*Zingiber officinale*), *Food Chemistry*. 2007; 102:764-770.
 43. Talpur AD, Ikhwanuddin M, Abol-Munafi AB. Nutritional effects of ginger (*Zingiber officinale* Roscoe) on immune response of Asian sea bass, *Lates calcarifer* (Bloch) and disease resistance against *Vibrio harveyi*. *Aquaculture*. 2013; 400:46-52.
 44. Tan BKH, Vanitha J. Immunomodulatory and antimicrobial effects of some traditional Chinese medicinal herbs: A review. *Curr. Med. Chem*. 2004; 11:1423-1430.
 45. Venkataramalingam K, Godwin Christopher J, Citarasu T. *Zingiber officinalis*, an herbal appetizer in the tiger shrimp *Penaeusmonodon* (Fabricius) larviculture. *AquacultNutr*. 2007; 13:439-43.
 46. Weidner MS, Sigwart K. The Safety of a Ginger Extract in the Rat. *J Ethnopharmacol*. 2000; 73(3):513-520.

47. White B. Antimicrobial activity of ginger against different microorganisms: *Physician*. 2007; 75:1689-1691.
48. Zhou HL, Deng YM, Xie QM. The modulatory effects of the volatile oil of ginger on the cellular immune response in vitro and in vivo in mice. *J Ethnopharmacol*. 2006; 105:301-305.